izing thoroughly with a 10 per cent. solution, I noted no change in the dilated pupil, but this did not influence me in any way, as the cocainization had little or no effect in diminishing the hypertrophy. There was nothing of moment in the operation. The day after, the pupillary dilation was greatly reduced; two or three days afterward the dilation was as great as ever, but I could justly account for this, by the fact that the pressure of the scab covering the point of operation was as great as that of the original hypertrophy; I did not take this away at once, but waited two or more days and then removed it, after which the pupillary reaction gradually became normal.

Allow me to add that in this case I found a hysterical deafness, that is to say, the watch, speech and tuning-fork tests gave normal reaction; but, as she said, "while she could hear what was said, she could not always understand, and was obliged to have sentences repeated." I would like to say too, that I was not sure that the pressure caused by the hypertrophy was the cause of the pupillary dilation, but there were ample indications to justify me in the removal of the pressure. The results were gratifying.

Case 2.—Miss B., a nurse, aged 27, was also of a rather nervous temperament, but not so markedly as in the first instance; the left pupil here was dilated to 7 mm. The right measured but 4 mm in diameter. On examining the left nostril, a diffused redness of the mucous membrane was apparent. There was no hypertrophy, but the anterior head of the middle turbinated body was so bent as to produce marked pressure against the septum. Having the experience of the previous case before me, I determined to remove as much of the turbinated as caused the pressure, which I did by means of cutting forceps; the results in this case were not so satisfactory as in the first, as the pupillary dilation diminished only gradually, and the last time I saw the patient, while it was greatly diminished in size, the left pupil was slightly larger than the right.

Dr. Bettmann's Notes:

Case 1.—Mrs. P. came to me in May, 1892, complaining of inability to read the markings of the thermometer. Upon examination I found the following conditions: Vision, both eyes equal to 20-20, accepted + 0.5D both eyes. Right pupil dilated (medium); left pupil one-half again as large. No reaction to light or on convergence. Can not read ordinary newspaper print. With $\times 1.5D$ reads fine print at twelve inches; muscular equilibrium normal; ophthalmic examination negative. Diagnosis: paralysis of sphincter muscles of irides and accommodation. No history of diphtheria, rheumatism, or syphilis. Recommended galvanic current and injections of strychnin. This treatment was carried out for two weeks, without producing any improvement in her condition. I then referred her to Dr. Pierce, who operated with instantaneous effect on the ocular symptoms. The day after the operation the pupils were normal in size and reaction and she read the finest type at close range. A few days later—as soon as a crust formed over the wound—the abnormal conditions of the eye returned, to disappear again permanently as soon as the scab came away.

The changes in Case 2, Miss B., were similar to those already reported; the reaction to light very slight on convergence. Vision = 6-12, with $\times 1D$. V. 6-6 both eyes. Can only read fine print with her glasses. The ophthalmoscope revealed nothing abnormal. Advised the patient to consult Dr. Pierce for examination of the nose. Dr. Pierce operated, and the operation was followed a week later on exposure to cold by an acute catarrhal otitis media and a mastoid-periostitis, both of which yielded readily to treatment. The ocular symptoms had previously disappeared.

A third case I wish to report, was a patient referred to me by Dr. Fenger of Chicago. A farmer from the interior of Illinois; strong-framed, muscular, plethoric. Complained of a constant pain in the right half of the tongue. He consulted Dr. Fenger, fearing cancer, who, not finding any pathologic change in the tongue referred the patient to me. On examination, I found an extensive hypertrophy of the inferior turbinated body on the right side, causing complete obstruction and a marked degree of pressure in that region. Operation was performed by means of galvano-cautery. In three days the pain in the tongue had entirely disappeared, and has remained absent ever since—a year having elapsed since the operation was performed.

REPRODUCTION OF THE UPPER AIR-PASSAGES BY PLATING CASTS OBTAINED BY THE CORROSION METHOD.

Read in the Section on Laryngology and Otology, at the Forty-fourth Annual Meeting of the American Medical Association.

BY B. ALEXANDER RANDALL, M.D.

PHILADELPHIA, PA.

While it is no new thing to obtain, in metal or other materials, casts of the cavities or structures of the body, and exquisite results have been long and by many workers obtained in this field, there yet remains much to be thus demonstrated to the scientific world at large, and far more in the way of local demonstration. It is with this latter object that I desire here to bring forward some of my results, believing that they will be new to some and interesting to others who work in the upper air tract. Made in fusible metal, which can be readily cast into the soft tissues of the recent cadaver, these metal casts have a delicacy and a durability far superior to anything which is possible with the more frequently used wax mass; and the secondary process which I have here employed has marked advantages and seems to constitute a new and important step in this matter.

All of these casts have the grave disadvantage, which grows with the increased complexity of the object, of being negatives only, and a comprehension of their real teaching can be gained only by a process of mentally turning them inside out, which many persons find very difficult and unprofitable. They are often needlessly costly, too, since one can lock up many pounds of expensive alloy in a single cast. But if the solid cast can be electroplated and its metal then fused out to leave the thin shell of deposit, we have a repetition of the negative almost as true and quite as instructive as before; while we have absolutely reproduced, in a permanent and convenient form, the hollows of the organs originally molded. Strong as is the fusible metal, it is yet so heavy that casts made of it are very apt to fracture if dropped, and my idea, as a teacher, is that preparations too delicate to take some chances of rudeness of handling are of little value except to the original student. They may be marvelous of beauty and accuracy, full of new and instructive points, but all of this is lost to those who can not handle and closely study them. These electrotypes can be featherweight reproductions if desired—delicate as the empty shells of a collection of birds' eggs—yet safe for any reasonable handling or transportation. Carefully sectioned, before or after the removal of the fusible metal within, they can afford new possibilities of study and reveal much which might escape observation, even when both interior and exterior are thus accessible.

The first results which I have now in hand to bring forward are in many respects rude and unsatisfactory; yet are delicate enough to reveal all the shortcomings of the original casts. They illustrate that the primary casting in fusible metal should be done with all care and accuracy, and with due consideration of the aims, difficulties and possibilities of the method. The metal had best be fused in a water-jacketed funnel capable of containing several pounds, so that a superabundance may be present to secure full fluidity of the material and completeness at one casting without requiring any supplementary

pouring, even if a good deal should be lost through some unexpected leakage. This last mishap calls for the aid of some efficient assistant who shall stand ready to stem such flow, and do anything to aid in perfecting the result. Thus cast, with foresight in arranging for the escape of air from all the cavities and a normal disposition of the preparation so as to avoid distortion, exquisite results can be obtained from adequate material; the main difficulty in America, being to obtain such material in any quantity, and at times when the experimenter really has the leisure to utilize it. This has rarely been my good fortune, hence my preparations fall far short of what the method might afford.

As to the electroplating, the main point is that it should be slowly done, in order to obtain due evenness and smoothness. Any forcing of the process will give lumpy distortions which may greatly impair the accuracy and beauty of the operation.

BICHLORID OF MERCURY IN THE TREATMENT OF DIPHTHERIA.

Read in the Section on Laryngology and Otology, at the Forty-fourth Annual Meeting of the American Medical Association.

BY CHARLES F. MCGAHAN, M.D.

AIKEN, S. C.

I wish to call attention to the treatment of diphtheria by the bichlorid of mercury, not that I regard it as a specific in the disease, for I am sure that all of us who have had much experience with this most treacherous malady will agree with me that there is no such thing as a specific for it. Various modes of treatment have been followed by me, and frequently with such success in different epidemics that I felt on several occasions I had found a cure, but each time my hopes were crushed by having the treatment fail entirely in the next series of cases which fell under my observation. I have made this statement so that all can see that I am not reviving this treatment, in the expectation of having it considered a "cure-all" for the disease, but simply because I believe that if the profession would use it more extensively, and in larger doses, that we should have better results.

In one of the latest works on general therapeutics we find, in laryngeal diphtheria, "one one-hundred and fiftieth to one one-hundredth of a grain of bichlorid may be given every hour, for not over four days, and it must be carefully watched." This dose is not large enough to be of any service, if we have a true case of diphtheria to deal with. In fact, we can diagnose a case of diphtheria by the way the patient bears the bichlorid of mercury. I will give the history of one case, so as to illustrate the manner of giving it:

On February 13 I was called to see a patient, a lady over sixty years of age, who was suffering from a sore throat. Upon examination, I found the fauces very red, and not then suspecting diphtheria I prescribed peroxid of hydrogen (Marchand's) one to four, to be sprayed four times a day, and an alterative. When calling on the 14th, I found both tonsils covered with membrane. The patient was immediately put upon one-fifteenth of a grain of bichlorid of mercury every hour and a half, and to be sprayed every hour, alternating with bichlorid of mercury one one-thousandth and peroxid of hydrogen one to two. The disease gained rapidly and by the next

day the pharynx, larynx and nasal passages were covered with membrane. For the first four days the patient took one-fifteenth of a grain of bichlorid of mercury every hour and a half, and for the four following days she took one-fifteenth of a grain of bichlorid of mercury every two hours. Besides this, during the eight days her throat was sprayed every alternate hour with bichlorid of mercury one one-thousandth, and enough solution was atomized each time to vaporize one-tenth of a grain of bichlorid of mercury, making one and one-half grains of bichlorid of mercury sprayed on the throat every day, and some days it amounted to even more, for when the disease was at its worst I used the bichlorid of mercury every hour.

In spraying, the larynx, pharynx and nasal cavities were sprayed. During the eight days over eight grains of bichlorid of mercury were given by the mouth, and about twelve grains sprayed into the larynx, pharynx and nasal cavities.

The only bad effect I had from the bichlorid was great depression which was counteracted by giving whisky freely. The throat was so covered with the membrane that my patient could not swallow for thirty-six hours, and nourishment was given by the rectum, and bichlorid through a tube passed in through the nose. Her recovery was very slow, and she suffered with paralysis of the vocal cords for over two months, but it gradually yielded to treatment.

In conclusion, I wish to impress you with the importance of giving bichlorid of mercury in large doses, in diphtheria, and do not be afraid of it.

CLINICAL EXPERIENCE WITH SOLUTIONS OF PYROZONE.

Read in the Section on Laryngology at the Forty-fourth Annual Meeting of the American Medical Association.

BY JAMES E. NEWCOMB, M.D.

ATTENDING LARYNGOLOGIST DEMILT DISPENSARY; ASSISTANT IN THE ROOSEVELT HOSPITAL, OUT-PATIENT (THROAT) DEPARTMENT, NEW YORK, N. Y.

About thirty-five years ago, Sir Benjamin Ward Richardson introduced into medical and surgical practice the use of solutions of hydrogen dioxid. For a quarter of a century or more, he was their only advocate. With considerable distinctness he pointed out their application to various clinical conditions. He carefully recognized their limitations and, while making for them broad claims, did not regard them as a panacea for all ills. During the past few years, he has had many imitators of his advocacy. They have gone far beyond him in their claims of therapeutic results. They have, unfortunately, fallen short of him in honest conservatism. They have forgotten the counsels of this therapeutic pioneer, have made wild and untrustworthy statements, and have thus tended to bring the remedy into disrepute. They have had as allies some of the manufacturing chemists, whose wily rhetoric has been far more specious than their appreciation of the problems of clinical medicine has been accurate.

Richardson strongly insisted that it was impossible to prepare more than a ten-volume stable solution in water; to exceed this limit required a degree of acidity so great that, on exposure to the air, the solution rapidly ran down in oxygen strength, the H_2O_2 becoming practically H_2O . A true and honest ten-volume solution has only a very slight degree of