most part in the second week, the temperature of which was 2° below its average; that of the preceding week having been 6° below its average.

November.—Barometer, 29.642 in.; being $\frac{1}{4}$ of an inch below its average. Temperature was 48.9° ; being $6\frac{1}{2}^{\circ}$ too great. Degree of humidity was 83; being 6 too small. The daily range of temperature was 10.4° ; being 0.3° too small. The fall of rain during the month was 6 in.; which was 3.7 in. above the average. The number of deaths daily was 138, or 4129 in the month; the expected number being 4912. The mortality during the first two weeks of the month was above its average, but on the third week it was much less.

December.—Barometer, 29.758 in.; being 0.282 in. below the average. The temperature was 47.6; exceeding its average value by 8.8° . The degree of humidity was 79; being 10 below its average. The daily range of temperature was 9.7° ; exceeding its average by 0.8° . Rain fell to the depth of 2.2 in. The number of deaths daily was 148, or 4597 during the month; while the corrected average was 5791—1194 less than the average. The temperature of the months November and December was very high, the pressure was low, the degree of humidity was low, and the fall of rain was in excess.

The number of deaths in the year was 54,022; being about 4000 less than the calculated number.

The mean temperature of the year was 50.6° ; being 2.2° above the average.

The mean reading of the barometer for the year was 29.905 in.; being 0.046 in. below its average.

Rain fell to the depth of 34.4 inches.

REMEDY FOR INFANTILE ASPHYXIA.

BY W. P. JONES, M.D., NASHVILLE, TENN.

ONE pleasant afternoon, not long ago, I was sitting quietly in my office, when my cogitations were interrupted by the hasty steps of a large, fresh, Yankee-looking gentleman, who said his wife was suffering very much, and wanted a doctor. I followed him, and on the way was informed by the husband of the sufferer that he had only been married a few days—an interesting patient, thought I, and probably quickened my steps. On arriving at the house, I was conducted into the room, where I soon became apprehensive that the woman, though just married, was absolutely in childbirth, and that my services as accoucheur would be brought into immediate requisition. After making the usual examination, and becoming thoroughly satisfied that my conjecture was well founded, and that it was the first presentation, I sat down to await the issue, and was not long detained; for immediately very severe pain supervened, and a foetus was expelled; without, however, any other vital indication than bodily warmth, and very feeble, almost indistinct, circulation in the cord.

The ordinary means, such as stimulating baths, inflating the lungs, friction, &c., were used, probably a half hour or more, without any ap-

parently beneficial effect. At this crisis, the father (or man who should have been), entered the room, bearing upon his breath the odor of alcohol. I called to him, and told him to come and apply his lips to those of the child, and gently blow into its mouth. He looked astonished, and exclaimed—What, me, sir! Yes, sir, said I, you are just the man, come on. He did so. He inflated the lungs, and I expelled the air repeatedly; and together we at length succeeded in resuscitating the child, which I am constrained to regard as wholly attributable to the stimulus of the alcoholic inspiration. It is at any rate worthy of a trial, after the ordinary means have failed; and in the absence of a drunken husband, we suggest that the accoucheur inhale the vapor of alcohol, or spirits of camphor, immediately previous to inflating the lungs of the child.—*Southern Journal of the Medical and Physical Sciences*.

STRICTURE OF ŒSOPHAGUS.

BY WILLIAM JOHNSON, M.D.

AMONG the many ills to which flesh is heir, strictures of those canals which are lined by the mucous membrane, are far from being the least interesting, whether considered with respect to the sufferings both mental and physical, which they produce, or the imminent danger to which they often subject the unfortunate individuals, who are the victims of their visitation. Memory recalls many cases illustrative of these truths, particularly two fatal cases of stricture of the *œsophagus*, which occurred in this region of country since my settlement here. The first case I did not myself see, but had the relation from his widow. He died a few weeks after my settlement. He was young, but absolutely starved to death, after protracted sufferings, in full consciousness of his dreadful condition. The other case I saw but once, and that a short period previous to her death. She, too, died from inanition. Prof. Dorsey, in his *Surgery*, relates a case of stricture of the *œsophagus*, where the post-mortem examination revealed an almost entire obliteration of the canal; a probe could scarcely pass through the strictured part. Velpeau and other writers cite cases of complete obliteration of the passage.

It is not my purpose to enter into speculation, as to the causes of these affections, whether they be owing to ulceration in some portion of these canals, producing narrowing by the contraction of their cicatrices in healing, or congestion and infiltration in the adjacent tissues while the ulcerative process is still going on, or whether simple spasmodic action be prominent in their production. But any light which can be thrown upon this subject by the record of individual cases, is due to the medical public, and it is under an imperative sense of duty that I am induced to present the following case of stricture of the *œsophagus*.

I was consulted, the 23d of last July, by M—— C——, aged about 22 years, on account of stricture of the *œsophagus*. Her attention was called to it, about two years since. She was eating a piece of pickled cucumber, and was choked, and still more so very shortly afterwards, in eating a pear. Since then the disease has been gradually increasing.