20 years. Among chimney-sweeps the comparative mortality is above the standard by about one-third part. By far the greatest excess of mortality in this occupation is attributable to cancer, for which the comparative figure is $2\frac{1}{2}$ per cent above the standard. Chimney-sweeps appear to be exceptionally addicted to alcoholism, and their mortality from phthisis, diseases of the nervous, digestive, and respiratory sys-

tems and from suicide, is also high. On the other hand they appear to be but slightly liable to influenza, Bright's disease, and to accident. The comparative mortality of Civil Service messengers is 21 per cent below the standard, and that of gamekeepers is remarkably low, being in the main working period of life 42 per cent less than the standard.

Taking the general body of occupied males between

the ages of 25 and 65 we find the mortality lowest among clergymen, priests and ministers, then come gardeners, nurserymen, seedsmen, gamekeepers, farmers, graziers, railway engine-drivers, stokers, firemen, laborers, the highest mortality being among innkeepers, publicans, costermongers, tin miners, and general laborers.—Journal of the Royal Society of Arts.

THE PREPARATION OF TOBACCO.*

ITS CULTIVATION AND ITS CHEMISTRY.

BY DR. W. DONSELT.

TOBACCO, an annual plant, of the order of *Solanaceæ*, is indigenous to tropical America, and is cultivated in America, Europe, and Asia. As soon as it is ripe, it is slowly dried with the greatest care in specially constructed buildings, so-called drying barns, until it becomes "barn-ripe." As a general rule, in Germany, the leaves are strung separately on cords and hung up; in America the entire plant is cut off just above the root and suspended upside down in the drying place. The leaves remain alive under this treatment, when hung in the drying barns, and processes occur in them that are essential to the production of a perfectly barn-ripe article. Thus, for instance, the plant consumes the starch by respiration, the albuminous substances are decomposed, and the quantity of amides increased; the color of the tobacco, at first green, gradually changes to brown.

Before the drying proper, the tobacco is often allowed to sweat, and to accomplish this the leaves are piled in heaps on one another. Soon after this, heating takes place in the interior of the heaps, due to the fact that the still living leaves begin to breathe. By this mode of treatment it is claimed that the tobacco leaf attains a better ripeness and dries better. Above all things, however, the temperature of the heap must be carefully watched, for if the leaves get too hot they stew and turn black.

In the drying proper, micro-organisms play no part. But if the drying is not carefully enough conducted, e. g., if the particularly thick ribbed leaves, which are slowest in drying, are not placed far enough apart, so that the air and natural heat will have free access to them, the leaves very rapidly spoil. As Müller-Thurgan found, it is a fungus of the species *Pleospora* that attacks the leaves and causes the so-called "barn blight." We may in such case note the gradual appearance on the leaves of dark colored to black spots, growing gradually larger, which, with sufficient moisture, may even cover the entire leaf. The leaf is soft and moist, and tears at the slightest touch. Sturgis found, in cases of rotting of this character, two different types of bacteria, a rod bacterium and a micrococcus, living together on the leaves.

These diseases attack only the tobacco leaves; the stems of the entire plants hung up to dry are attacked by the so-called stem-rot. On the stems white, velvet-like spots appear, which gradually extend farther along the leafribs. Behrens, who investigated these spots, found that they consisted of the mycelium of *Botrytis longibrachiata* Oudem. This fungus gradually destroys the entire tissue of the leaf.

In addition, the quality of the tobacco is disadvantageously affected by the artificial fertilization of the fields with Chili saltpeter. This promotes, for instance, the occurrence of stem rot and "barn blight." The tobacco leaf is certainly made thicker and fleshier by resort to saltpeter fertilization, and in consequence contains more water. This prolongs the drying process in the sheds and affords the fungi of putrefaction an opportunity to establish themselves. In addition, the substance of the tobacco by this method of fertilization becomes a peculiarly favorable field for *Botrytis*.

The tobacco being now barn-ripe, passes into the hands of the manufacturer or dealer, who then subjects it to fermentation. The tobacco bunches are piled together in great heaps, each containing many hundred-weight, upon which the entire mass begins gradually to become heated. This spontaneous heating, due to fermentation, will advance in the interior of the heap to a temperature of 142 deg. F., as Suchsland found, but such a high degree of heat should be avoided; it should not be allowed to exceed 122 deg. F. Should it occur, the heaps must be taken apart, the outside leaves placed in the center and those in the center on the outside of the heap. Work is continued in this manner until the entire mass appears to be sufficiently fermented.

During the fermentation of the tobacco leaves, which proceeds under the generation of considerable spontan-

eous heat, a fairly considerable loss of substance occurs. Thus, in fermented raw tobacco, there is a lack of carbohydrates soluble in water, like sugar and the organic acids, for instance the malic and citric acids still found in barn-ripe leaves, whereas the amount of starch remains unchanged during the fermentation. In addition, the albuminous substances are decomposed, amides being developed; a decrease in the amount of nicotine is also said to be caused thereby. This, however, can occur only as a result of the activity of fermentation organisms, for in tobacco nicotine is not found in a free state, neither in the unfermented nor fermented leaf, but occurs in the form of non-volatile compounds. The nicotine can therefore not be volatilized by heat during fermentation, as confirmed by nitrogen determinations before and after fermentation. The amount of nitrogen, if we take into consideration the loss of dry substances during fermentation, was the same. The micro-organisms that are active in the fermentation use probably a portion of the nicotine as nutriment and transform it into other combinations. Behrens found, for instance, that the mold fungus *Botrytis cinerea* can supply its requirements of nitrogen from the nicotine, just as bacteria placed in a nicotine nutritive solution gradually effect its complete elimination.

As regards the volatile acids in tobacco, it was found that after fermentation they had doubled in quantity. Lactic acid could not be discovered in fermented tobacco; on the other hand, butyric acid was probably formed.

In regard to the organism germs found in the fermentation the bacteriological investigations of Vernhout are worthy of special notice. He used for his observations East Java tobacco. During its fermentation he first of all directed his attention to the thermophile bacteria and he consequently kept the cultures constantly at 122 deg. F. At this they contained invariably the same kind of bacillus. This was actively mobile and formed endospores belonging to the group of the potato bacillus, and it grew likewise at ordinary temperature. In addition, it did not grow as well in a nutritive medium containing tobacco decoction as in one free from tobacco, and in bouillon developed ammonia. Only later Vernhout discovered that the tobacco used by him in the preparation of the tobacco decoction he employed had been subjected to a fermentation, and the experiments which he subsequently instituted with a decoction of barn-ripe tobacco showed that the above-mentioned thermophile bacillus, *Bacillus tabaci fermentationis*, grew well in it. The temperature optimum for growing and flourishing was between 111 deg. and 122 deg. F. The temperature limits were between 77 deg. and 136 deg. F.

Vernhout essayed in the following manner to inoculate a pure culture of the *Bacillus tabaci fermentationis* on tobacco leaves: He halved the tobacco leaves, which were but slightly fermented, on the line of the central rib and then, after further reduction, distributed the halves in Petri dishes in such a manner that the right half of the leaves came into one, the left half into the other part of the double dish. After the halves had been sterilized at 248 deg. F., they were inoculated with pure culture of the *Bacillus tabaci fermentationis* and all the dishes left in a thermostat at 122 deg. F. Of seven experiments only three furnished favorable results and, as a fact, in one of these experiments the tobacco leaf halves, after about twelve days, developed a marked odor of sweet rye-bread, a flavor that is almost invariably noticed in tobacco fermentation. Unfortunately Vernhout left Java too soon, so that the investigations could not be continued.

Another investigator, Koning, observed the fermentation of Dutch tobacco. He found invariably numerous distributed in fermenting tobacco *Bacillus subtilis*, as well as mycoides, also the *Bacillus tabaci*, in three different types. Of these two belonged to the Proteus group, the other, the only spore-forming variety, to the Subtilis group, which played the chief part in the fermentation. Koning also, like Vernhout, inoculated sterilized tobacco with pure cultures of these

different organisms and then allowed them to ferment for six weeks. After this period, he determined that the tobacco inoculated with the *Bacillus tabaci*, belonging to the Subtilis group, was the best, which was also confirmed by practical experts.

In addition, Koning also isolated a *Diplococcus tabaci*, which took part in the fermentation and the optimum of growth of which was 75 deg. F. At a higher temperature of the made-up tobacco heaps, the activity of the diplococcus ceased and in the same manner the bacteria of the Proteus group died off during the fermentation; the bacillus of the Subtilis group, on the other hand, continued to grow. Of the bacilli mentioned, the *Diplococcus tabaci* is said to improve the combustibility; the *Bacillus tabaci*, of the Proteus group, the aroma. Koning, who tested these bacilli in the course of various experiments, determined that an improvement in aroma and taste was invariably recognizable, in some cases he observed even an improvement in the combustibility and in each case the inoculated tobacco was better than that which had not been inoculated. As regards the inoculation of the leaves with bacteria, Koning did not use for this purpose bacteria suspended in water, but a mixture of the bacteria with tobacco powder, which was then blown into the tobacco bunches.

In 1899, Loew determined that the fermentation of tobacco leaves was not effected by living organisms, but by oxidizing enzymes of the tobacco leaf itself. Where bacteria or other organisms obtained access during this process, they caused damage. According to his observations, the number of bacteria decreased with the advancing fermentation, in place of increasing, as other investigators had found. Loew found among the oxidizing enzymes in the tobacco leaves an oxydase which at once colored tincture of guaiac blue, also a peroxydase which only colored tincture of guaiac blue in the presence of peroxide of hydrogen.

The tobacco oxydase died when a watery solution was heated to 149 deg. to 151 deg. F., the peroxydase at 188 deg. to 190 deg. F. Both were present in fresh, barn-ripe, and fermented tobacco, but after a few years were destroyed. In a Florida tobacco, kept for two years, Loew found only the peroxydase. According to his opinion these oxidizing enzymes act during the barn-ripening and fermentation on the tannin and the nicotine like the diastase on starch. Consequently, chemical changes occur in the tobacco leaf. According to an experiment the nicotine is attacked by the tobacco oxydase, ammonia being formed. It is possible that during the fermentation new quantities of oxydase are formed from a zymogen. To what extent the peroxydases, catalases, participate in this remains to be determined by precise investigation.

As regards the disturbance in the fermentation process, too large a proportion of moisture, as far as we have to deal with organisms, may prove dangerous. There may easily occur then, owing to the action of decomposing agencies, decomposition phenomena. Thus, in the fermenting rooms, the thread fungus *Botrytis* was found, in a damp bale also the bacteria of butyric acid.

After fermentation, the tobacco is subjected to different processes. The different sorts are mixed together, the tobacco is kilned, extracted and allowed to age. As far as the aging of tobacco is concerned, saccharine substances are frequently added, yeast and invariably water. These are the so-called "sauces." Directions for this are given by Wagner, Kissling, and others. Another method of aging consists in so-called petunizing. For this purpose leaves that are of little value for cigar making and are still faultless as to aroma are selected and allowed to remain about eight days in water, where they begin to rot. After the tobacco crop has completed its fermentation and the leaves are dry, they are made up into separate bunches and sprinkled with a little water, but so that the leaves are not too wet. Too great a softening of the leaves must be avoided, otherwise decomposition occurs too readily. The leaves are then done up in bundles and the superfluous water allowed to drip off. After the

* Pure Products.

bunches have hung for a few hours in the drying house they are packed, pressed firmly together, into boxes. Here a fermentation takes place incited by the fermentation organisms introduced into the leaves with the water of maceration.

The effect of the petunizing fluid is due to the carbonate of ammonia developed during the fermentation of tobacco leaves or stems and which increases the alkalinity of the tobacco treated. According to Loew, it increases the activity of the oxidizing enzymes. For this reason carbonate of ammonia is frequently used directly in petunizing fluid, to which other additions, such as rum, molasses, wine, etc., are made.

Splendore found that the simultaneous treatment of tobacco with petune and carbonate of ammonia improved both its flavor and aroma. In addition, the same investigator discovered that the tobacco was improved by pasteurizing in live steam at a temperature of 140 deg. to 212 deg. F. Here the characteristic odor of a well-conducted fermentation, the bread aroma, occurs. Pasteurization increases the durability of the tobacco; mold fungi, yeast, and bacteria which exist on the green leaves are killed, according to Splendore. In addition the flavor is improved and the color made uniform and darker.

As far as the aging or maturing of the tobacco is

concerned, this is sometimes of no advantage as far as quality is concerned. Very often it is the cause of the development of organisms that cause cigars to become moldy and greatly detracts from their quality. Behrens discovered that the moldiness of cigars was caused extensively by the common *Aspergillus glaucus*.

After all that has been advanced it is very probable that bacteria, in the fermentation of tobacco, play a part very similar to that noted in the spontaneous heating of hay. Unfortunately we have not as yet succeeded in finding fine bacteria that will bring good fortune to the business, nor a domestic tobacco that could be improved by pure culture.

VON ZEPPELIN'S FORGOTTEN PREDECESSOR.

ZAMBECCARI'S WONDERFUL AIRSHIP IN 1803.

THE name of the man who apparently first succeeded in steering an airship has long been forgotten by the generality of mankind and is preserved only in the French "Biographie Universelle," in a little-known book of travels by Kotzebue, and perhaps in a few technical works. It is worthy of remark that this pioneer in aviation was neither a physicist nor a mechanic, but a naval officer.

Count Francisco Zambeccari was born in Bologna in 1756. After completing his education, which was a very good one, he entered the Russian navy. In 1787 he was captured by the Turks and held prisoner in Constantinople for three years, but finally obtained his liberty through the intercession of the Spanish ambassador. Thenceforth he devoted his entire attention to the problem of directing the course of a balloon. His predecessors, with the exception of Leonardo da Vinci, had constructed fantastic devices which had no basis in experiment or observation of nature. Zambeccari, on the contrary, investigated the peculiarities of atmospheric currents and was the first to conceive the idea of controlling the course of a balloon by rising or sinking from an unfavorable to a favorable current in which oars could be used to advantage. In order to carry out this idea it was necessary to construct a balloon which could be raised and lowered at will, and this Zambeccari succeeded in accomplishing. The inventor was killed in 1812, by the wreck and explosion of his airship, in which he had ascended during a storm.

Kotzebue had found him in bed, suffering from the effects of an earlier accident. Kotzebue says: "I learned the details of his ever-memorable flight through the air and saw the lamp which enabled him to rise and fall at will. It is undeniable that a balloon in equilibrium with the surrounding air can be moved horizontally by an exceedingly small force. This force in Zambeccari's balloon is furnished by small attached vanes. I was assured by several eye witnesses that he undertook to fly to and return from two towers, situated in different directions from the starting point, and accomplished both tasks with a very small expenditure of force."

One of Zambeccari's memorable flights was made in 1803. He was enabled to construct his airship by the interest of the celebrated mathematicians, Saladini and Canterzani, who induced the government of Bologna to grant him a subsidy of about \$1200. The flight had been announced to take place on September 4, but bad weather and small defects in the balloon had necessitated a postponement until October 5. On that day the weather was again unfavorable, but Zambeccari was driven by the jeers of the spectators, as he said afterward, to attempt the flight. The balloon was carried out to sea, where it and its half-drowned occupants were secured by fishermen and taken to Pola.

Although this attempt had ended in so disastrous a failure, the professors, Saladini, Canterzani, and Avanzini, to whom the matter had been referred by the government, brought in a favorable report, which was printed November 9, 1804. This report is not accessible to the present writer, but Kotzebue, who appears to be well informed, gives substantially the following account of it.

The referees lay down the following law: "If a partly-filled balloon is in equilibrium with the air at any point, for example, near the ground, it will remain in equilibrium when it is transported to any other point of the atmosphere, so long as the gas is allowed to expand or contract." In other words, the volume of the partly-filled balloon varies inversely as the density of the surrounding air. The referees acknowledge that this law was discovered and demonstrated by Zambeccari. Then they speak of his lamp, of the absence of danger in the use of fire and inflammable gas in the same apparatus, and of the manner in which Zambeccari, with the aid of the lamp, was able to ascend, descend or remain stationary at will,

without using a single valve, losing any inflammable gas or throwing out any ballast. Then they describe the rudders, by means of which the inventor had previously propelled a light car suspended from the roof of a church, in the presence of many persons. They assert that Zambeccari, in his last flight, hovered a long time over Bologna, ascended and descended at will, and described an arc round the city from south to west, solely by means of his flame and without throwing out ballast, for they had observed him carefully with telescopes. In conclusion they state that Zambeccari has succeeded in proving his theory by actual flight and that the misfortune that has twice overtaken him can detract nothing from his fame or the value of his invention.

The balloon used by Zambeccari was of the type known as a *rozière*, consisting of a *montgolfière*, or fire balloon, in combination with a gas bag, the latter giving a constant buoyancy, while ascent and descent could be accomplished by regulating the flame of the hot air balloon, without expending gas or ballast. Just what the "oars" were with which he drove his craft is not clear.—Umschau.

EXPLOSION AND FLAMELESS COMBUSTION.

THE conditions under which explosive mixtures of gases can be ignited by electric sparks and by glowing bodies are now attracting attention in all countries. It is a matter of great importance in view of the rapid introduction of electric machinery in coal-mines, and already there is a considerable amount of experimental data available for those interested. The latest experiments are those of MM. H. Couriot and Jean Meunier, who have made an extensive series of experiments of the action (1) of incandescent wires or filaments, and (2) of electric sparks, on mixtures of methane with various proportions of air, the gas being sometimes at rest and sometimes in motion. A current of a mixture of 80 per cent of methane with 20 per cent of air could not, they found, be ignited by an incandescent wire or by the spark produced when the wire melted owing to the intensity of the current, though in contact with a flame the mixture readily ignited. Whatever the proportion of methane and air, the only effect produced by an electric spark, if any effect were produced, was an explosion, not quiet combustion. Under no circumstances were they able, in these experiments, to cause an explosion by means of an incandescent wire. When an explosion was caused, it was found to be due to the breaking of the wire, and the consequent production of a spark. The most readily explosive mixture contains 9.5 per cent of methane, and slight explosions were observed with so low a proportion as 4.5 per cent. The maximum limit of explosibility is 12 per cent of methane, and this was shown by the fact that they never obtained an explosion when the percentage reached 12.25. So long as the proportion does not fall below 5.5 per cent combustion during the explosion is complete. If the current by which the wire is heated is shunted with a parallel shunt, the production of the spark causes no ignition when the resistance in the two branches is equal. If the resistance of the shunt is high, the production of a spark always leads to the ignition of the mixture; on the other hand, when the resistance of the shunt falls below a certain value, the production of a spark again causes an explosion. In order to avoid an explosion, the intensity of the current must not exceed a certain maximum, which depends on the resistance of the two branches of the conductor. (Comptes Rendus, 145, pages 1161 to 1163.)

In explanation of the observations that an inflammable mixture of methane (marsh gas) and air is not ignited by an incandescent wire or filament unless the wire fuses and a spark is produced, it is suggested that the incandescent filament attracts the molecules of oxygen and repels those of the hydrocarbon in the mixture; the zone of gas, therefore, surrounding the

filament is too poor in the hydrocarbon to be inflammable, and slow or rapid combustion occurs with the formation of water and carbon dioxide. When, however, the filament employed is sufficiently large in volume to raise by its radiation the temperature of the surrounding inflammable mixture beyond the inert zone to about 650 deg. Cent., it appears that an explosion may occur. Hauser has recently shown that, while an incandescent ferro-nickel wire 0.3 millimeter in diameter does not ignite a mixture of 9 per cent of methane with air, a soft-iron wire 0.9 millimeter in diameter caused an explosion in six out of seventeen times.

M. Jean Meunier has continued these researches: (1) determining the limits of inflammability of explosive mixtures of ether and air, also Boudouard and Le Chatelier; and (2) investigating combustion without flame and the inflammation of gases at the end of a metallic rod. (Comptes Rendus, 146, pages 539 and 540.) In the last-named an interesting experiment with an alcohol-fed incandescent burner is described. The burner is supported on four tubes, through which cotton wicks pass, supplying the alcohol. It is of annular form, the flame rising from a central cylinder which is covered with a convex disk, perforated with closely-spaced holes. An iron stem 8 centimeters long, supporting the mantle, rises from the center of the disk. It is observed that after continued use the light becomes less bright, and that the darkening of the mantle commences at the moment that the disk reddens. After removal of the mantle the author succeeded in getting the disk to redden without production of a flame. Once the disk is red hot, its incandescence increases with the supply of alcohol vapor. It is impossible to ignite either a pad moistened with alcohol, a wick soaked in light petroleum, or a wooden match, by bringing them near the hot disk, although the wood of the match is charred. This flameless combustion occurring on the surface of the disk is to be explained by the retention of either the combustible gas or the oxygen on this surface, as suggested by Couriot and Meunier in the case of the non-explosion of an explosive mixture by an incandescent filament. Application of a lighted match to the upper end of the metallic stem results in the production of a flame at this point, which continues to burn so long as it is protected from drafts, showing that the oxygen is retained by the disk, and that the excess of combustible gas rises around the stem without mixing with the surrounding air and without igniting, although the stem is at a fairly high temperature. The shape of the flame can be altered by varying the form of the upper extremity of the metallic rod. The experiment, besides its important bearing on the mechanism of combustion by incandescence, seems to indicate that gases have a certain amount of cohesion. It may be noted in passing that the late Mr. Thomas Fletcher, of Warrington, on more than one occasion, several years ago, publicly performed experiments showing the flameless combustion of coal-gas.—Engineering (London).

According to a recent consular report the large deposits of peat in the world, which have hitherto been of practically no value, may prove to be of great worth if the recent experiments of a French chemist, Raynaud, in distilling alcohol from peat prove to be practical, as now seems probable. The cellulose of the peat is hydrated and converted by means of sulphuric acid into a soluble carbohydrate, which is fermented by means of a special yeast, which has hitherto been kept secret. According to statements made by Prof. Ramsay, one ton of dry peat is said to furnish about 43 gallons pure spirits and about 66 pounds of sulphate of ammonia of 100 per cent. It is asserted that the manufacturing costs of 1 gallon of alcohol of 97 per cent are at most 10 cents, whereas the cost of the process of distilling from potatoes is about 35 to 40 cents.