blindness, usually since youth. The central vision is sometimes normal, sometimes reduced. In some the visual field is restricted. There is often, but not always, consanguinity in the parentage. Several instances occurred in brothers and sisters. The greater part of the fundus, but not the periphery, is studded with minute dull white dots, while the macula is nearly free when central sight is still good. In many of the cases pigment figures were seen in the periphery of the fundus sometimes described as not characteristic of retinitis pigmentosa. In cases with reduced sight the nerve appeared moderately pale. It has been repeatedly observed that the disease can remain stationary for many years and that it progresses rapidly.

Cases of this kind have been reported by Gayet,⁹ Dor¹⁰ and Nettleship.¹¹ Fuchs¹² subsequently interpreted these cases as atypical retinitis pigmentosa. Further reports have since appeared by Liebrecht,¹³ by Griffith,¹⁴ also by Tr. Collins and Swanzy,¹⁴ and Wuestefeld.¹⁵ The latest of the same kind are reported by Wicherkiewicz¹⁶ and by Takayasu,¹⁷ who saw two instances in Japan.

CASE 4.—Patient.—Miss L., aged 29, anemic, complained of headaches, especially in the afternoons, and of fatigue of the eyes. Did not see well at night.

Examination.—V. R. E. = 20/20; L. E. = 20/40, improved to 20/30 by Cyl. + 1, axis 135. Homatropin showed in addition H. = 1 D. in both eyes. Nerves and vessels normal. Near but not quite at the periphery of the fundus was a zone showing pigment figures in the form of delicate branching threads. In the L. E. there were two dozen white specks in the more central part and several areas of exceedingly fine white dots. A fainter dotting was also seen in the R. E. The headaches were permanently relieved by appropriate glasses. No change after one year.

Subsequent History.—Eight years later I saw the patient again. Her night-blindness was a trifle more troublesome, but her central vision was as good as formerly. The fields were still perfect. The ophthalmoscope showed no certain change in the appearance of the retinal pigmentation. The white dots, however, could no longer be found.

The case can not be considered ordinary retinitis pigmentosa on account of the atypical form of pigmentation as well as the persistence of practically normal vision for eight years in a patient now 37 years of age. As far as I know this is the first instance of this form of retinitis punctata watched during such a long time, and also the first in which the white dots have ultimately been found to disappear.

DISCUSSION.

Dr. M. WIENER, St. Louis, said he has had little experience with the punctate retinitis, described by Dr. Gradle as a retinal infiltration which disappears and sometimes reappears to fade away again, leaving no trace or scar. He has seen a punctate infiltration of the retina which he would term a senile form, in which the swelling would subside and leave little sharply outlined dots, as though they had been punched out with a conductor's punch. These may, or may not, affect the visual acuity according to what degree the macula be affected. Where the spots occur around the disc or between the disc and the macula, which is the most frequent site, we may have no disturbance of vision. But should they lie in the macula we have a marked reduction of the visual acuity. Dr. Wiener has examined the fundus of patients with no other

trouble than their presbyopia and found these atrophic spots, the patient unconscious of having had any inflammatory trouble in the retina. Then again, as Dr. Gradle pointed out, the retinal dots may be associated with some other trouble, as optic neuritis or with diffuse retinitis. He said he believed that in a great many cases the retina is only secondarily affected, the trouble being primarily in the choroid; the infiltration occurring almost at the same time, or a little later, in the retina. The atypical albuminuric form he has not seen.

Dr. R. S. LAMB, Washington, D. C., called attention to what he believes to be the foundation cause of retinitis punctata, the condition of the arteries. He asked if Dr. Gradle takes the blood pressure in any of these cases. He said he should think that the retina would be the first to be affected, rather than the choroid, because of the size of the vessels.

Dr. H. GRADLE, Chicago, said that in the two instances which seemed atypical forms of albuminuric retinitis there were probably vascular changes. In none of the others was there any reason to suspect this.

SURGERY OF TUMORS OF THE BLADDER FROM A MODERN ASPECT.

G. KOLISCHER, M.D., AND L. E. SCHMIDT, M.D.
CHICAGO.

Operative interference in tumors of the bladder is not at the present time conducted in a uniform manner, nor is it based on any generally accepted rules, as is the case in other domains of surgery. Each operator has notions and fancies of his own and operates accordingly, employing in many instances methods which, if applied in other branches of akiurgy, would bring ridicule on his head. Therefore, the possibility of approaching this subject in a more orderly and systematic manner, as well as the great scientific advantage to be derived therefrom, has induced us to attempt, by taking into consideration the work that we had observed in various genito-urinary clinics, by availing ourselves of the pertinent literature, and by analyzing our own experiences, to establish certain basic principles on which, as a foundation, might eventually be built up a rational and consistent surgery of tumors of the bladder.

From the very first one essential principle becomes apparent: benign tumors should be approached from the inside, malignant tumors from the outside of the bladder. The intrinsic qualities of a benign tumor are such that it is manifestly sufficient if its mucosa, submucosa and pedicle alone be removed; it is not necessary in operating on such a tumor to go through the entire thickness of the bladder wall and resect all its layers.

CYSTOSCOPE OR INCISION?

In approaching benign tumors from the inside of the bladder we have at our command two methods: one, the endovesical, through an operative cystoscope; the other through an incision in the viscus, which renders accessible the implantation of the growth. In the endovesical method, after proper dilatation of the bladder, the operative cystoscope is introduced, the tumor is focused and the galvanocautic snare is developed into a loop within the bladder. Then, in case the tumor is not too large, the loop is passed over the growth and down to its base and constricted until it fits snugly around the pedicle. The electric current is now turned on and the interpolated resistance is diminished until little bubbles appear around the loop, after which the pedicle is seared through by slowly constricting the loop. Then, if deemed necessary, the galvanocautic loop-carrier is withdrawn and a straight, solid galvanocautery is in-

9. Arch. d' Ophth., iii, 386.

10. Id., 481.

11. Trans. Ophth. Soc., U. K., iv, 162; vii, 301; viii, 163.

12. Arch. f. Ophth., 39, iii, 65; and Arch. of Ophth., 1888, xxvii, 484.

13. Klin. Monatsb. f. A., June, 1895.

14. Trans. Ophth. Soc., U. K., 1897.

15. Ztschr. f. A., v, 2.

16. Trans. of Lisbon Congress.

17. von Graefe's A. f. Ophth., lxi, 281.

serted in its place, by means of which the stump of the pedicle is again cauterized.

The severed tumor either is picked up by forceps introduced through the cystoscope or, after removal of the cystoscope, is pumped out through an evacuator; if very small it may be left to be expelled by the urinary stream. The suggestion of using a steel wire snare, instead of the incandescent platino-iridium loop, to crush the tumor and thus prevent subsequent hemorrhage (B. Klose) seems to have been based largely on theoretical speculation, for actual experience has not justified its claims. In case the tumor is rather large it may be removed piecemeal by means of the galvanocautic snare; when the pedicle is reached it is seared through by means of the loop or burned down with the solid galvanocautery.

Although the instruments almost exclusively used by the operators who are doing endovesical work are those with indirect view, we desire to state that, in our opinion, which is based on manikin practice and experience on the living subject, it is of decided advantage to use the cystoscope of direct view in operations of this character. As a matter of fact, the cautery of Nitze's operative cystoscope almost entirely obstructs the view of the point of its application; this defect has led to the suggestion of using a fenestrated galvanocautery (B. Klose). With the instrument of direct view it is much easier to pass the wire loop over the tumor and down to the insertion of the pedicle, first, because the loop and the tumor are in the direct line of view, and, second, because of the short distance between the window, the loop-carrier and the growth. The same reasoning holds good in the use of the solid galvanocautery. It must be admitted, however, that in operating endovesically on a tumor located in the vertex of the bladder the cystoscope of direct view can hardly be used; in such cases, which, however, are extremely rare, the so-called long wire-loop of Nitze's operative cystoscope should be employed. Judging from our experience on the manikin, it would seem as though many of the disadvantages observed in the operative cystoscope of Nitze might be avoided by using an instrument of indirect view, constructed after the pattern devised by Casper.

There are some writers who object on principle to all endovesical operations so far as tumors of the bladder are concerned, basing their objections on the possibility of subsequent hemorrhage, the less extensive excision of the base of the tumor, and their belief that, since such an interference calls for specialized study and skill, it will never become a routine surgical procedure (Cathelin). Technical objections coming thus from men who have had no personal experience with the endovesical methods have no weight. The theoretical objections as to the insufficient extensiveness of the operation are refuted by the statistics of those who are actually doing endovesical work (Nitze, Wenrich, Klose, Casper, Keersmaecker). Among these Nitze stands pre-eminent; there are no statistics which compare even approximately with the splendid results obtained by this master and his pupils. In the light of such work even those who are most pessimistic as to the outcome of operations on bladder tumors, however radical the excision may be, are beginning to realize more and more the necessity of employing the endovesical procedure (Keydel).

EXSECTION.

The other method for the removal of the benign tumor involves its bloody execution after the bladder has been opened by an incision. The only route through which

such an operation should be approached is the suprapubic, for in this way free access is obtained to all parts of the bladder. We desire to submit, as the fundamental principle of this operation, that it be conducted with one object always in view, i. e., the complete closure of the bladder after the tumor is extirpated. In order, however, that primary union and uneventful recovery ensue, it is of paramount importance to bear certain details carefully in mind, remembering always that a liberal incision through the anterior wall of the bladder is requisite for smooth, clean surgical work inside the viscus; nowhere is button-hole surgery more out of place than in bladder work.

The anterior aspect of the bladder should, therefore, be freely exposed in order to gain space for the generous incision; perivesical fat and the peritoneal duplication are to be stripped off until a sufficiently large area of the bladder wall is exposed. The bladder wall is caught by a few sutures, which are used later on as guy ropes. The bladder is incised between these sutures and all squirting blood vessels are ligated. The tumor and its base are now brought into full view and made easily accessible either by catching the base of the bladder with sutures and pulling on them so that the field of operation is brought into the incision and close to the surface, or by having an assistant insert two fingers into the rectum or into the vagina, as the case may be, and thus force the base and the fundus of the bladder upward (Fig. 1). Sutures are now run beneath the insertion of the tumor and are promptly tied one by one in the path of the operator as he is excising the base of the growth (Figs. 2 and 3).

This detail in technic prevents the hemorrhage which otherwise would flood the field of operation and necessitate extensive sponging and continual hunting for the excision wound, and thus avoids a traumatism of the mucosa which might interfere with the prompt and complete closing of the bladder. Care should be taken not to grasp the tumor with the fingers or the forceps, since, as a rule, such growths are very friable and are apt to bleed profusely when crushed or torn. If the tumor be located close to the vesical portion of one of the ureters one end of the ureteral catheter is run a few inches up into the ureter, previous to the insertion of the preventive stitches, and the other end is led through the internal orifice of the urethra and thence outside by way of the urethra. Should some of the preventive sutures encircle the vesical end of the ureter the fact would become apparent on attempting to remove the ureteral catheter; it would be engaged. In this case the catheter is simply left in place for a week or so, until the healing process has progressed sufficiently, and then withdrawn (Figs. 4 and 5).

After the main tumor has been removed it is very important to inspect the entire inner surface of the bladder in order to discover smaller-sized papillomata, should such be present. Particular attention should be paid to the parts opposite the main tumor, since we often find these smaller growths in those areas which come in contact with it when the bladder is empty. This condition is of such frequent occurrence that the suggestion of contact infection forces itself on the mind. All such smaller tumors, if present, should be removed in the manner described above; the erroneous diagnosis of a relapse of the removed tumor is often due to the fact that these have been overlooked and not extirpated. After the viscus has been cleared of all growths, large and small, and after the excision line or lines have been

closed by sutures, the incision of the bladder is closed by means of two layers of sutures. The essential feature of this suturing is to bring together only denuded and bleeding surfaces and to prevent any impaction of the mucosa into the suture line.

Fig. 1.—Benign tumor exposed.

The closed viscus is now so suspended to the abdominal wall as to exclude any dead spaces and thus prevent the accumulation of secretion; it is important, however, in order to prevent pressure necrosis, that the sus-

Fig. 2.—Preliminary sutures run underneath the base of the benign tumor.

pension sutures do not interfere with the suture line closing the vesical incision (Fig. 6). The abdomen is closed in the usual manner. As a rule patients urinate naturally a few hours after the operation; it is only when a patient does not do so that catheterism at proper

intervals is resorted to. The permanent catheter should never be used, as it offers no advantage; on the contrary, it always leads to irritation, quite often to infection.

Fig. 3.—Suture line after removal of the tumor.

TRANSFORMING BENIGN INTO MALIGNANT GROWTHS.

It is often contended that the simple removal of benign tumors, such as is indicated above, leads in the majority of cases to recurrence, and that all recurrences are of a malignant character; on the other hand, there

Fig. 4.—Ureteral catheter inserted into both ureter and urethra to prevent tying up of the ureter by the preliminary sutures. Ur. ureteral orifice; C, catheter.

are opinions which differ entirely from this contention, some coming from authoritative sources (Lubarsch, Birch-Hirschfeld). A kind of middle ground is taken by Wendel:

We are compelled to believe that not infrequently bladder tumors will appear which are, in outward appearance and in histologic and clinical characteristics, unquestionably benign, but which later on might become transformed into cancer. In all such cases there is a stage in which surgical interference

Fig. 5.—Suture line after removal of the tumor with catheter in place. Ur, urethra; C, catheter; U, ureteral mouth.

Fig. 6.—The closed bladder suspended to the abdominal wall by two sutures.

may occur early enough to remove these tumors with the same favorable outlook as is the case in other benign tumors.

As a matter of fact, most authors have relinquished the idea that benign tumors suffer relapse of malignant

character (Rafin). All men of actual experience concur in the opinion that benign tumors should be operated on immediately after their discovery (Rafin, Malesherbes, Hegue, Escat, Desmos, Carlier, Hamonic, Nitze, Fischer, Casper, Fedoroff, Bary, Fedenat, Albarran and others).

Operations for the removal of malignant tumors should be guided by principles derived from sound surgical reasoning and not by traditional superstition. We believe that all operations of this character should be started from above, but at the same time we wish to say emphatically that the traditional method of making a suprapubic incision into the bladder and of then hunting for the malignant tumor from the inside should be absolutely condemned. If a surgeon were dealing with a pyloric carcinoma, would he first make an incision through one of the curvatures and endeavor, by

Fig. 7.—Excision of section of bladder wall carrying a malignant tumor. B, bladder; F, section.

working his way through this opening, to dig the cancer out from the inside of the stomach? Certainly not! Why should not the same reasoning hold good in operations for malignant tumors of the bladder? The incision into and through the bladder wall should be made as close to the location of the tumor as is consistent with the obvious demand that the lines of excision should run through apparently healthy tissue (Fig. 7). Making a routine median incision into the bladder regardless of the location of the tumor means inflicting an unnecessary injury on the bladder. The recognition of these facts calls for a topographic diagnosis of the tumor before operation and demands a free exposure of that portion of the bladder which is going to be in the field of resection. Sometimes the base of the tumor can

be definitely outlined, or the presence and extent of infiltrations ascertained only by palpation after the bladder is made accessible by detachment and incision.

The steps of the operation should be as follows: The suprapubic incision is made; the perivesical fat, which is thus exposed to view, together with the peritoneum, is stripped from the bladder to such an extent that the anterior aspect of the viscus is freely exposed. A few sutures are now drawn through this part of the bladder to serve as guy ropes in facilitating its further manipulation. The denuding of the bladder and its detachment from the adjacent tissue is continued until its exterior surface, in so far as it corresponds with the seat of the tumor inside, is exposed, thus giving additional leeway for resection through healthy tissue. The flap formed by the perivesical fat is now tilted back over the peritoneum and fastened there with a few stitches; in this way a firm protection is furnished for the peritoneum against subsequent infection.

The bladder is now dragged to one side by means of the guy ropes and an incision is made into its wall as close to the site of the tumor as is compatible with surgical demands. Then, by means of the index finger passed into the bladder through this opening, and the thumb outside, the extent of the second incision is determined by palpation; this second incision completes the resection and removes from the entire thickness of the bladder wall a sector which carries with it the tumor. It goes without saying that the excision will, as far as possible, be made in such a way as to facilitate the approximation of the edges of the wound. The resec-

Fig. 8.—Suture line closing up wound after resection of flap out of the lateral wall of bladder.

tion, in case it does not remove more than one-third of the bladder, should be followed by total suturing of the viscus (Fig. 8). After the organ is closed, a cigarette drain is run down to the deepest point of the bed of the wound, the bladder is suspended to the abdominal wound, and all is closed up save for the point of drainage. The permanent catheter should never be used in cases of partial resection and subsequent complete suturing of the bladder; the patient either urinates naturally or is catheterized at appropriate intervals.

WHEN THE URETERAL OPENING IS INVOLVED.

Should the tumor involve a ureteral opening, the bladder is dissected out to such an extent that not only complete excision of the tumor base becomes possible, but also the vesical end of the ureter is laid bare. The ureter is clipped off at its insertion into the bladder, and

then the exsection of the tumor base, including the ureteral mouth, is accomplished. The stump of the ureter is now reimplanted into the upper end of the excision wound, after one of the standard methods, the wound of the bladder is closed as completely as is feasible (Figs. 9 and 10), and the wound cavity attended to appropriately.

In case a more extensive resection of the bladder wall is necessary, it becomes impossible to close up the organ entirely. However, an attempt should always be made to reform the bladder, as far as possible, by suturing and by attaching the edges of the remaining portion of the viscus to the abdominal muscles. It is astonishing to observe what a small remnant of the bladder will suffice to start the formation of a serviceable organ inside of a few weeks. We have on record three cases, two cancers and one sarcoma of the bladder, in each of which more than two-thirds of the viscus was removed. After about

Fig. 9.—Section carrying tumor involving ureteral mouth cut out of bladder wall. A, puncture line indicating line of excision; B, bladder after excision; C, flap with tumor and ureteral stump; D, outside aspect of bladder after implantation of ureter and closing up of the excision wound.

eight weeks in each instance, all was healed over and the patient was able to retain from 250 c.c. to 300 c.c. of urine; the action of the bladder was perfect, the stream being projected with force, the viscus completely emptied. One is induced to believe that muscle fibers grow from the bladder remnant into the cicatricial tissue which forms over the mutilated organ, and thus impart contractile power to this newly formed portion of the bladder.

In those cases in which a radical operation is no longer possible, there are two methods of relieving the patient's suffering. One method is to open the bladder from above and cauterize the exulcerated surfaces with a Paquelin; as a rule, this intervention does not benefit

the patient to any great extent, at least not for any length of time, for the urine flows into the bladder and over the excoriated and irritable surface of the tumor, and thus necessarily leads to permanent pain. The other method, which furnishes more satisfactory results, consists in establishing permanent kidney fistulæ and thus diverting the urine from the bladder; the patient is at once relieved from all pain and is made quite comfortable. A belt carrying two small pouches keeps the patient comparatively dry.

The choice of anesthesia is of the utmost importance in all cases of bladder surgery. Endovesical operations can, as a rule, be carried on under the influence of an antipyrin anesthesia supported by the rectal application of morphin. For the cutting operations, we always employ, as a matter of routine, the anesthesia produced by laughing gas. This anesthesia, besides being free from danger, so far as the heart is concerned, possesses a great advantage in that the patient returns to full consciousness immediately after the completion of the operation; this enhances the possibilities of natural micturition, of very early breathing exercises, and of the early administrations of food and drink. Up to the present time we have never observed any untoward influence of gas-anesthesia on the kidneys.

Fig. 10.—Inner aspect of the bladder showing the implanted ureter. Flap operation.

The next question that naturally arises is whether or not it is possible to determine, previous to the operation, the character of a bladder tumor; we do not hesitate a moment to answer in the affirmative and to say that such diagnostic work can be done with accuracy sufficient for all practical purposes. Any man with experience in cystoscopic diagnosis can recognize a benign tumor from its appearance (Nitze, Weinrich); tumors of doubtful nature are to be considered and treated as malign. In this connection it should be borne in mind that the benign papillomata comprise the majority of bladder tumors (Civiale, Thompson, Guyon, Soemmering, Kuester).

It was formerly held by a great many surgeons that, because of the poor permanent results and the quickness of the relapses, patients with malign bladder tumors should never be operated on. This view is shared at the present time by some writers. In our opinion, however, the soundness of such reasoning is, to say the least, open to discussion because, in the first place, a great many patients were and are still being operated on in an irrational manner, and, in the second place, because the diagnosis is quite often made so late that the chances for a successful operation are greatly reduced; the more

general employment of the cystoscope, however, will greatly improve matters as to the latter point. Furthermore, it should be taken into consideration that the use of rational operative methods and the uniform employment of gas anesthesia will certainly reduce the primary mortality of these procedures.

As it is, there are two cases of vesical cancer on record in which total extirpation of the bladder accomplished a cure without relapse for five years (Hogge), and for fifteen years (Pawlik), respectively. Rafin reports five cases of partial resection of the bladder for carcinoma that remained cured for more than three years. In Czerny's sarcoma case the patient lived for twelve years after the operation without relapse. Quite a few surgeons report cases of malign bladder tumor which, after partial vesical resection, have remained cured for from one to three years (L. Davis, Pousson, Pitha, Leschnew, Albarran, Alessandri, Schmidt, Kolischer).

Mikulicz, in his last publication before his death, declared that he felt convinced that he had relieved the sufferings and prolonged the life of his patient every time he had operated for cancer of the bladder. Surely so impressive a message gives warrant for our belief that, even though nothing else were accomplished than the adding of a few years of comfort to the sufferer's life, the operation would be fully justified.

CONCLUSIONS.

In conclusion we desire to submit, as the result of our work along these lines, the following fundamental propositions:

1. All benign tumors of the bladder should be approached from the inside, all malign tumors from the outside of the viscus.
2. In all malign cases in which the loss of substance is not too great the bladder should be closed completely by sutures after the removal of the tumor.
3. In cases of malign tumor the incision into the bladder should be made in accordance with the location of the growth, as defined by the cystoscope.
4. The permanent catheter should be absolutely abolished.
5. Gas anesthesia should be employed exclusively.
6. A constant cystoscopic surveillance should be maintained over any bladder that has ever been operated on for tumor.

ONYCHOMYCOSIS TRICHOPHYTINA,

WITH REPORT OF TWO CASES.*

A. RAVOGLI, M.D.

CINCINNATI.

The nails, like any other epidermic tissue, are exposed to the invasion of the trichophyton, yet cases of the kind occur rather rarely. Saubouraud, in the department of the ringworm affections in the Hospital St. Louis, considers onychomycosis trichophytina a rare disease. Neuberger¹ referred to the fact that Anderson, in Glasgow, among 11,000 cases of cutaneous diseases, of which 178 were of trichophytic nature, did not have a single case of trichophyties of the nails. White, of Boston, in 5,000 cases of diseases of the skin, with 180 cases of tinea trichophytina, did not have one case of trichophytic onychomycosis. On the other hand,

* Read in the Section on Cutaneous Medicine and Surgery of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

1. Ein Fall von Onychomycosis Trichophytina, Derm. Ztschr., xi, 551.