place by wearing a truss, in less than three, four, or five years.

Now you will hardly be surprised if it occasionally happens, that a protrusion takes place while the patient is wearing the truss, which places him in a state of considerable danger; for although a person wears a truss, a force may push down a portion of the bowel below the side of the pad, and thus strangulation may take place; and I should observe, that if the neck of the sac has become contracted by the pressure of a truss, and a protrusion then takes place, it is very likely to become strangulated, because the orifice is so much more contracted than it was before. Hence persons have sought for some means of radically curing rupture and heretofore various means were adopted for that purpose, which are now become obsolete.—The older surgeons, with a view of radically curing rupture, dissected down to the swelling, insolated the hernial sac, and tied a ligature round its outside near to the orifice at which the parts were protruded from the abdomen; or they cut into the parts, returned the protrusion, and then sewed the sac up with sutures; and some who went a shorter way to work, performed the operation of *castration*—cut away the spermatic cord, testicles, tumour and all! Such means have been employed for *radically* curing ruptures. Now you will easily understand, that all those proceedings must involve a risk equal to that of strangulated hernia. If any of those measures were adopted, they would put the life of the patient in as much danger as the operation for strangulated hernia could do, and this is only performed to save the patient's life; it is not done with the view of radically curing the complaint, for it does not secure to the patient a radical cure—he must wear a truss afterwards. You will easily see therefore, why all measures of that kind have been abandoned. Instances have been known in which patients have submitted to have a truss that pressed very powerfully upon the sac so as to bring on a state of inflammation there, and it is possible that inflammation thus produced, might cause agglutination of the sac, and prevent the reproduction of the tumour. But this is also a very uncertain and dangerous mode of proceeding, for if you were once to excite inflammation in a part of the peritoneum, you can have no security for limiting it just to the point to which you wished it to extend; it might spread over the whole cavity of the abdomen, and the patient might lose his life by such an attempt to rid him of the hernia. It is therefore now considered that a patient who has a reducible hernia must be contented with the palliative relief which the application of a truss will afford.

MR. CASTLE'S CASE OF STONE, TREATED
BY BARON HEURTELOUP.

To the Editor of THE LANCET.

SIR,—I have yet to send you several interesting cases of stone, successfully treated by Baron Heurteloup, according to his improved system of lithotritry. From amongst them I have first selected that of Mr. Castle, an eminent surgeon of Clare, who, after witnessing the case of his friend Major Moore, entrusted himself to the Baron's care.

Mr. Castle has drawn up a sketch of his own case, and I think his paper, accompanied by the Baron's notes, will form a document not altogether uninteresting to your numerous readers. One truth, at least, will be evident, on a careful perusal, that the domain of lithotritry has been much extended by the labours of Dr. Heurteloup. These reflections are made as a salvo to the reputation of the beautiful operation, in case of unsuccessfulness in any of these first essays; and it behoves us to be guarded in our deductions, lest we ascribe, to the insufficiency of the art, that which may be the effect of particular want of skill, or of possessing inferior instruments.

In his first endeavours at lithotritry, every surgeon is naturally anxious to possess the best instruments, and as little doubt can now be entertained of the superiority of the Baron's armoury, it is to be hoped he will shortly give that extensive publicity to his machinery which surgeons have been so long and anxiously expecting.

I have the honour to be, &c.

D. O. EDWARDS.

Westminster Hospital, Aug. 14, 1830.

Mr. CASTLE to Baron HEURTELOUP.

My dear Baron,—I have the pleasure to present you with an abstract of my case, as well for your satisfaction as for that of such sufferers from that terrible malady as may wish for authentic information, respecting the success of your method of treatment. In drawing up the enclosed history, I have confined myself to a statement of results only, expressed in general terms. For further particulars, I must refer to those gentlemen who witnessed your operations, and have carefully noted the results; with that view I have given the names of three medical gentlemen, fellows of the Royal College of Physicians of London, who will, I have no doubt, feel pleasure in affording to inquirers all the information on the subject in their possession. I cannot conclude without, in the strongest manner, expressing my conviction of your consummate skill, and my warmest acknowledgments of gratitude and

esteem for your unwearied attentions and kindness. I leave you perfectly at liberty to make use of this letter and accompanying case in any way you may think proper; and I am, my dear Baron, most sincerely,

Your obliged and faithful servant,

ANDREW CASTLE.

To Baron Heurteloup, 1, Vere Street,
Cavendish Square.

Mr. CASTLE'S Case.

About the year 1824, I became subject to nephritic attacks, more particularly in the winter. Since that year, no winter has passed without one or more returns of nephritic pain, and of other signs of urinary calculus. In January, 1830, the diagnostic symptoms of stone in the bladder were first observed. In June following I went to Dublin, where my friend Mr. Crompton, surgeon-general for Ireland, ascertained by the sound the presence of a calculus. By his recommendation I proceeded to London, to put myself under the care of Baron Heurteloup, of whose success I had repeatedly heard, and of which an example had fallen under my own eye in the person of an officer of my acquaintance. On the 24th of June I was first sounded, and, on the 26th, first operated on by the Baron. Between the 26th June and the 25th July following, I underwent eight lithontritic operations, after each of which I was able to return on foot to my lodgings without inconvenience. At various periods between the operations, fragments passed with the urine; a few times I was obliged to call for the Baron's assistance, to enable me to get off fragments larger or more angular than usual, which that gentleman effected with facility and singular dexterity.* During the whole of the time, from June the 26th to July 25th, I found my sleep, appetite, and strength, &c., as good as at any time for the twelve

months preceding, with the exception of two or three days of the last week, during which I, as well as most of my acquaintances, suffered from the oppressive heat of the weather. Immediately after the eighth operation, July 25th, the last fragments came away, and no symptom of calculus remained; and on the 27th, no particle or fragment could, after the most minute examination, be detected. The fragments that passed at various times between June 26th and July 25th, are now in my possession, and weigh altogether 243 grains, exclusive of a great deal of minute particles, too fine to be easily collected; the fragments appear to consist almost entirely of uric acid. I must here remark, that the first operations were performed with the three-branched instrument; and although the Baron appeared to me to seize the calculus with great facility, yet he did not succeed in destroying it with his usual rapidity, owing, no doubt, to the unfavourable shape of the stone, which appears to have been flat; in consequence of this he abandoned the use of that instrument, and had recourse to another, acting on a different principle, called "*brise-coque*," invented, I understood, by himself, by means of which he speedily relieved me completely;† each

* There exists in some individuals a particular disposition to retain fragments of stone in the urethra for a considerable time, an accident which occasionally annoys them more than the operation of crushing the stone itself. These fragments, when thus entangled, require to be immediately replaced in the bladder, or to be extracted entirely; and it often occurs, that these two indications are very difficult to fulfil. I have devised several instruments for these purposes, which render the necessary manœuvres extremely easy, and, above all, exempt from danger of wounding the canal. I beg here to observe, that lithotrity is little understood by those who imagine it consists in the destruction of calculi merely, which is frequently the most easy part of the process.—Dr. H.

† Mr. Castle was subjected *eight* times to the employment of lithontritic instruments, and I commenced by having recourse to the "*three-branch instrument*" of MM. Leroy and Civiale, with which I made *five* essays, and I concluded by resorting to the *brise-coque*, which only and alone was sufficient to bring the operation to a fortunate issue. In the first sitting, with the three-branch instrument, I seized the stone three times, and twice it escaped at the moment I was about to make the perforation; the third time I succeeded in making the drill penetrate into its interior, but this effected nothing more than the discharge of a very small quantity of powder, and of a narrow oblong fragment, which was evidently the edge of a flat stone. In the second sitting, I laid hold of the stone four times, and it as often escaped. The third seance was just as fruitless in its results. I grasped the hostile body once, but at the moment I was closing the branches for its permanent fixture, it slipped away, and probably fell into one of the lateral pouches of the bladder. This viscus was very large, and I did not succeed in recovering the stone. In the fourth sitting, I took the stone three times, twice it fled out of the embrace of the branches, and the third time I was enabled to hold it sufficiently fast to admit of the action of the bow, but, unfortunately, the stone being held in the same position as in the first sitting, the drill entered the same hole which it had made before. The fifth

operation performed with the brise-coque being followed by a quantity of fragments considerably larger than was ever produced by the three-branched instrument. On the whole, from what I could observe relative to the brise-coque, I think it an instrument of great value. It does not admit of the calculus slipping away after its seizure, as, to judge from my own case, the three-branched instrument appears, at least occasionally, to do. I found the brise-coque to cause less inconvenience and pain; and I further consider it *peculiarly* useful for the destruction of fragment. In conclusion, I have only to add, that my friend Dr. Hume was present at every operation excepting the first, and constantly visited me in the intervals. Mr. Biggs, the Baron's pupil, witnessed every operation, and Dr. Boyton and Dr. Clendinning were present on several occasions.

ANDREW CASTLE,
Licen. Royal Coll. Surg. in Ireland, and
Surg. of the County Clare Infirmary.

The histories of medical cases drawn up by the patients themselves being generally the most interesting, I have thought I could not do better than copy, for publication, Mr. Castle's autograph. The opinion of this scientific practitioner cannot fail to have weight, for what person can be so capable

sitting was perfectly ineffectual, for the stone being twice taken, escaped each time, and I finally withdrew the instrument.

Seeing that in *five* different *seances* I had scarcely obtained twenty or thirty grains of detritus, and that the stone constantly slipped away, I had recourse to the brise-coque. The stone was of middling size and evidently flat, and this was proved at once by the fragment expelled, and afterwards by the frequency of the stone's escape from the claws of the three-branch instrument.

In three *seances* with the brise-coque, which, according to Mr. Castle, produced less pain than the first instrument, I completely destroyed the stone. It was immediately with facility seized, and destroyed with remarkable celerity.

One of the first obstacles which opposed themselves to the progress of lithotrixy, was the difficulty of demonstrating by comparative operations, the pre-eminence of one instrument over another. All the cases which contribute to enlighten opinion on this subject are valuable, and that of Mr. Castle possesses great interest in this respect, for it establishes in an incontrovertible manner, the high importance of the brise-coque, and its superiority in the instances of flat stones, over the instruments of Drs. Leroy (*d'étoiles*) and Civiale.—Dr. H.

of judging of the efficiency and importance of any curative means, as a medical man who has himself undergone the process?

When Dr. Heurteoup arrived in England a year ago, the series of means which he put in practice for the destruction of urinary calculi, excited much admiration. I then took the liberty of declaring in this Journal, that this operation could not be performed with every chance of success, without adopting, in the attack upon the different kinds of calculi, the several instruments constructed by him.

The happy issue of Mr. Castle's case furnishes, like many others treated by my friend, a strong support to that assertion. Indeed, what can be more illustrative of the utility of this powerful mechanism than the present history? Constructed by its author to be employed in cases where the 'instrument à trois branches' or 'pèrce-pierre' could not be used, in consequence of the flat form of the calculi, its utility has been here manifested in its fullest extent. We have seen that to effectuate the cure of Mr. Castle it was necessary to have recourse five times to the *pèrce-pierre* of the Chevalier Leroy, and that at each sitting the stone was captured three or four times but escaped as frequently. That on the first and third *seance* the perforator reached the stone but once only, and once again in an incomplete manner, and that until the five sittings scarce thirty grains of sabulous discharge were produced. After these sterile attempts Dr. Heurteoup employed the brise-coque, and in three *seances*, which lasted but three or four minutes, a large quantity of detritus is ejaculated, and a speedy cure succeeds.

Besides the great force which the brise-coque exerts in the destruction of these bodies, its action is much more rapid than that of the *pèrce-pierre*. This arises from the wonderful facility with which the movements of the new instrument are executed, as well in seizing as in breaking the stone. To be convinced of this, it is only necessary to see the Baron operate with the two instruments, a sitting with the "instrument à trois branches" continues as long as twelve or fifteen minutes, which is a long time.

The frequent occurrences of flat stones will explain to us why certain operations undertaken by other surgeons with the aid of Leroy or Civiale's instrument, are so tedious, occasion so much suffering to the patient, and are rarely attended with perfect success. Such operations ought, indeed, to be considered only as exercises, and their results cannot be admitted to indicate the quantity of good which humanity may derive from this new province of surgery.