elimination is a necessary procedure. In all cases it is well to examine the urine, noting any change from normal; sugar especially should be looked for. Some cases of carbohydrate intolerance, while showing no sugar in the urine, present quite constantly an increased amount of sugar in the blood.

Other patients are intolerant of sodium chlorid in large quantities, and questioning will often bring out the fact that these patients indulge overmuch in the use of ordinary table salt. Still other patients eat excessive quantities of nitrogenous articles of food, and a diminution in this element is of distinct advantage.

Especially in persons past middle life suffering from eczema, nephritic changes should be looked for, and if present, properly treated. Certain other patients with eczema present an anaphylactoid if not a true anaphylactic reaction against specific proteins or proteins of certain groups, among which may be mentioned egg albumin, pork, and the protein of cow's milk. It would lead too far ahead to enter on a discussion of the various proteins which, in susceptible persons, may cause an outbreak of eczema, but the careful clinician will bear these possibilities in mind and will inquire into the dietary habits of his patients. Often he will find that the patient himself may connect the ingestion of certain articles of diet with the appearance of the eruption. A patient recently under my care reacted in a marked way to the ingestion of walnuts. Besides the eczematous condition which followed he also presented marked edema of the nasal and conjunctival mucosae. It is well, therefore, to start a patient off with an absolutely bland diet, which is readily digestible and which contains little or none of the substances that experience has shown may be productive of eczema.

There are undoubtedly patients in whom the predisposition to eczema can in a greater or less degree be controlled by organotherapy. I have repeatedly seen patients materially aided by small doses of thyroid substance, and several neurotic and underdeveloped young women, I am sure were markedly benefited by the use of corpus luteum extract. However, the physiology and therapeutic uses of the ductless gland substances are in general so poorly understood that it is impossible to give clear-cut indications for their use, and the suggestion is offered merely as one that they may sometimes be helpful in the management of eczema.

25 East Washington Street.

Infant Mortality in War Zone.—A report on the subject of the sufferings of French and Belgian children under German domination in the occupied regions has been made to the French Academy of Medicine by Dr. Albert Calmette. It was found that out of 18,036 children who at the time of the armistice were attending the public or private schools of Lille, rather more than 8,000 had to be sent to hospital or convalescent colonies by the ministry of the invaded regions. This selection, made by specially qualified doctors, established that in all the scholastic groups at the end of the occupation the growth of 60 per cent. of the children had been definitely arrested, while about 40 per cent. showed clear signs of tuberculosis of the lymph glands. In a single school (Ecole Fombelle) inspected in March, 1919, of 210 children examined only one was normal; 163 suffered from deformity or other disease; 139 had swollen glands; forty-two had rickets, and six had pulmonary tuberculosis.—*Med. Officer*, Nov. 29, 1919.

Clinical Notes, Suggestions, and New Instruments

BLOOD TRANSFUSION APPARATUS

L. L. STANLEY, M.D., SAN QUENTIN, CALIF.

This blood transfusion apparatus was devised and has been successfully used at the California State Prison, at San Quentin, Calif. It is of the ball valve plan, and may be used with any capacity Luer syringe.

Fig. 1.—Details of blood transfusion apparatus.

When the plunger of the syringe is drawn out, the ball A (Fig. 1) engages in the socket, not permitting any fluid to pass it, while the ball C is displaced upward to position B, allowing the blood to come from the donor into the syringe. When the syringe piston is pushed in, the ball in the lower chamber engages the socket at C, preventing the fluid from returning to the donor. At the same time the ball A is released from its socket and assumes a position in the upper chamber at D, allowing the blood forced from the syringe to flow into the veins of the recipient.

Fig. 2.—Blood transfusion apparatus and buret.

The donor and the recipient are placed on adjoining tables, with the arm of the recipient slightly higher than that of the donor, for the balls engage by gravity, and the valve must be held in a vertical position. The syringe is thoroughly sterilized, and the inside is coated with Ablemann's citrated ointment, which consists of:

	Gm. or C.c.
Adeps lanae	10
Aqua	10
Sodium citrate	10
Petrolatum	q. s. ad 100

The valve connections and needles are boiled together.

With the syringe connected, enough physiologic sodium chlorid solution is drawn in to displace any air, and leave a

few cubic centimeters in the barrel. The cubital veins of both parties are located, and tourniquets applied. The needle is placed first into the vein of the donor, and when blood begins to flow, is connected to the apparatus at *E*. Similar procedure is done with the recipient, and when the connection is made at *F*, the operator may pump the blood.

The apparatus is so arranged that it may be taken apart and thoroughly cleaned. It is well to have two of these valves, one to quickly replace the other, in case the syringe should begin to work hard, or the blood begin to clot. The first valve is easily disconnected at *E*, *F* and *G*, and the second one substituted.

Should it be desirable to give citrate solution with the blood, a buret (*I*, Fig. 2), to which is attached a rubber tubing with a No. 28 hypodermic needle at the end, is provided. When the apparatus is connected up, the needle is plunged into the connecting tube at *H*. As the blood is drawn in from the donor, likewise the citrate solution is taken in from the container *I*. The percentage of the mixture is regulated by the size of the needles at *H* and *E*, and the strength of the citrate solution.

The advantages of the apparatus are:

1. The amount of blood transfused can be measured.
2. The blood is not exposed to the air.
3. The method is direct.
4. Only veins are used, and repeated transfusions from the same donor may be made.
5. Any amount can be transfused, and done very quickly.

In this prison, many injections of arsphenamin are given each week. When the patients are ready, sufficient solution is prepared for ten consecutive injections.

With this apparatus, using a 50 c.c. Luer syringe, one dose is drawn from the mixing vessel through the donor's tube, and injected into the patient, through the recipient's tube. This is more satisfactory than the gravity method, and saves considerable time.

UNIQUE FOREIGN BODY IN NOSE*

IRVING WILSON VOORHEES, M.S., M.D., NEW YORK

N. A. W., a colored man, aged 34, a native of the British West Indies, came to the West Side Dispensary, Jan. 17, 1920, with a history of yellow discharge from the right nostril during the previous two months. Examination revealed a large black mass in the right nasal fossa, very firm and hard, but freely movable in its bed. The septum was pushed over to the left, and the mass was bathed in yellow pus. It seemed reasonable to suppose that we had to do with a mass of necrotic bone, probably a sequestrum from the nasosphenoidal wall following a tertiary lesion. Pledgets of cotton saturated with 10 per cent. cocaine were introduced into the nose and left for ten minutes, at the end of which time our diagnostic acumen received a rude shock.

When 17 years of age, the patient tried to invent a gun. He took a piece of bicycle pipe about 1 foot long, into the open end of which he screwed a threaded bolt borrowed for the occasion from some part of a bedstead. He then bored a touch-hole in front of this breech plug. Into the muzzle he loaded a charge of gunpowder, rammed it down, followed this with a few BB shots, which were also rammed down, and then invited his cousin to apply a flame to the touch-hole while he held the apparatus in the position of aiming. The charge exploded and so did the piece of pipe. The breech plug blew backward and had never been seen or found, but small pieces of iron penetrated the left side of the nose in the region of the ethmoid. The scar from these fragments is still quite visible but insignificant. The boy was in bed for seventeen days, and was visited twice daily by a physician, who did not, however, locate the piece of iron. For seven months thereafter the boy experienced pain in his head and between the eyes about 10 o'clock each morning. Considerable discharge came from his nose. This was worse, he thinks, in summer. He came to the clinic chiefly because his friends chided him about a stench from his head, and because of "frequent colds."

*Patient and specimen presented before the Nose and Throat Section, New York Academy of Medicine, Jan. 21, 1920.

After removal of the cocaine pledgets, the black mass above mentioned was grasped with strong nasal grasping forceps, and much to our surprise a solid heavy object was gradually withdrawn from the nose, occasioning only slight discomfort in its passage. When washed off under the water tap, the improvised "breech plug" was identified as the missing iron bolt which seventeen years previously had been suddenly projected into the patient's nose.

The dimensions and weight of the bolt were: length 5.5 cm. (2½ inches); diameter, 1.5 cm. (½ inch); weight, a little over 1 ounce.

There is probably some latent sinus trouble, which we intend to look into as soon as roentgenograms can be made.

It seems hardly possible that any human being could carry such a foreign body in any part of his anatomy for seventeen years and not be conscious of its presence.

Special Article

TYPHOID IN THE LARGE CITIES OF THE UNITED STATES IN 1919

EIGHTH ANNUAL REPORT

THE JOURNAL presents its eighth annual survey¹ of typhoid fever mortality in cities in the United States having more than 100,000 population. The cities included in the summary number sixty, and are the same as those reported on last year.²

It should be especially noted that the calculated typhoid rates for 1919 presented in this report are based on relatively unsatisfactory population estimates. It is now nine years since the 1910 census, and probably in a number of instances the population estimates are quite wide of the mark. The U. S. Census Bureau makes no estimate at all for the population of Denver, Memphis, Portland, Ore., Seattle and Spokane. Special methods of estimation have been used by the Census Bureau for the population of Cleveland and Washington.

The margin of error is probably not great in a few cities, such as Albany, N. Y., but in many others it will doubtless be shown by the 1920 census enumeration that calculated death rates for 1919 will have to be considerably revised. For these reasons, typhoid death rates can be subjected only to a broadly critical comparison this year.

The nine largest cities in the country (Group 1) show a very notable improvement in typhoid death rates as compared with 1918. For the first time, all the cities in this group have a rate below 10. In only one instance (Philadelphia) are the 1919 figures higher than those for the preceding year. Baltimore has the lowest typhoid rate in its history, and rises into the first rank (cities having a death rate of less than 10). The improvement in Baltimore is all the more noteworthy in that the recently annexed territory has had a much higher typhoid morbidity than the older portion of the city. It is stated that for the first

1. The preceding articles were published, May 31, 1913, p. 1702; May 9, 1914, p. 1473; April 17, 1915, p. 1322; April 22, 1916, p. 1305; March 17, 1917, p. 845; March 16, 1918, p. 777, and April 5, 1919, p. 997.

2. The number of typhoid deaths has been sent us by the local officer of health, and the rates have been calculated on the basis of population estimates made according to the method of the U. S. Census Bureau. It may perhaps be noted that the figures kindly furnished us by the municipal officials include the deaths of nonresidents as well as residents occurring within the city limits. In some instances this undoubtedly gives an exaggerated impression of the amount of typhoid fever in a community, but at present statisticians are agreed that "the attempt to eliminate the deaths of nonresidents would often result in an understatement of the true mortality" (Bureau of the Census, Mortality Statistics, 1912, p. 13).