pulse and temperature were concerned were prompt and good, but the greatest improvement noticed was in the early disappearance of pain, vomiting and nausea, of tenseness, tenderness and distension over the abdomen and of the restlessness and anxiety so painfully evident in every movement and look. She recovered and is now about ready for an interval operation which offers her every possible prospect for health and longevity.

I must add just a few words condemning two drugs frequently prescribed in peritonitis: 1. I desire to endorse absolutely all Ochsner says about the harmful effect of salts in this disease and, judging others by myself, I take it that the habit of starting the treatment of these conditions with salts will be the hardest lesson the physician will have to unlearn, for that plan is part and parcel of the fixed routine of 95 physicians in 100. It will be even more difficult to impress the laity with the importance of leaving off the salts. 2. Opium, as frequently ordered, is also to be condemned. It does arrest peristalsis, to be sure, and some of you will say that it favors the very objects I seek; but unfortunately it does much more and in those respects its effects are so supremely injurious that the sum of its action is far from good. The point of this matter is, however, in the fact that, if the lavage and absolute fasting are maintained, both the salts and the opium are quite unnecessary, as the intestinal and stomach contents are removed and the peristalsis arrested by a better, simpler and far safer proceeding. This does not preclude the moderate use of morphin and codein hypodermically for the relief of pain, anxiety and restlessness in the early stages of the disease and its treatment, but if the lavage, fasting and avoidance of cathartics is strictly maintained the necessity for anodynes is soon past.

SUMMARY.

1. The treatment of septic peritonitis by surgical measures on the one hand or by rational medical measures on the other must be determined in each case first through a carefully-made differential diagnosis as to the source of the infection; second, through an estimate of the apparent virulence and diffusion of the infection, and third, through consideration of the elapsed time since the infecting material was injected into the cavity.

2. Infections from the intestines, vermiform appendix and oviduct are apt to be dangerous in the order given, because of the relatively greater mobility of the first and then the second and third, and of the greater relative distribution of their contents, which favors and promotes diffusion of infection and prevents plastic closure of the leaking viscus.

3. Physiologic rest of the intestines favors limitation of the area of peritoneal involvement and the saving of life, and is best attained by the maintenance of an empty bowel and the avoidance of cathartics; this means lavage, fasting and no salts.

If enemata are used for clearing the rectum, as may be required, they must be small in amount and injected with the utmost care from a low bag.

4. Cases of diffuse peritonitis with profound local and general symptoms must be regarded as being inoperable, inasmuch as the interests of the patient and the requirements of good surgery both demand that untimely surgical measures be not employed where better results may be had with simpler and safer ones.

5. Early operation, before diffusion has taken place, is safe and in most instances to be advised if the environment be favorable and an experienced surgeon at hand. Interval operations are certainly advisable after recovery from a diffuse peritonitis from any source.

THE NEUROSES OF THE HEART.*

HERMAN H. HOPPE, M.D.

CINCINNATI, OHIO.

In presenting for your consideration the subject of functional heart affection, I have a two-fold object in view: First, to illustrate by appropriate cases the undoubted forms of heart neurosis, and secondly, to call your attention to borderland conditions where we are frequently in a quandary as to whether the case would be classified as myocarditis or whether the cause lies in a functional disturbance of the nervous mechanism of the heart's action.

The subject of myocarditis and its clinical phenomena have lately been the subject of much discussion and the chronic form at least is of much interest to us, especially in our office practice where it often is important to know, and difficult to determine whether a given irregularity of the heart is of organic or functional origin. In the same way, when diseases of the muscular walls or of the valves do not enter into consideration at all, it is often impossible to determine whether the nervous derangement of the heart's rhythm is of an organic or functional character. If we omit from our consideration the subject of fevers and direct chemical poisons affecting the heart, we will render the subject more circumscribed and at the same time not detract from the interest in the matter. Various factors are essential to a normal condition of the heart's action. The muscles must be sound, the valves normal, the nervous mechanism perfect, and the heart must be in its usual position. When any one of these conditions is altered, the heart's action becomes changed, irregular, rapid, slow, tumultuous.

A nervous affection of the heart is caused by a faulty condition of the nervous system locally in the heart, the nerves leading to the heart, their center in the medulla, or by what is more frequently the case, by a derangement of the nervous system as a whole, such as we see in neurasthenia and hysteria.

The nervous ganglion system of the heart may be the central point of the disease, and according to Krehl, although this is still in doubt, future investigation may confirm this idea. The nervous condition of the heart is surrounded, as a rule, by all of the general and special signs and symptoms of neurasthenia and hysteria, but sometimes it is the only symptom. In order to understand the neuroses of the heart it is necessary to give a brief survey of its physiology. Twenty years ago the physiology of the heart was looked upon as a closed chapter; the Traube theory was universally satisfactory. He taught the two-fold action of the heart: *a*, the muscular motor system, *b*, the regulator system, viz., the accelerator influence of the sympathetic and the depressor influence of the vagus. To say that the theory is overthrown, at least in part, is to give you old news. Gaskell, Woolbridge and Tigerstead were the first to show that the experiments on which muscular theory was based were faulty, but it remained for His, Jr., Romberg and Krehl to put forward and apparently substantiate the theory of the autonomy of the heart. Engelman in a complete paper gave the theory its present standing. The myogenic theory of the heart's action is critically reviewed in an admirable paper by Leyden. It teaches that the heart is practically one muscle fiber with the power of automatic action. The ganglion cells and all the nerve fibers of the heart are sensory in char-

* Read at the Cincinnati Academy of Medicine, Dec. 16, 1901.

acter. Carl Huber of Ann Arbor in an admirable paper which has the support of Waldeyer, tries to show that nerve fibers do enter and terminate in the nerve fibers of the heart by a small knob-like termination. But the consensus of opinion is against this theory. While the heart is rich in sensory fibers and ganglionic cells there are no motor fibers, no motor centers; the first decided blow to the neuron theory. The motor stimulus comes from the blood itself, perhaps from the chlorid of sodium alone. Kronecker, who had performed an experiment of pricking the center of the heart just below the aortic valves and thereby throwing the former into an irregular and tumultuous action, and finally causing it to stop altogether, seems to have found a motor center. But the most careful investigation, however, failed to reveal any accumulation of cells or nerves in this locality, and as long as ten years ago at the Berlin International Congress he explained the result by the irritation of the vasomotor nerves of the coronary artery. While the rhythm of the heart's action is automatic, this can be increased or diminished by influences, chemical, physical, or psychical in character, coming either through the sympathetic nerve or from the vagus.

We can divide the neuroses of the heart into three divisions: 1, neurasthenia cordis; 2, tachycardia; 3, bradycardia.

NEURASTHENIA CORDIS.

Under the head of neurasthenia of the heart various manifestations are seen. We have: *a*, typical neurasthenia of the heart, *b*, arrhythmia of the heart, *c*, pseudo angina pectoris, *d*, nervous bruit. Of these conditions, the neurasthenic heart is the best known and the most frequently encountered. This condition of the heart rarely occurs as the only sign of the neurasthenia. It is the local manifestation of a general condition. The neurasthenic condition being present, no matter what the cause, the local phenomena on the part of the heart are usually the product of the psychic influences. The most frequent condition which we see is the cardiac hypochondria, fear of heart disease. It is usually based upon the local precordial sensation produced by palpitation, irregularity or pain. Fear is the predominant psychic element of neurasthenia and the fear of heart disease is often combined with a marked disinclination to examination for fear of having apprehensions confirmed by the physician.

Palpitation, while frequent, is not an invariable sign of a neurasthenic heart. Its origin is often chemical or mechanical, due to displacement by a distended stomach. But when it is accompanied by a sense of fear and anxiety, I should say that it is at least associated with a general neurasthenic condition. Palpitation may occur without any cause whatever, often in the middle of sleep. It is associated with precordial distress, varying in degree from a slight pressure to pain so crucial and severe as to simulate angina pectoris. At the same time there is great mental distress, a pallid, moist countenance, and facial expression of great anxiety. At times the palpitation and precordial anguish is associated with a sense of constriction in the throat, a sense of strangulation which makes the patient jump from the bed, often from a deep sleep, into the middle of a room, uttering guttural noises or rushing to a window and throwing up the sash for air. These attacks come on suddenly and last from a few minutes to an hour or more. These paroxysms are only one phase of the neurasthenic heart. In some cases the heart is constantly twenty to thirty beats above the normal and

there is a sense of dissatisfaction with the breath taken, a constant effort to breathe deeper and a feeling that relief would be attained if a deeper inspiration could be made. This dyspnea may last for hours, days, or even weeks, the only relief being obtained by sleep.

The typical neurasthenic heart is weak, irritable and easily exhausted. When at rest the pulse is normal, but the slightest exercise, such as stepping up two or three steps to a platform, will raise the pulse rate from twenty to fifty beats per minute without any discomfort to the patient. In less than a minute after the patient has come to a halt, the pulse rate returns almost to normal. The same result is often produced by psychic influences, by coming into a physician's office and being examined. The sphygmographic tracing will show this weakness and irritability of the heart. The worst case of neurasthenia that ever came under my observation was a blacksmith, who worked daily at the forge for a whole year while he was under treatment. His pulse was so irritable, weak and rapid that the mere walking into the clinic in the presence of the students raised his pulse rate to 120 or more. This shows that there are exceptions to the rule. Just as irritable weakness, whether in the mental or physical sphere, is the characteristic of neurasthenia, so it is also of the neurasthenic heart. Krehl raises the question whether nervous affection can produce changes in the size of the heart. If so, how often and to what extent, and can it produce hypertrophy or dilatation? We might answer at once that when hypertrophy and dilatation are present there is an organic heart lesion, a diseased condition of the muscle itself. At present, however, there can be no certainty upon this point, principally because autopsies in nervous subjects are exceedingly rare. Krehl says that he has seen dilatation of the heart especially in masturbators and that with a cessation of the habit the dilatation has disappeared. He has also seen dilatation occur in ordinary neurasthenic paroxysms and disappear with the cessation of the attacks. Theoretically the explanation is very plausible. It is well known that an irritation of the vagus nerve has marked influence on the tone of the heart muscle. Why should not psychic influences have the same effect, the dilatation being the result of diminution of tone. Krehl is not so positive about the occurrence of hypertrophy; he thinks that he has seen it, but does not wish to place himself on record because of the difficulty of diagnosing small degrees of hypertrophy. Before the diagnosis of dilatation or hypertrophy can be made in neurosis of the heart, we must carefully exclude all organic lesions and must likewise bear in mind that we may have at the same time both an organic lesion and a neurosis. It is well to remember, however, that we may have as a result of simple neurosis of the heart undoubted dilatation and perhaps hypertrophy.

ARRHYTHMIA CORDIS.

In palpitation of the heart under the influence of psychic emotion, there is frequently not only an increase in the rhythm but there is a decided irregularity both in the rapidity of the action as well as in the occurrence of intermissions, and when the heart quiets down again we have a regular rhythmic pulse. In arrhythmia, on the other hand, we have a pulse that is irregular in rhythm, as well as intermittent, in the absence of all excitement, psychical or physical. Kronecker's experiment seemed to indicate that the cause of the irregularity in nervous patients may be due to some vasomotor disturbance of

the coronary circulation. Directly or indirectly, no matter how, the cerebrospinal system regulates the force and frequency of the heart, but the rhythm is the peculiar inherent automatic quality of the heart muscles and nerves. The most frequent cause of arrhythmia therefore is either a diseased condition of the coronary arteries or myocarditis. There can, however, be no doubt that we frequently find arrhythmia as a functional affection of the heart. Husley has shown that even in apparent health, variations not at all physiological occur, but marked arrhythmia to the point of intermitting is always pathological. The nervous heart may be intermittent; we may have a pulse bi-geminus, or trigeminus, or delirium cordis. All of the cases that I have seen have been neurasthenia with marked vasomotor disturbance. The great difficulty in neurasthenia lies in excluding myocarditis or organic nervous trouble. Certain cases had been diagnosed as Graves' disease but this was held to be untenable because of the irregularity of the heart, and of the fact that at times the heart's action was reduced to 80 beats or less per minute which never occurs in the former affection. The sphygmographic tracing of Graves' disease with a pulse rate of 130 will illustrate the difference in the two conditions. The rhythm of the heart is always perfect in Graves' disease. Myocarditis and disease of the coronary artery must be ruled out and this can only be done with great difficulty, especially in the start. We must rule out all acute diseases which might have caused myocarditis and arteriosclerosis.

Normal pulse.

Normal pulse, case of brain exhaustion.

Krehl lays much stress on allorhythmia, viz., an intermittance every fourth, sixth or eighth beat, recurring regularly. The intermittence of myocarditis occurs with great irregularity. In nervous arrhythmia there is a marked consciousness of the irregularity, which is usually absent in organic heart trouble. Arrhythmia of nervous origin sometimes disappears under excitement, while bodily exercise invariably increases the irregularity in myocarditis. Myocarditis is, as a rule, associated either with hypertrophy or dilatation, whereas in arrhythmia it is the exception. We must remember, however, that patients with myocarditis may have neurasthenia, and very often the same cause which produced the neurasthenia may also produce a myocarditis, e. g., influenza. In cases of this kind, time alone will clear up the doubt. Neurasthenic patients may have arteriosclerosis and this may complicate the diagnosis. Each and every symptom and sign should be carefully weighed in the balance and even then we may remain in doubt as to the exact nature of the arrhythmia. We can see the importance of a careful diagnosis by the bearing which it has on the treatment, for in neurasthenia an outdoor life is demanded and a gradually increasing amount of exercise, which may prove fatal to the case of sclerosis of the coronary artery or myocarditis.

NERVOUS ANGINA.

The nervous ganglia and nerves of the heart are, according to present teaching, purely sensory and therefore we may have in the heart, in addition to the other disturbances of sensation, distinct pain. This pain may

occur in paroxysm, simulating neuralgia, or it may be more or less constant with periodic exacerbations lasting from minutes to hours. This pain now and then occurs independently of any organic heart trouble and is known as nervous angina pectoris. There is no better authority than Romberg and he holds, in view of the many negative autopsies, that angina pectoris is always a neuralgia of the heart, that aortic stenosis and sclerosis of coronary arteries are merely contributing causes. There is even much dispute as to the seat of pain, the vagus, sympathetic, phrenic, intercostal nerves and the spinal cord having been looked upon as the seat of pain by undoubted authorities. I have seen one case of mild alcoholic brachial neuritis of the left arm which was associated with periodic attacks of precordial pain, radiating into the shoulders, which Oppenheim of Berlin saw with me and confirmed the diagnosis of pseudo angina pectoris. Prolonged rest and withdrawal of alcohol has up to the present time relieved the attacks. We are on very uncertain ground when we view the pathology of angina pectoris, and whether we hold the theory of coronary disease, of neuritis or neuralgia, we have undoubted authority to uphold us in our opinion. The most widely accepted view, however, is that true angina is caused by organic changes and most frequently by atheroma of the coronary artery. Angina pectoris, like tachycardia and bradycardia, is a symptom and not a disease. My object is not to treat of angina as such, but merely a pseudo-angina occurring in neurasthenia, which has a very close resemblance to the true form.

Whittaker says with truth that precordial pain, no matter how great its severity, does not constitute angina. We must have associated with it dyspnea, great anguish, mental anxiety and the fear of impending death. The cardiac crises of locomotor ataxia are purely neuralgic in character. We may have mild attacks of reflex angina arising from various disturbances of the liver, pelvic and gastro-intestinal canal. When angina is seen in alcoholic patients, I should hesitate to make the diagnosis of a purely functional neurosis. We may have a neuritis or even coronary arteriosclerosis. Tobacco, tea and coffee may produce attacks of pseudo-angina.

Since pain alone, its intensity, location and duration are the same for true angina as for the false we must look around for some differential point. True angina usually occurs in individuals who have other evidences of organic heart lesions, muscular or valvular. False angina occurs in individuals who are neurasthenic or hysterical. In the attack this point will not avail much, but outside the attack, in the interval, it matters a great deal.

Huchard gives the following points: First, every angina produced by an effort of any kind is true angina. Second, every angina which occurs spontaneously is false angina. Third, an angina which occurs in the night, though independent of effort, is a false angina. Nervous patients are not quiet during an attack. They walk up and down from chair to lounge, from lounge to bed, wringing their hands, lamenting and crying. The attack might be associated with other signs of neurasthenia.

thenia, as palpitation of the heart. Areas of anesthesia may be present. Pain may affect the whole left side. Neurasthenic cases are often associated with typical neurasthenic breathing. Most functional, reflex or neurotic anginas occur in young individuals; true angina in the old. Hysterical attacks of angina are often associated with typical hysterical mania, in which delusion of the senses of sight are very frequent, always being absent in true angina.

BRUIT AS A NEUROSIS.

Can a purely functional change in the rhythm of the heart produce a bruit? We see bruit frequently in the

Neurasthenia cordis.

various forms of anemia and in chorea and attribute it to the quality of the blood. But what ground have we for this assertion? If it be true, why do they not occur in all cases of anemia? Is it not just as probable that in these cases there is some disturbance of the mechanism of the heart, which produces the bruit or some anomaly in the muscular contraction which leads to a temporary faulty closure of the valves?

The following is a case of nervous bruit:

Mr. G., age 41, married, three children, has always been

Neurasthenia cordis.

healthy. No evidence of venereal disease, never sick, is applying for life insurance, was examined by Dr. Minor and myself at patient's office. On examination of the heart, a loud bruit was heard, even before the ear or the stethoscope was applied to the chest. It could be heard at least an inch from the chest. The heart was found normal in size and location. Its action was strong, somewhat rapid, pulse 100. Second sound slightly accentuated. The examination in other respects perfectly normal. Patient seemed only slightly excited. Urinalysis negative. Insurance refused. Second examination made in my office.

Neurasthenia cordis.

Heart perfectly normal in size and sound. No bruit. Patient attributed the disturbance at the examination twenty-four hours before to a dinner which he had attended on the night previous. On demand of applicant, the medical director of the insurance company came from New York and a joint examination was made after a lapse of ten days, the heart was again found normal, and the insurance granted in full amount.

It is difficult to explain, and we have not the time to go into details concerning the numerous theories that have been advanced, to explain the nervous derangement of the mechanism of the heart action.

Krchl, Martius and others look upon the result of

physiological experiments on animals and their application to the human heart with much distrust, not to say contempt. Very little reliance is placed by them upon the direct influence of the vagus and sympathetic nerves in the production of derangements of the heart's action. The whole subject is still in obscurity, but they seem to lay more weight upon the ganglionic system of the heart itself as the seat of all disturbances of action. Barker believes that the experimental and fetal researches of His, Jr., and Romberg have found a satisfactory anatomical basis of the physiology of the nerves of the heart and a starting point whence perhaps those puzzling clinical problems in connection with the cardiac neurosis may be advantageously approached. His, Jr., in his illustration of the ganglionic system of the heart, has likewise shown a wonderful blood supply of the ganglionic system. "Every ganglion is surrounded (as Whittaker so beautifully puts it) by a thick network

Neurasthenia cordis.

of small blood vessels and numerous branches penetrating into its interior. Thus the heart ganglia are in no sense degraded by being denied the office of presiding over the motion of the heart. They have the higher one of perceiving the first influence of failing nutrition or toxic impression."

The neuroses of the heart are, as a rule, found in those forms of neurasthenia in which the vasomotor disturbances are most common. Patients with cold, moist hands and feet, with bluish-red discoloration of the skin of the extremities, the tache spinal, the pallid, easily flushed faces, are the ones who most commonly have the cardiac disturbances. It is not difficult to understand that if the arteries and veins surrounding the cardiac plexus are in the same state of passive distension and sluggish activity, how readily all the neuroses of the

Neurasthenia cordis.

heart can be explained on the ground of a deficient nutrition, producing a changed activity of the cardiac ganglia.

The prognosis and treatment of neurosis of the heart are essentially the same as in neurasthenia, the discussion of which we will not be able to enter here.

TACHYCARDIA.

Nothnagel, Bouveret and Martius have made known the symptom complex, which we designate as essential tachycardia. Tachycardia is a symptom of various affections and not a disease; it occurs in a moderate degree in various organic heart affections; in any disease of the neck or mediastinum compressing the vagus; in central disease of the medulla and in poisonings. But it occurs also under circumstances in which it is the essential symptom of a paroxysmal attack, which has come to be recognized as a distinct affection occurring periodically in individuals otherwise well. In its true form

it has to be encountered but once to dispel any doubt as to its right to a separate classification. Real tachycardia occurs in individuals who are, to all intents and purposes, well between attacks. In others there has been found a nervous diathesis without there being present any marked hysteria or neurasthenia. The attack comes on suddenly, the individual feels a sensation of a sudden start or a darting pain in the chest, then a sense of fatigue, fear and anxiety. He goes to bed and lies perfectly quiet, but when the attack lasts a long time, there is more or less restlessness. Oppenheim, however, reports one case with exceedingly high pulse, the patient being able to go about his business and even walk for miles in an attack. The main symptom is the pulse and the pulse rate. I should say from the few cases that I have seen, that in genuine cases the pulse rate is never below 160, usually between 180 and 220 and Krehl says, as high as 300 per minute. The pulse is weak,

Case of obesity in young woman—no neurasthenia.

feeble and small. The heart's action is feeble and the impact against chest wall scarcely perceptible. Binswanger, however, reports a case with a pulse rate of 150, pulse hard and with strong impact against the chest wall. The heart sounds are clear, but so rapid as to be almost indistinguishable. There is no bruit. The presence or absence of dilatation of the heart is a much discussed question. I have seen one case with acute dilatation of the heart in which there were all the symptoms of the typical paroxysmal attack of tachycardia. In the other two cases reported below, I was unable to find any dilatation. Martius claims that dilatation is always present. Krehl says that it is not and that he has seen cases in which it was not present. There seems to be an agreement among authorities, however, that muscular weakness and loss of muscular tone is an

Pulse tracing one hour after admission. October 10.

essential condition of the attack and Krehl argues that if this is the case, theoretically at least, it would not be improper to imagine that an acute dilatation does occur and that while it may be hard to detect by percussion, it is very important to bear this in mind in making a prognosis as to the individual attack. The whole discussion, however, proves the first statement of this chapter, viz., tachycardia is a symptomatic manifestation of various conditions. The following three cases will illustrate the various conditions under which this may occur:

Miss L., age 29, single, dressmaker. Mother well, father dead, was nervous. Has one brother who is a periodic drinker, who is nervous and hysterical. Patient has always been well, has worked hard and steadily at her occupation, has always been of a nervous temperament. Present trouble began two years ago. The attack always begins with palpitation of the heart, a feeling of oppression on the chest, difficulty of breathing, great anxiety, restlessness, and much physical exhaustion. First attacks lasted but a short time, never more than a half

hour, returning at intervals of a month. Each succeeding attack was longer and more severe. During the last year it was always necessary to administer a hypodermic injection of apomorphia, which would put an end to the attack. The last few attacks have lasted from three days to a week, during which period patient was confined to bed. On Dec. 5, 1899, I was called to see this patient. The attack has already been in existence for three weeks. It began suddenly like all the previous attacks with the following symptoms: Sudden feeling of oppression in cardiac region with sense of great palpitation. Intense apprehension, restlessness, pallor and dyspnea. Unable to sleep, unable to eat, vomits frequently. The violent palpitation has stopped but twice in three weeks and only for a few hours at a time. Examination: Patient is very pale, lips white, eyes sunken, with fear, anxiety and restlessness on the face. Complaints of violent action of heart and the feeling of suffocation. Action can be seen through chest wall; the apex beat is normal; no bruit can be detected. Pulse is over 200 per minute and usually so rapid that it is impossible to count it. No hyper-

Taken five seconds later than preceding chart.

trophy or dilatation. Radial pulse very thin and rapid, not compressible. No enlargement of the liver, no disturbance of the peripheral circulation. In previous attacks hypodermic injections of apomorphia gave prompt relief, but in the present attack none. In the sixth week the patient had a sudden attack of right side hemiplegia with difficulty of pronunciation. In the course of a week the hemiplegia improved, but there was no change whatever in the tachycardia. On the seventh day after the onset of hemiplegia, the patient suddenly died. It will be seen that this was an extraordinary case of tachycardia. The early attacks seemed to be essential, but judging from the fatal termination, preceded by the hemiplegia, there must have been a bulbar lesion which produced both the tachycardia and the sudden death. No autopsy could be obtained.

CASE 2.—Infant, age 7 months. Parents are both healthy. Has one older sister and brother, both well. Has been well since birth. Three days ago had an attack of gastro-intestinal

Taken October 14.

disturbance. Vomiting and diarrhea. Bottle fed. Temperature 100, pulse 100. I was called at midnight. Child collapsed suddenly. The face is pale, the lips somewhat blue and cyanosed. Is unconscious. Eyes half open, respiration 48. Radial pulse at times absent, at others, faintly felt, very small, thready, easily compressed. Heart normal in size. Heart's action varies from 180 to a frequency at which it can not be counted. The liver is markedly enlarged. The body generally pale and limp. Salt water injection made hypodermically. Very little improvement. Injection of 1/180 gr. strychn. sulph. gave prompt relief. Child recovered consciousness, pulse becomes stronger, 120 per minute. In 24 hours a similar attack. This, however, continues 12 hours, the pulse being constantly 180 and over per minute. Child unconscious, in state of collapse. Tendency to convulsions, with a scarcely detectable radial pulse. During the next two days has two similar attacks of short duration with recovery. The treatment in this case consisted of many neutral saline injections with three doses of strychnia strophanthus 3/4 gr. every 3.

hours and ten doses of calomel gr. 1/20. Diagnosis, acute dilatation of heart.

CASE 3.—Mrs. K., age 58, married, four children, has had attacks of illness. For the past few months has been more nervous than usual. Has been under treatment for nervous indigestion. Complains of violent heart's action and a sense of prostration, great mental distress and fear of death. Her face is somewhat flushed and has an anxious expression. No pain. Pulse weak, small, thready, 190 per minute. No bruit, no enlargement. Gave seidlitz powder and potassium bromid gr. 30 every 3 hours. In four hours no relief. While sitting at bedside, without any apparent cause, the pulse suddenly dropped from 180 to 78 beats per minute. This change took place instantly. Has had in all three attacks. The second attack lasted twelve hours, the third, three hours. In each attack the relief came spontaneously. Last attack five years ago, since which time patient has been comparatively well, with the exception of more or less neurasthenia.

It is important in the individual case to determine the cause of the attack. Myocarditis is very often the cause

Case A.—Arrhythmia—irregular, violent, intermittent.

of tachycardia. In most cases there is some disease of the valves, coronary arteries or muscles, even dilatation or neuritis. While this is easily apparent on the autopsy table, it is very difficult to find during life. In essential tachycardia there is an absence of all indications of disease of the heart or nervous system in the intervals. In the attack we may find dilatation, but whether this is the cause or the effect is a matter of dispute. We must always remember the great difficulty that exists of diagnosing the slight deviation from the normal, either in the vessels or in the muscles of the heart. Martius says that all the attacks of tachycardia are the direct result of dilatation of the heart. Krehl takes a middle standpoint. The dilatation often follows the onset of the attack, may even be the cause, but is often absent. With Romberg he holds that the cause

When at perfect rest, pulse 72, irregular and intermittent.

is to be looked for either in a spasm or in a diseased condition of the coronary arteries, in a derangement of the nervous mechanism of the heart itself, and not of distant nervous system.

The diseased condition of the heart itself, viz., of the muscles, valves and arteries, can, at most, be only a predisposing cause, for, with the enormous frequency of the latter condition, tachycardia should be a very common instead of the rarest of cardiac affections.

Oppenheim has called attention to a group of symptoms which he says is due to irritation of the sympathetic nerves of the heart and of which tachycardia may be the most prominent symptom, viz., dilatation of the pupil on one side, wide separation of the eyeballs, pallor and reduction of the temperature of the face, with tachycardia. In making the differential diagnosis between essential tachycardia and that dependent upon disease, we must especially during the interval look for

all those signs and symptoms which may point to an organic lesion of the muscles of the heart, general arterial sclerosis, disease of the medulla oblongata or the occurrence of one of the acute infectious diseases which might have caused a neuritis of the vagus nerve. Whittaker says the irregularity of the pulse during the attack would be an indication of dilatation of the heart. Fraentzel says that tachycardia due to irritation of the sympathetic can be cut short by morphia and that produced by vagus affection by pressure on the vagus (Whittaker).

BRADYCARDIA.

This is exceedingly rare as a nervous affection. The bradus signifies slow—a slow heart. Bradycardia usually signifies a stenosis of the aortic valve, often complicated by myocarditis chronica or coronary artery sclerosis. There are at present two cases in the Cincinnati Hospital with pulses respectively of 32 and 36, both cases being of organic nature. In the clinical records

In paroxysms of palpitation, pulse goes as high as 140 per minute.

of the Cincinnati Hospital I have been able to find one case in the past ten years, also organic in character. A slow pulse of 60 or 64 is very rare, however. One of my patients who has been a neurasthenic for the past 20 years has a pulse habitually of 54 or 56. This patient has very marked attacks of palpitation and all the other manifestations of neurasthenia cordis, but only in paroxysmal attacks which occur at intervals of months, induced usually without cause, but at times by indigestion. The attack is invariably attended and followed by the fear of having organic heart disease, which is invariably dispelled by a formal examination and a positive statement to the contrary. Still I would not like to classify this case as one of bradycardia.

I have seen another case of slow heart in a neurasthenic with periods of intermittent pulse, rate 60, in

These last four tracings were taken at one sitting.

which no cause but a psychic one, the fear of heart disease, could be found. The intermittent pulse was present for a period of months. Tobacco was alleged as a cause, but its withdrawal or its continued use seemed to make no difference in the attack. We can also rule out myocarditis in this particular case. I have known him for seven years and he is still in good health with no evidence of organic lesion. I should look upon these two cases as unusual, because they occur in neurasthenia. Another patient now under my care for insomnia, fear of insanity and general nervous and physical exhaustion had a pulse of 54 for 10 days. At present it is normal.

Physiological bradycardia is seen often. Napoleon is said to have had a pulse of 40. Ruhlin, the pugilist, is said to have a pulse of 44, even after exercise. Bradycardia as a sign of organic heart disease or as a symptom of organic brain or cervical myelitis or injuries to the chord in the cervical region is not infre-

quent, but as a neurosis pure and simple it is exceedingly rare. Bradycardia is most frequently encountered in stenosis of the coronary artery. The bradycardia seen after the acute infectious diseases are probably due to mild myocarditis, while that always seen in conjunction with jaundice is probably due to the direct action of the circulating bile on the heart or on the nerve centers.

Reflex bradycardia is usually seen in gastro-intestinal disturbances and this is due to disturbance of the splanchnic circulation, throwing increased pressure and work on the heart. Attacks of bradycardia are described by nervous patients very frequently. These attacks are paroxysmal, not like tachycardia, but have more the character of palpitation or they may be associated with palpitation. The general signs are those of precordial pressure, anxiety, a sense of constriction, fear of suffocation and a subjective feeling as if the heart were beating very slowly and forcibly, combined with the fear of

Case of exophthalmic goiter, pulse 130, heart's action rapid, rhythm and strength normal.

its stopping altogether. The extremities become cold, the face pallid and covered with perspiration. In other cases there is distress in the stomach, nausea, retching, vomiting and subsequent collapse (Whittaker). The attack comes on suddenly and ceases suddenly. In milder cases there is simply abdominal distress, tenesmus, at times diarrhea. Sometimes bradycardia is associated with pain, which simulates very closely that of angina pectoris. Bradycardia is at times complicated with epileptiform convulsions. Whittaker says that these convulsions may be due to brain anemia, but he believes that the bradycardia and the epilepsy in the given case are both due to the same arrangement of the central nervous system. Before we can make the diagnosis of nervous bradycardia we must rule out all or-

Case of irritable, neurotic heart: pulse at rest, 80; under excitement, 130. Pulse regular and rhythmic, but action weak. Strong, muscular man.

ganic disease of brain, cord, heart or distant organs. The prognosis of functional bradycardia upon a neurasthenic brain is not unfavorable.

The treatment of bradycardia of a functional character is identical with that of neurasthenia. In the paroxysm we can use the various sedatives, bromids, asafetida and valerian.

BIBLIOGRAPHY.

- Krehl: *Specielle Pathologie und Therapie*, Nothnagel, vol. xv, 1. His and Romberg: *Fortschritte der Medicin*, 1890, S. 374. His and Romberg: *Transac. Int. Med. Congress*, Berlin, 1890. Krehl and Romberg: *Archiv für Experim. Pathologie*, xxx, S. 49. Leyden: *Deutsche Med. Wochenschrift*, 1898, No. 31. Martius: *Tachycardie*, Stuttgart, 1895. Bouveret: *Revue de Médecin*, 1889, p. 755. Romberg: *Herzkrankheiten*. Nothnagel: *Wiener Med. Blätter*, 1887. Whittaker: *American Practice of Medicine* (Wood), vol. iv. Kronecker: *Trans. Internat. Med. Congress*, 1890. Oppenheim: *Lehrbuch Der Nervenkrankheiten*, 1900.

ACUTE CONGESTIVE OR INFLAMMATORY GLAUCOMA.*

CHARLES J. KIPP, M.D.

NEWARK, N. J.

Not long ago I was asked by a medical practitioner to see an old lady who had for some days been suffering from a bilious attack, and who, since the day before, was nearly blind. He thought, as the eye seemed to be pushed somewhat out of the socket and she was blind, that she had probably during the retching and vomiting broken blood vessels in the orbit and in the eye. I saw the patient soon after and learned that she had been vomiting almost incessantly for two days, that she had great pain in the left side of the head, in the teeth and the eye. She was greatly prostrated. On examining her eye, I found that she could scarcely see the flame of a candle held within a few feet of her eye. The eye was very sensitive to light. The eyelids were somewhat swollen. The eyeball was slightly protruded from the orbit. The ocular conjunctiva was greatly ingested and swollen and a fold of it hung out of the palpebral fissure. The cornea was very cloudy and its epithelium was rough; touched with a piece of paper it seemed less sensitive than normal. The anterior chamber was very shallow, the iris swollen, and the pupil nearly double the size of that of the other non-affected eye. The opacity of the cornea prevented an examination of the fundus. A faint red reflex was all that could be seen. The eyeball was very hard, much harder than the other one. I made the diagnosis of acute inflammatory glaucoma, and advised an iridectomy as soon as possible. I instilled a few drops of the solution of the salicylate of eserine, which had, however, but little effect. The iridectomy was done under ether anesthesia a few hours later; the pain and the vomiting did not return after she came out of the ether, the wound healed rapidly, and three weeks after the operation the vision was restored to the former acuity.

I have related this case here, because cases of this group of acute glaucoma are most likely to fall in the hands of the family physician. The incessant vomiting, the violent pain in the head and the prostration are usually attributed by the patients to some stomach trouble, a bilious attack, or some disease of the head. The loss of vision, if noticed at all, and the redness and painfulness of the eye are frequently looked upon as a cold in the eye and not part of the general illness, and the family physician is sent for. I have seen a number of such cases, and I must admit that in some of these cases the physician took the same view of the eye trouble as the patient, and discovered his mistake only when the eye was totally blind and beyond help. In the majority of cases of acute glaucoma the onset is not as severe as in the case related. The pain is not as violent, vomiting may be entirely absent, and the headache may be much less severe. The outbreak is, moreover, in many cases preceded by attacks of cloudiness of vision lasting from a few minutes to several hours. The attacks come on at irregular intervals and are usually thought to be brought on by a sleepless night, an attack of indigestion, overuse of eyes in near work or some emotional disturbance. A restful night often restores the eye to its previous good condition, at least in the estimation of the patient, but a careful examination often shows some impairment of central vision, or slight contraction of visual field after the attack. Dur-

* Read before the Newark, N. J., Medical and Surgical Society, December, 1901.