And now, gentlemen, having given you such a history of this troublesome, and sometimes dangerous malady, as my scanty sources of information would admit of; and also a plain, but faithful detail of this case, I have but to beg of you to consider calmly, what is the value of the suggestion I have thrown out this evening. It may be said, that I am sanguine in my estimate of the remedy I propose. If I am so, it is for this reason: I found my patient in a critical, a perilous situation; her husband and friends looked to me to rescue her from that state: the ordinary means for relieving her had failed in my hands; and at one moment I felt, that not only was my professional character at stake,—for one false step would probably ruin my prospects for life,—but that the life or death of my patient depended, in a great degree, on the decision I came to. The idea of *inflating the pelvis*, flashed across my mind: having once conceived it, the execution of it was easy, and the result has been most favourable; for, from that moment she has not had one unpleasant symptom.

If, in the course of your practice, you find my suggestion of value to you, and that by its adoption, you are enabled to release poor suffering woman from the methods heretofore resorted to in the treatment of this affection, “many of which,” says Dr. Denman, “are severe, and some extremely cruel;” it will ever be a source of unfeigned happiness to me, that I have been thus enabled to bring it before you.

ART. IV.—*Contributions to Surgical Pathology.* By WILLIAM HENRY PORTER, Professor of Surgery in the Royal College of Surgeons in Ireland, Surgeon to the Meath Hospital and County of Dublin Infirmary, and Consulting Surgeon to the City of Dublin Hospital.

[Read at the Surgical Society of Ireland, 1st February, 1840.]

ANEURISM OF THE CAROTID ARTERY.

IN the details of the case which have given occasion to the following reflections, it may appear an almost fatal imperfection,

that I have not been able to verify the correctness of my views, by a *post mortem* examination of the parts engaged in the disease, and perhaps the omission may, in some respects, diminish the interest that would otherwise attach to it ; but although the case ended fatally, it was not in my power to procure permission to inspect the body, and the failure of my endeavours to obtain this object, is one of the principal reasons to induce me to lay it before the Profession. A length of time may elapse before I shall have an opportunity of seeing or treating an exactly similar case ; perhaps (as they are very rare) I might never again meet with an example, and therefore I am anxious to call the attention of my brethren to the subject, in order that some one else may complete that which I shall only seek the merit of having commenced. There is another circumstance that invests this case with a more than ordinary degree of interest. Although attended with several unpromising symptoms, I freely confess I undertook the operation with very sanguine expectations of success. After it had been performed, every thing seemed to progress very favourably ; even when the unpleasant occurrence of a suppurating sac made its appearance I was not much alarmed ; but when death took place, from a cause that I had scarcely anticipated, and but slightly dreaded, it at once led me to reflect most seriously on the circumstances that could render an operation, generally so successful, in this particular instance so entirely abortive. The case to which I allude was one of aneurism of the carotid artery—the chief point to be verified by dissection is, whether of the external or interval ? I believe it was of the latter, and the reasons on which that opinion is grounded will be found hereafter.

In considering the causes that may interfere with, or prevent the successful issue of an operation, and the recovery of a patient, it is rather an unusual view of the subject to regard the situation of the tumour as exercising a very decided influence, or to aver that the principle on which we treat an aneurism in the ham or in the thigh with almost a certainty of success, may not

be applicable to the management of a similar affection in a different locality. Yet such, I fear, is practically correct ; and although other collateral circumstances must be taken into account, as contributing their share in the production of this unpleasant result, yet I believe the internal carotid artery will be found so placed, as to afford a peculiar facility for the occurrence of these untoward complications, and that operations performed for the cure of aneurisms of this vessel, are attended with a degree of uncertainty scarcely to be expected elsewhere. In order to the correct understanding of this position, it will be necessary to say a few words on the manner in which an aneurism is cured by a ligature being placed on the trunk of the vessel, at the cardiac side of the tumour.

When such ligature is applied, it merely removes the impulse of the heart from the circulation through the vessel beyond it. The blood necessary to the nourishment of the limb or part at its distal side, is then conveyed thither by the anastomosing branches, which are generally too small to convey the impulse of the heart at the same time, and thus whatever blood enters the sac of the aneurism is no longer thrown in with a jet or *per saltum*. It thus can exercise no influence either on the fluid blood within the tumour or on the elastic qualities of the sac or its coverings : it is allowed to remain quiescent within its cavity, and no portion of it is forced back into the general circulation again. So circumstanced and at perfect rest, the natural tendency of the blood is to become coagulated, and the coagulum thus formed, being restrained by the sac and its coverings from extending itself in any other direction, is pushed towards the point where there is least resistance ; that is, against the open vessel which it compresses and causes to become obliterated, just as would happen to any bleeding artery under the continued pressure of a clot. In order, says Scarpa, that the compression may produce the union of the two opposite sides of an artery with each other, and at the same time the radical cure of the aneurism, it is necessary that, besides the vitality of the

coats of the artery, the degree of pressure applied upon it be such as to place the two opposite parietes of the injured vessel in firm and complete contact, and that it be at the same time capable of exciting the adhesive inflammation in the proper coats of the artery. Without the concurrence of these circumstances the compression does not prove beneficial, or only produces an incomplete cure ; for, whenever the compression is not sufficient to place the two opposite sides of the artery in complete and firm contact, and does not excite in them the adhesive inflammation, including also the root, properly speaking, of the aneurism, the canal of the lacerated or wounded artery remains open and pervious as before the use of the compression. If this view of the case be conceded, (as probably it will,) it must follow, that any thing capable of disturbing the blood within the sac, will interfere with the process of coagulation, and thereby delay, if it does not prevent, the cure ; and that any condition or position of the sac, which will allow the pressure of its contents to be directed otherwise than against the injured vessel, must have a similar effect. Either of these circumstances, taken singly, may prevent the cure ; when both are combined, success is scarcely to be hoped for.

When pulsation is observed to return in the aneurismal tumour in a short time after the application of the ligature, it is obvious that the great object of completely cutting off the impulse of the heart has not been attained ; that blood, to a greater or less quantity, not only enters the sac, but that some is returned again from it, and that the blood within it is constantly disturbed, and therefore kept in a state unfavourable to coagulation. This phenomenon of the re-appearance of pulsation, has of late years been frequently observed, and many of the causes that may conduce to it are now well known ; some of them being of a nature entirely to prevent the cure of the aneurism, such as the existence of two large trunks in the limb, by an irregular arterial distribution, or where one or more large branches arise from, or otherwise communicate with the sac, and others which

only delay the sanative process, but do not (at least necessarily) completely interfere with it. These latter are, firstly, the circumstance of the ligature being tied so loosely as not only not to stop the current of blood through the vessel, but not completely to cut off the impulse of the heart from the distal circulation; and what is of more importance, as bearing on the present subject, the existence of such an extensive and free communication by anastomosis as will convey to the tumour, by a circuitous route, the impulse which the ligature was intended to remove. I know not how far such a communication may be established in an aneurismal limb by a preternatural increase of size in the collateral vessels: such has been spoken of by authors, but I have no evidence of its existence in any one particular case to an extent sufficient to produce the phenomenon alluded to. However, I believe this unhappy condition obtains, with respect to aneurisms of the internal carotid artery in the neck, and that the free and extensive anastomoses through the vessels of the brain in their natural and normal state, will be quite sufficient, in some instances, as in the following case, to delay the process of cure, in others perhaps, to prevent it.

On the 21st of August, 1829, I performed the operation of securing the trunk of the common carotid artery of the right side in a woman of the name of Rourke, in the Meath Hospital: as this case has been already published in the fifth volume of the Dublin Hospital Reports, I shall merely remind the reader of some facts in connexion with it, which it may be necessary to refer to hereafter. It was a case of aneurism of the carotid of fifteen years' duration, consequently its growth had been extremely slow, and it might be reasonably inferred, that the aperture leading from the artery into the sac was very small; it was firm, hard, and solid, containing scarcely any fluid blood, and on examining the fauces, no pulsation could be observed within. The progress of this case was attended with some unpleasant consequences, such as a return of pulsation in the tumour in four hours after the operation,

and suppuration of the sac ; however, eventually my patient recovered, and left the hospital early in the following March to resume her former occupation as a servant. She had never been a very healthy person, and was afterwards frequently an inmate of the hospital for pectoral complaints : however, at length she died, on the 7th September, 1836 ; and I may mention, as a singular instance of strength of mind overcoming the prejudices so constantly met with in persons of her condition, she bequeathed her body to me, and by permission of the Inspector of Anatomy, it was sent to the School in Park-street.

On examining the neck, it was recollected that in the course of the operation, the sternal and a large portion of the clavicular attachments of the sterno-mastoid muscle had been divided, yet during the remainder of life, the patient never seemed to experience any inconvenience or imperfection of motion in consequence. The condition of this muscle, then, first attracted attention. At the inferior part of the neck, and for two inches above the clavicle, there was not a trace of fibre remaining of the muscle, and its place was supplied by a dense and strong fascia attached to the clavicle, below and into which the remnant of the muscle was inserted above. In short, the inferior portion of the muscle seemed to have been converted into this fascia ; on dividing it, the jugular vein, and the ligamentous-like substance into which the artery had degenerated, were exposed:

The remnant of the artery exhibited one continuous and unbroken cord from the bifurcation of the innominate to the division into internal and external carotids, so that, although the vessel must have been divided by the separation of the ligature, it had united again, and the exact spot at which it had been tied could not be ascertained. The internal carotid also was obliterated up to the spot where the ophthalmic artery was given off within the skull. In close connexion with this, was the remnant of the aneurismal sac, a small, firm, fibrous tumour, of an oblong shape, and nearly of the size and form of a very large almond ; it lay a little below the posterior belly of the digastric

muscle, and on the lingual nerve, to which it had some connexion, but not very intimate. The external carotid was pervious, but as compared with the vessel of the opposite side, greatly diminished in size, as were all its branches, excepting only the thyroid, which was pretty nearly of its natural dimensions. In consequence of the injection not being very perfect, the anastomoses were in general not minutely developed above the thyroid artery, but the communication between this and the branch ascending from the subclavian was extremely free, and the inosculations of these vessels within and on the surface of the thyroid gland very numerous, much more so than between the vessels of the opposite sides of the neck.

The subclavian on the right side was at least of twice the diameter of that on the left; the vertebral was enlarged in the same proportion; and the ascending branch of the thyroid was also increased in size. The chief external communication was between the ascending and descending thyroid arteries, which seemed to be the medium by which the external carotid and its branches were supplied: the quantity of blood formerly brought to the brain by the internal carotid, was afterwards furnished by the vertebral, which became enlarged throughout its whole course, until the formation of the basilar artery, after which the entire circulation of the brain was perfectly normal.

This dissection proves that in this case at least, the re-appearance of the pulsation after operation, was caused by the free and extensive anastomotic circulation through the brain, and that, without the slightest appreciable enlargement of the collateral vessels. The dissection was seen and examined by Dr. Hart, now Professor of Practical Anatomy in the Royal College of Surgeons in Ireland, and the preparation remains in the Museum of the School of Medicine and Surgery, Park-street. So far then, it has been shewn that the locality of the internal carotid is unfavourable to the cure of aneurism by operation, by at least, delaying its progress; I have now to shew how the same

influence may prevent it altogether. But previously, it may be advisable to mention the leading particulars of a case in which such an unfortunate failure actually took place.

Matthew Markey, æt. 38, of low stature, and very strong make, admitted into the Meath hospital on the 19th September, 1838, with a very large aneurism, occupying nearly the entire of the left side of the neck. It extended from about three quarters of an inch above the clavicle to the mastoid process, was bounded posteriorly by the trapezius muscle, and anteriorly it pushed the larynx considerably to the right side. The entire circumference of the neck over the most prominent part of the tumour, was fourteen inches and three-quarters: from the thyroid cartilage across the tumour, to the spinous process of the fourth cervical vertebra nine inches and a quarter: from the same points, the measurement on the opposite side amounted to but five inches and a half. Examined by the mouth, the appearances of the tumour were most alarming: the pulsation could be distinctly seen, and the blood almost felt under the mucous membrane; it seemed ready to give way, and burst into the mouth every moment, and so remarkable, and so urgent was this symptom, that on requesting my friend Dr. Graves, then the physician in attendance on the hospital, to examine this patient, I received a note from him, strongly pressing the necessity of immediate operation, lest such a catastrophe should take place. It is needless to detail the other symptoms; they were such as are usually observed, except that the tumour was very soft, the blood within it evidently fluid, and of course, the pulsation extremely violent. This peculiarity might, in some respects, be explained by the history of the case.

The disease may be said to have existed but a few days. Only five weeks had elapsed, since he first perceived a small hard tumour like a kernel, near the angle of the jaw, perfectly moveable, without pain, and (as he stated) without pulsation. In the course of ten or twelve days, it became uneasy, but not actually painful, and he poulticed it, in the expectation that it

would suppurate and break : it, however, increased in size, although slowly, and occasioned a good deal of annoyance in the motions of the head. It had then become distinctly pulsatile. Only seven days before admission, while at work, and after exerting himself considerably, he was suddenly attacked with most excruciating pain darting from the tumour across the forehead, and towards the vertex. He was immediately obliged to quit his employment and return home, where he discovered that the tumour had increased in size to a surprising extent, and that it pulsated with great violence. He suffered dreadfully for the next three nights, not sleeping, or even being able to lay down his head. He was then attacked with hoarseness, which amounting at times to nearly a total loss of voice, alarmed him so much as to cause him to apply at the hospital for relief, and he was admitted on the day above specified.

On the 22nd September, the operation of tying the common carotid was performed. As the space between the tumour, and the clavicle was extremely limited, I made a transverse incision at the root of the neck, parallel to, and above this bone, commencing internal to the superficial jugular vein, and extending forwards about two inches in length. Having thus exposed the sterno-mastoid muscle, I divided its clavicular attachment cautiously on a director, and came down on a very strong and resisting fascia, which, having slightly torn with the forceps, I also divided on the director. The edge of the sterno-hyoideus muscle could now be distinctly seen, which being carefully divided, I tore the sheath of the vessels partly with the nail of the fore-finger and partly with the end of the director. The vessel was now exposed, and although the wound was deep, I could easily pass the needle I generally use, and of which I have given a description on a former occasion, round the artery, which was tied as tightly as I could draw the ligature. During the operation I experienced no inconvenience from the jugular vein : I might almost say I never saw it. I certainly saw the pneumogastric nerve, because I looked carefully for it ; but the pleura

did not rise up in the neck, as I have experienced on other occasions. Altogether there was much less difficulty in the operation than might be anticipated ; the patient bore it well : was but twenty minutes on the table, and walked up stairs to his ward afterwards, refusing any assistance. On the ligature being tied, the usual phenomena occurred ; the pulsation ceased in the tumour : it became diminished in size ; and the patient declared himself relieved from pain.

I had mentioned in my clinical lecture on this case, that I anticipated a return of the pulsation in the tumour at an early period after the operation, and the suppuration of the sac at one more remote. In the former of these expectations I was disappointed ; pulsation did not return, although the tumour remained soft, and its contents evidently fluid. As to the full extent of this assertion, however, there may be some exception taken. Mr., now Sir Philip Crampton, who took a great interest in the case throughout, always said that he perceived a weak pulsatile thrill in the tumour ; and on looking at it in profile, I sometimes saw, or fancied I saw, a slight motion corresponding with the action of the heart, like that which might be exhibited by a large swelling in the immediate vicinity of an artery ; but on examining it with the hand, I never could feel a distinct pulsation ; if such existed, therefore, it must have been extremely weak and indistinct.

It is unnecessary to enter minutely into the details of this case, which at first appeared to progress as favourably as could be desired. On the fifteenth day after the operation the ligature came away with the dressings ; and on Saturday, the 20th October, exactly four weeks after the vessel had been tied, I find the Hospital Report to be as follows : " Patient's health is now very good ; he is up all day and walks about the grounds ; sleeps well during the night ; has no pain or uneasiness ; the discharge from the wound daily diminishing in quantity, and assuming a more healthy character." But on Monday, the 22nd, matters began to assume a different aspect. He complained of

pain and stiffness in the neck, with headach, furred tongue, and general constitutional derangement. The sac had begun to inflame. On Saturday the 27th, (five weeks after the operation,) “the pain and swelling of the neck had greatly increased; the skin tense and shining, of a deep red colour over the centre of the tumour, more faint towards its border; the apex soft and elastic, with a distinct sense of fluctuation: he described the pain as being most excruciating, and of a hot and throbbing character. He had intense head-ach; foul tongue; bowels obstinately costive; pulse 96, hard and full. He also suffered from constant cough, difficult respiration, and painful deglutition. The sac had suppurated, but as this was an occurrence which had frequently taken place in my experience before, I acknowledge it occasioned me little uneasiness, and I prepared to treat the case in the manner I had treated others with uniform success.

I made a free incision into the tumour with a view to discharge the matter, turn out all the coagula, and then by applying pressure externally, seek the obliteration of the sac. The incision gave exit to a large quantity of pus mixed with *fluid blood*, and I found I had opened into a large cavity which scarcely contained any coagulum at all. I laid down the sides of the wound and endeavoured to apply pressure by means of compresses retained by numerous straps of adhesive plaster. This latter indication, however, could not be accomplished. Direct pressure caused an intolerable sense of suffocation, and the consequence was, that under the moderate degree employed, the sac suppurated freely, and the discharge became profuse. Still I imagined I had no cause for apprehension beyond the wearing and wasting hectic that would probably be established; and was therefore surprised at being called at 3 o'clock on the morning of the 30th, with information that my poor patient was bleeding profusely from the wound I had made in opening the abscess. I hastened to the Hospital, and found him literally bathed in blood; and, notwithstanding the exertions of a most active and intelligent

pupil who was resident there at the time, he must have lost between forty and fifty ounces. The bleeding was kept under by the pressure of this gentleman's hand, but immediately on its being removed, the blood spouted forth with considerable force. On examining the cavity, with the aid of a very imperfect light, I discovered several streams of arterial blood passing in different directions through a broken clot at the bottom of the sac, and my first impression was, that some branches opened into and communicated with the cavity ; and as the man must have died of hæmorrhage, whilst I should be endeavouring to secure these, I determined on trying to place the patient in the same condition as if the sac had never been opened, and trusting to the pressure of the coagulated blood for the suppression of the hæmorrhage and the obliteration of the vessels. I therefore passed four needles through the lips of the wound, and applied the twisted suture, which held them firmly together, and effectually stopped the bleeding, and I left my patient safe for this time, but pale, weak, sunken, and evidently unable to bear the loss of more blood.

Friday, the 2nd October. The blood burst out again, welling up profusely from the bottom of the wound, but coming without impetus. As I had now the advantage of the assistance of my colleague, Mr. M. Collis, I determined to explore the cavity, with the view of securing the bleeding vessel or vessels, and if possible preventing any further loss of blood. I opened the full extent of my incision, and began to clear out the cavity, when the blood burst forth in a stream, equally frightful and uncontrollable, flowing from a rent in the vessel that my finger could not cover. In this predicament not a moment was to be lost. The patient had lost so much blood, that a single minute would probably decide his fate. As for seeking to tie a vessel lying at the bottom of an enormous cavity, and fully at a distance of five inches from the surface, it appeared to be wholly out of the question ; and the vicinity of the pneumo-gastric nerve and the deep jugular vein, rendered a plunge of the

needle or the employment of the actual cautery equally objectionable. I had no resource but to fill this enormous cavity with compresses of sponge, which should be maintained in their places by closing the integuments over them, and this latter could only be effected by means of needles and the twisted suture. Straps of adhesive plaster were totally useless; glue spread on leather was tried, and found equally inefficacious: nothing remained but to stitch the wound in the manner specified, and it certainly had the effect of perfectly restraining the bleeding for the time being. However, I knew that there could be no permanent benefit derived. I knew from the examination I was enabled to make, that the blood proceeded from the original aneurismal rent, that the artery which was not affected by the usual processes of nature during the five weeks that intervened between the operation and the suppuration of the sac, would scarcely become obliterated under the pressure of the sponge, and that the needles must ulcerate the parts and cut their way out long before any permanent benefit could be achieved. Immediately on the wound being perfectly closed, the pulsation, which had disappeared, or been so weak as to be only perceptible to the most delicate touch, returned in the tumour as vividly and as violently as before the operation had been performed at all.

It would be tedious to dwell on the minute reports of this awful and melancholy case; suffice it that the patient still continued to bleed at intervals. According as a needle would separate or a compress be disturbed, blood would burst forth with greater or less violence, and although the hæmorrhage was always promptly restrained, yet, these successive losses, in conjunction with pain, loss of sleep, and extreme anxiety, reduced him so low, that he expired, without a struggle, on the evening of the 12th October—having lived thirteen days from the first appearance of the bleeding. Every effort was made to procure a *post mortem* examination, but in vain, and I have been left to speculation to account for the singular phenomena, and unexpected results that attended this most interesting and important case.

But it is not altogether without a parallel, and after some research I have discovered a case which will throw additional light on these cases of aneurism of the carotid artery.

On the 15th April,* 1831, Mr. Green, in St. Thomas's Hospital, tied the common carotid for the cure of an aneurism situated (as he believed) at the point of its division. The patient was a man advanced in life, being 65 years of age: the tumour had existed from the preceding Christmas, was slow of growth, and had attained only to the size of a walnut; in other respects the symptoms detailed bear a strong resemblance to those observed in the case just related. On the ligature being tightened, a manifest and instantaneous diminution of size took place in the tumour and in the force of pulsation, which was yet distinguishable; but the patient having been carefully removed to bed, in about an hour this had ceased altogether. It is worthy of remark that doubts were entertained by many who carefully observed the case, as to whether the pulsation ever ceased completely.

"On the twenty-fourth day the ligature separated, the noose thereof being perfect and firm, and the dressings having been applied to the wound daily. This has some exuberant granulations, not occupying more than half an inch, which have been touched with *argentum nitratum*. A feeble pulsation has been constant since the 20th April, (the fifth day after the operation,) and we are of opinion that it has latterly been more vigorous; the tumour itself is very materially diminished, but not to the degree that, at this distance of time from the obliteration of the canal of the vessel, we should reasonably expect."

"May 31st. Arterial pulsation has become more distinct in the tumour, but is yet weak. It is supposed, from the situation of the latter, (at the bifurcation of the common carotid,) which is favourable to such, that a communication exists in that part, between the external and internal divisions."

* For the details of this case, see Dr. Johnson's *Medico-Chirurgical Review* for April, 1832, (No. 32,) page 520.

“ June 14th,” (two months after the operation.) “ The aneurismal sac has grown larger within the last fortnight, and the pulsation remains equally, if not more powerful.”

“ July 28th. Patient is suffering from suppuration of the right tonsil ; a diffused swelling has taken place externally upon the neck, a little below the angle of the jaw. The tonsil was opened, and discharged pus freely.”

“ September 20th. The above local symptoms subsided in due course, but the aneurismal swelling retains the same trifling pulsation.”

I have merely extracted from the printed report of the above case, the points in which it bears some resemblance to that of my patient, and which seem to have some reference to the locality of the disease. It was supposed by Mr. Green to exist at the bifurcation of the carotids, a spot equally favourable with the internal itself, for the return of pulsation through the medium of the cerebral circulation. This pulsation did return, and according to the opinion of some, never totally disappeared. About the 28th July, symptoms occurred which might have been occasioned by the suppuration of the sac, and at the end of five months pulsation remained ; the disease had not been cured. The case seems to hold a middle place between that of Elizabeth Rourke, in which the progress of the disease was extremely slow, and the contents of the sac nearly solid, and that of Markey, which had proceeded so very rapidly, and in which there was probably no coagulum at all. It is to be regretted that the appearances of the tumour, in relation to the mouth and pharynx, were not described, and its ultimate result not ascertained.

Thus far, however, I have endeavoured to show that aneurisms of the internal carotid are unfavourably placed for the accomplishment of one part of the cure ; that a ligature on the trunk does not entirely cut off the impulse of the heart from the diseased vessel ; that such impulse may be conveyed through the circulation of the brain in its normal state ; and that the ef-

fect of this impulse must be, to disturb the blood within the sac, and delay, if it does not prevent, its coagulation. I have now to direct my attention to another circumstance of, at least, equal importance. In a limb, when the aneurismal sac becomes distended with blood after operation, the pressure exercised thereby must be directed against the injured vessel ; the structures external to, and surrounding such sac are, many of them, inelastic, all more or less resisting ; they will not permit a growth or extension of the tumour in any direction towards them, and consequently when the sac is filled, and more particularly when its contents are solidified, it must not only press against the ruptured vessel, but compress it to an extent and degree to occasion its obliteration. But with respect to the internal carotid, the case is widely different. However covered with fascia, and muscle, and skin, and other resisting structures externally, it is wholly unprotected in the direction of the pharynx, for, on making a vertical section of the head and neck, and dissecting from within outwards, I find that the internal carotid is very close to the pharynx. In its passage upwards, from the bifurcation to its entrance into the skull, it obliques slightly backwards and inwards, having behind it the sympathetic nerve and first cervical ganglion, where it rests against the spine : external it has the styloid process, the three styloid muscles, the digastric, the mastoideus, and the different layers of fascia ; a little in front it has the stylo-pharyngeus muscle, but internally, or towards the pharynx, it has nothing but the mucous membrane, the constrictor of the pharynx, some very loose cellular tissue, and the twigs of the superior laryngeal nerve ; thus the aneurismal sac has ample room to grow and increase inwardly, and consequently the pressure it is forced to make on the opening in the vessel may be so trifling, as not in any way to lead to its obliteration. This circumstance may explain (although in Mr. Green's case it does not appear that the mouth and pharynx were ever examined) why that case proved to be what Scarpa might term an imperfect cure, and why in the case of Markay, in five weeks after the

vessel was tied, and the direct force of the heart cut off, there should have been no advance whatever in the process by which the artery might be obliterated. It appears curious that the position of this vessel, with respect to the pharynx, should not have attracted more attention in the examination of aneurismal tumours ; for, reasoning from the anatomy of the parts, I should be disposed to believe that the symptoms of pulsation would always be most clearly observed from within. It was so in the present case, and I have since had an opportunity of seeing it under circumstances where it might be still less expected or looked for.

A young girl named E—— M—— was admitted into the Meath Hospital, with a varicose aneurism situated at the angle of the jaw, and extending downwards a short way along the course of the external jugular vein. It became very large and prominent when pressure was made on this vessel below, so as to interrupt the current of blood ; it became then excessively painful, and exhibited the usual thrilling sensations, both to the finger and the ear ; but when looked at from the mouth, a strong and continued pulsation, together with considerable tumefaction, was obvious to every eye. This disease had been produced by a stab of a scissors, inflicted seven years before, and no very decisive treatment was adopted for it in hospital ; in fact, the exact nature of the lesion was not understood, and it was only matter of conjecture, that a communication had been established between the external jugular vein and the internal carotid artery, with the intermediate existence of a varicose aneurismal sac. The case, however, is pathologically interesting, as affording an illustration of the facility with which such tumours may grow and increase internally or towards the mouth.

Whilst on this subject, I may be permitted to notice a circumstance in the case of Markay, in explanation of which, I confess myself unable to form even a remote conjecture. Why did the blood remain in a fluid state within the sac during five weeks ? Before the artery was tied, and whilst a large current

was forced into the sac with the full strength of the heart's action, it is not difficult to conceive that such constant and violent motion might interfere with the process of coagulation ; but that after such an interval, the blood, with the exception of one or two very small coagula, at the bottom, should have been found perfectly fluid, was scarcely to have been expected, and must have had a very unfavourable influence on the success of the operation. Yet such was the case, and it is no easy matter to explain why it should have been so. But the difficulty of solution attendant on this question, leads me to propose another. Is there a pathological difference in the blood of different individuals, giving to that of one a greater or less tendency or disposition to coagulate, than to that of another ? If there is such a difference of condition or constitution, a knowledge of the fact, and more particularly of the causes or circumstances that lead to its production, might prove of incalculable benefit in the management of disease in general, but more particularly those of the circulating system. And that there is such a difference, I am strongly disposed to believe, although being totally unprepared with satisfactory proofs, I dare not offer it even as an hypothesis—but rather as a suggestion that may lead others (as opportunity may offer) to investigate the pathology of the blood as promising to lead to invaluable practical results. That the blood of individuals suffering from different diseases, will exhibit different phenomena in the quickness with which the coagulum is formed, and the degree of firmness and solidity it reaches, no one will be disposed to deny, but the point to which I wish to direct attention is, that the blood of a person apparently in a healthy state, may not coagulate under circumstances wherein that of another individual would almost certainly do so.

Some years ago, a man was admitted into the Meath Hospital, having received a stab of a sharp-pointed shoemaker's knife about an inch below the right sterno-clavicular articulation, by which the internal mammary artery and vein were wounded. These vessels poured out their blood continually,

and in such abundance into the cavity of the pleura, that the lung became dreadfully oppressed, and it was deemed advisable to perform the operation of paracentesis on the fourth or fifth day after the receipt of the injury. The wound made in the operation was large, in the expectation that it might facilitate the escape of any coagulum, but the precaution was found to have been unnecessary, as the blood had remained in a perfectly fluid state, and flowed away with the greatest facility. The quantity of blood thus lost was enormous: it must have amounted to some quarts: and as the wounded vessels still continued to bleed, it is scarcely necessary to add that the patient soon died. On examination after death, a good deal of fluid blood was found in the cavity of the pleura, but not a particle in a state of coagulation. Is it not reasonable to conclude, that in this case there was some peculiar condition of the blood that rendered it incapable of coagulation? and might not this have been one cause of the continued and unceasing hæmorrhage, without the slightest effort on the part of Nature to arrest it?

On the 5th August, a man named James Wilson was admitted into the Meath Hospital, with an enormous aortic aneurism.

Three months previously he had perceived a tumour of the size of a hazel nut above the clavicle, and close to the sternum, which tumour increased in size daily, enlarging from below upwards: on admission it occupied the anterior and right side of the neck, extending as high as the thyroid cartilage, and slightly displacing the larynx. It pulsated violently. From the symptoms, the history of the case, and careful stethoscopic examination, it was decided that it was an aneurism of the aorta, and that palliative measures only should be adopted. He remained in hospital until the 8th September, when the tumour burst, and the patient died in a few seconds, with an awful gush of blood.

Post Mortem Examination.—On looking at the situation occupied by the tumour during life, it appeared to have shrunk and collapsed, and instead of being elevated, there was a very large cavity, in the centre of which was a dark spot marking the place

at which it had given way. The sac was eight inches in its long diameter by five in the transverse at its widest part, and sprung from the aorta at the root of the innominata, passing up behind the clavicle, which was in part carious. Not a particle of lymph, or fibrine, or coagulum of any description existed in the cavity of this immense sac, although upwards of four months had elapsed since the commencement of the disease, and the tumour had not increased with any very extraordinary rapidity.

From these cases, and from other facts and observations, with which it is unnecessary to swell this paper, already pushed to perhaps a tedious length, I have been led to believe that a condition of blood indisposing to coagulation, may exist in individuals otherwise apparently healthy, and exhibiting no symptom indicative of such an abnormal state. If such an opinion be correct, and if (as is generally conceded) the coagulation is an active agent in the suppression of hæmorrhage, and a necessary part of the process in the cure of aneurism, few subjects can be submitted to the attention of the pathologist of greater apparent importance, or more likely to repay the trouble of investigation.

ART. V.—Appendix to DR. GRAVES's Report on the Progress of Cholera.

DR. GRAVES feels great pleasure in communicating to the readers of this Journal, the following official Cholera Returns.

The Tables were constructed by the Commissioners appointed to adopt the measures most likely to arrest the progress of cholera in Great Britain, and whose duty it became to watch and record the progress of the disease. This duty they performed with a diligence and accuracy which stand unrivalled, and their labours have succeeded in furnishing a more valuable and accurate history of the march and effects of cholera, than any other country can boast. By the aid of these Tables, any person can