days of the starvation. Perhaps there is some benefit to be expected from hot poultices or ice bags over the epigastrium. It is much to be regretted that our views of the exact effect and value of these measures should not be more settled and unanimous than they are; some of us prize the hot applications, some use the cold, while others deny the value of either.

In the subacute and chronic cases the daily full flushing of the bowels by alkaline laxative waters is useful, but it is irrational to suppose that gallstones are thus washed away. Nor do olive oil or any other of the pretended expulsive agents have the smallest effect. The thing that happens is probably merely the elimination of effete matter from the body, thus increasing the physiologic resisting power. This process is aided by a restricted diet of the most proper foods and by general good hygiene. It is the custom in Europe to send patients with chronic gall-tract diseases to some of the spas (mostly to Carlsbad) for a course of the waters, to see if an operation can be avoided by the appearances of recovery. These measures are all useful, for the patient gets the benefit of an abstemious regimen, for a time at least, in addition to the waters, and thus is frequently able to improve in a marked manner.

To go to the springs, drink freely of water and live on a low diet, be much out of doors, and rest from work and perhaps from cares is always wholesome for these cases. Such a vacation can be taken once or twice a year with great propriety, but it has little power to postpone the day of a final surgical operation or death, if, on returning home, the patient goes back to his former habits of overfeeding, irritation of stomach, lack of depuration, and generally unhygienic life. This latter is just what a large number of patients are constantly doing. By this course they invite infection from irritating things in their gall tracts and so hasten surgical intervention. This last is no misfortune if the surgery can be done at the right time and under the best conditions; but they usually regard it as a misfortune and think they are fighting it off by their occasional water cures.

The only proper course for such patients, who would be fairly well and avoid surgery, is to live lives of absolute regularity and abstemiousness, rigidly temperate in everything except the use of water, and hygienic and moderate in every day's living. One day of overdoing, overeating or overdrinking, or of severe chilling of the body, may start an infection and precipitate an acute attack that will destroy life unless prevented by timely surgery.

If a chronic gallstone patient who refuses surgery wishes to manage his health in the usual erratic manner and so make an operation or death certain, it is perhaps his privilege as a citizen to do so. But it is only common fairness that he should know the inevitable tendency of his ways, and it is for the physician to show him this and to point out a wiser course, and then leave the responsibility with the patient, where it really belongs.

Operative Treatment of Lung Cavities.—According to J. Berry, in the *Clinical Journal*, June 6, 1906, incision and drainage constitute the only operation on lung cavities that is of practical importance. This is more feasible in cavities at the base than at the apex or deep seated in the lung. It is indicated in abscess where the results are good (mortality 20 per cent.) and in gangrene (mortality 35 per cent.). In bronchiectasis and tuberculosis the results are not encouraging. A patient with abscess due probably to the bursting of a liver abscess into the lung made a good recovery after operation.

NEPHROLITHIASIS.

A. H. CORDIER, M.D.
KANSAS CITY, MO.

In the searchlight of that greatest of all beacons, pathologic and surgical experience, new territory is being invaded and taken possession of by modern surgery, and fewer and less debatable grounds remain along the border line of internal medicine. In no locality is this better settled than in nephrolithiasis. Stones in the pelvis of the kidney or ureter, when known to exist, are looked on as dangerous foreign bodies and dealt with accordingly. Occupation, locality, or age does not, in my opinion, have any direct etiologic bearing on these cases. As a matter of fact, we do find more cases in persons between the ages of 20 and 50 years, but we find more people between these periods. I have found this condition to exist about equally in the two sexes. High living and sedentary habits are said to predispose to the formation of stones in the kidney, but my experience does not sustain this claim. The separation of the solids from the urine in the pelvis of the kidney gives rise to the condition known to the layman as "gravel" and produces only mild symptoms of mechanical irritation of the ureter, bladder and urethra. This condition may be transient, or it may lead to more serious stone formation by these little particles becoming agglutinated, by the presence of an infection, blood, mucus or pus, yet it is possible for a stone to form without the evidence of bacterial invasion or structural changes in the kidney. The nucleus of a stone may be composed of one salt of calcium, or uric acid, while the body of the calculus is made up of another. Carbonate of calcium stones are rare in man, and are deposited in alkaline urine, while the uric acid and oxalate of calcium are found to have their origin in acid urine. The chemistry of these cases, while very valuable and interesting from a scientific standpoint, is overshadowed by the diagnostic and surgical phases, to the busy practitioner and surgeon.

Oxalate of lime calculi are found most frequently, and usually produce the severer types of symptoms and pathology.

The right kidney, like other structures on the right side of the median line in the abdomen, is the one most frequently affected by stone, yet in rare instances both organs have calculi.

VALUE OF EARLY SURGERY.

To appreciate thoroughly the value of early surgery in these cases, one must understand the pathologic changes wrought by time in the ureter and kidney.

It is only a question of time in every case of renal calculi until an infection is developed, and, when once started, a nephritis is the result; if the ureter is occluded, dilatation of the pelvis takes place, and a pressure obliteration of the parenchyma is sure to follow unless quickly relieved. Perinephritic abscess, with extensive burrowing of the escaped fluids, takes place in some cases. The source of the infection may be from the bacteria in the blood, but usually is from an ascending ureteritis. A surprising number of infectious cases have their origin from the *Bacillus coli communis*, probably by direct invasion through an aseptic adhesion of the kidney to the ascending colon.

The number of stones and their sizes present a wide range, from one to dozens, from a mere speck of sand to a stone weighing many ounces.

I have found that the cases of uric acid stones, as a

rule, have only one small, irregular, angular, speculated stone, while the oxalate of lime stones have been large, multiple stones, with roughened surfaces, taking the shape of the kidney pelvis and calices. The phosphatic variety is usually an irregular, friable, semi-mushy stone found in old suppurative cases (Fig. 1). The carbonate of calcium stone, a rare variety, is smooth, multiple, irregular and very hard, producing kidney structural changes only slowly.

The range in weight of stones removed in my own work has been from a few grains to nearly four ounces. The calculus, as a rule, is found in the pelvis of the kidney or in one of the calices, yet it is of no uncommon occurrence to find a good-sized stone imbedded in the parenchyma. These are the cases that so often give rise to postnephritic abscess. These imbedded stones are often overlooked at the time of the operation.

Morris estimates that both kidneys are affected in about 10 per cent. of the cases. In none of my cases have I found both organs with stones in them.

The uric acid and oxalate of calcium stones form more easily than the pasty, semi-solid, phosphatic deposits.

A small stone may pass from the kidney and give rise

Fig. 1.—Nephritic calculi, carbonate of calcium. Weight, 1,540 grains.

to no symptoms, and a very large calculus may be found as a surprise in the pelvis of the kidney at a postmortem. Several subjective symptoms may be absent in the presence of a stone easily palpable externally.

A healthy kidney is sooner or later converted into a diseased one by harboring these foreign bodies. This change usually takes one of the three courses: Interstitial sclerosis, atrophic changes in parenchyma due to calculous hydronephrotic pressure or suppurative pyelonephritis, and rarely carcinoma or sarcoma as the result of chronic mechanical irritation. These pathologic changes have all shades of variation from a mild microscopic change to a complete destruction of the kidney with death of the patient from coma, amyloid changes in other organs, with a hectic accompaniment and death. While a stone remaining in the kidney is of much danger to that organ, the hazard is materially increased if the calculus, in its endeavors to reach the bladder, becomes lodged in the ureter. A stone thus situated may cut its way out of the ureter and lead to urinary infiltration, or completely block the flow of urine from that kidney and destroy the functions of the

organ by pressure of retained urine, or it may be the source of a calculous anuria by interfering with the function of the opposite organ. The persistence of pain is a matter of no small consequence in such a case.

DIAGNOSIS.

In making a diagnosis of stone in the kidney, it might be well for us always to remember that there are a number of other maladies producing symptoms misleading in character, and, again, a stone may exist without producing the typical stone history, or the symptoms may be manifested in a remote locality. We know that many obscure febrile symptoms are produced by a stone in the gall bladder; the same is true of nephrolithiasis. A stone that disturbs the surface epithelium and permits absorption may give rise to a urinary (septic) chill, the source of which may puzzle the diagnostician. I have seen a number of such cases.

If at a nephrolithotomy a large stone is found and there is a short symptomatic history, we all know that the stone was a long time forming; hence must have had a long quiescent period. I am firmly of the opinion that in such a case symptoms were present, but not correctly interpreted. The use of the microscope and the sifting of obscure evidence will often lead us to make a correct diagnosis where others have failed. Lumbago, a pain referred to the glans penis, or clitoris, increased desire to urinate, retraction of testicle, etc., all, as a rule, are sympathetic manifestations. A carefully compiled clinical history extending over a period of years will often clear the way to a correct deduction. An attack of pain in the side, or back, ten or fifteen years ago, may have been an attempt to expel a calculus that has remained quiescent in pelvis or kidney ever since, the patient being in danger all the time of the development of a suppurative process. Tenderness, as a symptom of renal calculi, is often absent, and too much reliance should not be placed on it. Tenderness, as a rule, is found only in those cases in which there exists a marked enlargement from inflammatory involvement of the kidney. Pain is present in the great majority of these cases. Of much diagnostic value is hematuria, as with few exceptions blood can be detected in the urine unless the ureter is completely occluded. In that event the presence of renal colic or pain, with an enlargement of the kidney, would aid in the diagnosis. The referred pains are often present during the passage of the stone along the ureter. The ilioinguinal, iliohypogastric and genitocrural are the nerves most frequently affected.

In renal colic the typical diagnostic symptoms will depend on the size and shape of the stone, as well as on its consistency and surface condition and on the elimination and nature of the attack, which lasts from a few minutes to several days, and disappears as suddenly as it appeared, the stones reaching the bladder or dropping back into the dilated pelvis of the kidney. These attacks may recur at any time, depending on the number of stones or the ability of any stone to block the ureter.

A feature that is noticeable in many cases is the increase of the pain and colicky spells from riding on a rough road. The pain during the passage of the stone is the most intense of any suffering and is usually accompanied by vomiting and profuse diaphoresis. If much blood or pus is present, a chemical test for albumin may be misleading unless the microscope be used also. Hyaline casts are found in a certain percentage of these cases and are evidence of stone irritation to kidney structure.

Rigidity of muscles on the affected side is often present, while in some cases, if the stone is large, it may be palpated. The *x*-ray, while a valuable adjunct, must not be relied on too much. The negative findings especially should be given little weight, as the thickness of the body and the comparative density of the stones presents such a wide range of resistance to the rays that their use may be misleading instead of an assistance. In doubtful cases, however, I would advise the use of the *x*-ray, and if the clinical history were stronger than the radiographic findings I would rely on the former.

Oxalate of calcium stones (Fig. 2) throw a dense shadow. The same may be said of the phosphate, with a nucleus of some one of the other urinary deposits. The uric acid stone presents the least resistance to the rays, as has been demonstrated by the radiograms recently made by myself and Dr. McCandless.

In arriving at a diagnosis of nephrolithiasis, a number of other conditions presenting very similar symptoms must be eliminated. With blood in the urine increased by exercise, pain along the ureter and severe colicky spells and a history of passing one or more stones, a diagnosis of stone is easy.

DIFFERENTIAL DIAGNOSIS.

The various diseases affecting the structures in the vicinity of the kidney may mislead one, but, as a rule,

Fig. 2.—Oxalate of calcium stone. Weight, 1,100 grains.

the symptoms of these affections are such as to point to an easy diagnosis.

Neuralgia and muscular rheumatism, while often found in the vicinity of the kidneys, are usually of a transient character and have symptomatic traits peculiar to themselves, as a bilateral tenderness and a stiffness of muscles, either from pressure or movements of the body. If the diagnosis can not be made otherwise, a course of antirheumatic treatment may assist. Spinal caries and other diseases of the vertebra, in their early history, are confusing when compared to the masked cases of stone in the kidney, but a resort to the microscope and a little time will usually eliminate the simulated diseases.

Neuralgia due to vasomotor disturbances in the kidney has led to a number of mistakes, and many patients have been operated on and no stone found; but, thanks to the splitting of the capsule and relief of the tension, most of these individuals are cured by operation. Patients who pass large quantities of oxalate of calcium crystals form a class of cases under the general head of neurasthenics and present many symptoms of early nephrolithiasis, the so-called "gravel" cases. In these cases it will often be hard to determine the exact extent of the pathology. Displacements of the kidney, presenting a "Deitel's Crises," are usually easily diagnosed.

While the late symptoms of tuberculosis of the kidney

are very typical, the early manifestations are equally misleading. With pus in the urine, frequent micturition, some hematuria and pain in the back, with colicky spells due to the passage of cheesy debris, the history certainly simulates stone.

A close scrutiny of other organs of the body should be made for tubercular foci, as the testicle, bladder, lungs, prostate and seminal vesicles, and tubercle bacilli in the urine should always be looked for. Hectic symptoms may exist in both conditions as a late manifestation. The presence of stones in a tuberculous kidney is not unusual, the phosphatic stones predominating. In most of my cases the right kidney has been the one affected, and the symptoms have been of such a character as to point definitely to the organ diseased, yet in some cases it has been rather perplexing to determine not only which kidney was diseased, but whether or not the other was affected also.

The knowledge of the exact location of the stone prior to operation is of value only in as far as it concerns the position of the incision, yet this is not of much value, as the kidney should be explored in all cases, even though the stone producing the symptoms is located in the ureter, the possible exception to this being in female patients in whom the stone can be removed through the vagina or directly by the bladder route. The cystoscope, while of great aid in diagnosing diseases confined to the bladder, has been found of little value in doubtful cases of stone in the kidney. However, it has its use as an aid to the introduction of the ureteral catheter in investigating some extremely obscure ureteral and kidney symptoms. The collection of urine from one or both kidneys separately is often necessary to determine the source of abnormal constituents found in that fluid.

Kelly and others, by the aid of wax-tipped ureteral catheters, have detected stones in the pelvis of the kidney. This procedure is one that requires much skill and will rarely be resorted to as an aid to diagnosis of stone in the pelvis of the kidney. The condition of the opposite kidney, or even its presence, should be demonstrated in every case in which the question of a nephrectomy is being considered. Palpation of the opposite kidney can often be accomplished, and the study of the character of the urine and the amount and ureteral catheterization will aid in making a diagnosis. The history of one kidney being hydronephrotic, while normal urine is being passed from the other, will point to one healthy kidney.

PROPHYLAXIS AND MEDICAL TREATMENT.

The function of the kidney being essential to life, it behooves the physician to preserve this organ by every possible and safe means. A crippled kidney from stone injury regains its function in a surprisingly short time if the source of the injury is removed, the rapidity of this recuperation depending on the length of time, the size of the stone, its location and the extent of the destruction of the pelvis and parenchyma. A recognition of these truths should admonish us to resort to early surgical interference when once the diagnosis is made.

Nephrolithotomy is considered one of the safest operations in surgery, unless by delays the case has assumed a grave character by the destruction of the urine-manufacturing portions of the organs.

No case should be permitted to go on until abscess in perirenal structures has developed or until urinary fistulae have formed, obliterating all normal structures surrounding the organs. Conditions such as mentioned

above add very materially to the difficulties and dangers of the operation, preclude in many cases the possibilities of saving the kidney, and make the convalescence a very unpleasant and tedious one, marked by suppuration, urinary fistulæ and more surgery.

A healthy kidney, when incised, heals as readily as any tissue in the body.

I have much faith in the prophylactic management of these threatened stone cases, but no reliance should be placed on methods advocated to dissolve and to remove stones when once formed. It is true that Nature in many cases will cause the passage of small stones, but we should not deceive our patients and delude ourselves in the belief that our efforts are of much avail in her behalf.

In a patient known to be suffering from "gravel" (passage of large amounts of oxalate of calcium or other crystals) the regulation of this diet, drinking of much water and the administration of the alkalies, potash, soda and benzoate of lithium, to prevent the deposit of these crystals, will be of much benefit. Piperazin, in from five to twelve grain doses, given three times a day, is highly extolled.

Urotropin (hexamethylenamin) is another remedy much used, as it does not derange the digestive organs and is a good urinary antiseptic.

The unfortunate feature of medication in these cases is the fact that only too often an incorrect diagnosis is made, and the medical treatment is persisted in to the extent that the very dangers we wish to avoid are permitted slowly to take place—that is, extensive injury to the kidney structure and the crippling of its function.

A patient suffering from an attack of renal colic will require large doses of morphin to relieve his pain, but it might be well to recall the fact that pain produces a tolerance of this drug, that the kidneys are one of the main sources of its elimination, and that the pain often ceases suddenly, leaving the patient in a condition of profound stupor from the cumulative action of the repeated doses of the drug at short intervals.

The medical treatment in these cases, of course, is palliative only, but must be resorted to, as in many cases while we are making the patients comfortable the calculus may pass, and with proper after-care another may never develop.

The operative indications in nephrolithiasis are fairly well marked in most cases, yet in some the case may assume a borderline position and require very careful analytic consideration as to its proper management.

OPERATIVE TREATMENT.

Hematuria of long duration or oft repeated may demand operation. Renal colic recurring often and severe would also indicate an operation. The presence of a stone in the pelvis of the kidney, whether producing colic or bloody urine, is a source of danger and should be removed. Chronic invalidism, when due to the presence of a stone, is often cured by the removal of a stone in the kidney that was not producing any local symptoms of a grave character.

When it is possible (and this is usually the case) a nephrotomy should be the operation of choice. The same incision is made for a nephrotomy as for a nephrectomy, that usually preferred being the oblique incision extending from an inch below the border of the last rib in front of the erector spinæ, downward the distance necessary. A few branches of the intercostal and lumbar vessels may require ligating; the nerves of the same name will also be encountered and should be

shoved aside when possible. The costoilic space of the diseased side may be increased by a good-sized sand bag or roll of sheets placed on the opposite side, the patient being practically in the Sims position.

The kidney can usually be delivered into the wound, and an exploration by palpation or otherwise made. In those cases in which the perirenal structures have been inflamed and converted into a vascular dense mass the delivery of the kidney is in most cases impracticable owing to the profuse bleeding when this vascular capsule is torn. The organ can then be explored by needling in all directions, taking care to avoid the larger vessels at their entrance into the hilum. While one may miss a small stone by this process, it will be the exception. If the presence of a stone is strongly suspected and the needle fails to find it, an incision on the convex surface of the kidney should be made, and the pelvis explored carefully with the finger. A stone with only the thickness of the ureter or pelvic structures can usually be detected by palpation, but small stone attached in the parenchymatous crater may be overlooked. If the stone is not found by needling or digital exploration, a metallic ureteral catheter should be gently introduced and the ureter sounded. This failing, I have for a number of years resorted to hydrostatic pressure to test the

Fig. 3.—Phosphatic stones.

patency of the ureter, and in two instances have succeeded in forcing stones into the bladder from the lower portion of the ureter. This procedure I always resort to after clearing the pelvis of stones as a precautionary routine to test the condition of the ureter.

The hemorrhage, on introducing the needle, is often quite profuse, but always stops of its own accord within a short time. The flow of blood from an incision into the kidney is alarming, but is quickly controlled by the pressure of the exploring finger, and if the kidney can be stitched at the completion of the operation the flow of blood can be checked. If a stone is detected its removal should be carried out with the utmost care lest it be crushed and small particles escape into the ureter or remain as a nucleus for the development of more stones.

If small particles have unavoidably been detached, or if there exist many small stones, the assistant should compress the ureter or twist the pedicle. A good gauze pack against the ureter will prevent any small particles getting beyond easy reach of the operator or his irrigating fluid. A gauze pack beneath the delivered kidney will also aid in controlling the bleeding during the

operation. On two or three occasions I have resorted to the use of rubber tubing carefully tightened about the pedicle of the kidney to limit the loss of blood. This will rarely be necessary.

After the stones have been removed, the kidney irrigated and the patency of the ureter established (if the case was not one of severe infection), the kidney may be closed as with any other wound, using chromicised gut for that purpose, taking care that the sutures do not go through the pelvis and thus act as a nucleus for further stone involvement. If it is a suppurating case the pelvis should be packed with gauze and a rubber drain put in. The gauze will aid in controlling the bleeding, especially if good firm pieces of gauze packing are put in around the kidney also. The wound is dressed in the usual way. This gauze should remain in place for several days, as a too early removal will induce a troublesome hemorrhage. Do not become discouraged at the duration of the drainage, as many of these old cases will drain for months, ultimately closing for good.

The removal of the kidney in any case is a matter that should receive very careful consideration. The indications for a nephrectomy are becoming fewer and rarer, as we understand the dangers of such a removal, and the recuperative powers of a crippled kidney if the source of irritation is removed.

When possible, the stone should always be removed through the parenchyma, as an incision into the upper ureter or pelvis will incur more risk in the development of an unhealed urinary fistula.

CONCLUSIONS.

1. Nephrolithiasis is more prevalent than is generally supposed.

2. A stone having formed in the pelvis of the kidney, if too large to pass, will sooner or later produce symptoms demanding its removal.

3. Suppuration in stone cases as a rule is a late process, and should be prevented by early surgical treatment.

4. There exists a special cause for stone development in the right kidney; 80 per cent. of my cases have occurred in the right kidney.

5. With a carefully obtained clinical history, the diagnosis of stone in the kidney is easy.

6. Obscure, persistent pains in the region of the kidney, in a patient who had a renal colic years ago, should lead to an exploration of the organ.

7. A wound in a healthy kidney heals quickly.

8. The operation of nephrolithotomy has a very low mortality.

9. Hemorrhage from a kidney is generally easily controlled.

10. At time of operation, the patency of the ureter should be established.

11. The kidney should not be removed if it has a patent ureter or unless it be practically destroyed by the disease.

12. Postoperative patience and faithful effort on the part of the surgeon, will result in the saving of many organs; otherwise removal will be necessary.

Typhoid Strumitis.—Inflammation of the goiter is described by Baradulin in a man 36 years of age who had had a goiter for 29 years. During the fourth week of typhoid fever an abscess developed in the right half of the goiter, which was opened, and bacteriologic examination revealed typhoid bacilli. —*Ruesky Vrach.* No. 14, 1906.

THE SURGICAL TREATMENT OF GASTRIC AND DUODENAL ULCER AND ITS RESULTS.*

WILLIAM J. MAYO, A.M., M.D.
Surgeon to St. Mary's Hospital.

ROCHESTER, MINN.

The contributions to the literature of ulcer of the stomach and duodenum have become so numerous that it will be impossible in a paper of this character to give credit for the many valuable advances which have been brought out by individual workers. I wish in this connection, however, to express my appreciation of their labors.

For more than twenty years the complications of this malady have received attention. Operations for benign stenosis of the pylorus and some acute secondary manifestations have occasionally been performed, but not until the last five years has there been a surgical invasion into the field of unhealed ulcers with a view of giving relief from pain, under-feeding and chronic disability which so frequently accompany the disease.

Acute ulcer of the stomach and duodenum properly belong to the domain of internal medicine, and surgery has to do only with the complications such as perforation, hemorrhage and obstruction.

The surgical treatment of acute perforations of the stomach and duodenum is now on a sound footing, the results depending on speedy diagnosis and prompt operative relief. Patients operated on within the first five hours usually recover; after ten hours the majority die. We have had 7 gastric with 2 deaths and 9 duodenal with 3 deaths.

Suture of the perforation and pelvic drainage with or without irrigation, the patient being subsequently maintained in the semi-sitting posture (exaggerated Fowler's) for several days, gives the best results.

One mooted point concerns whether or not a gastrojejunostomy shall be done at the same time, as a considerable percentage of patients following perforation will develop stricture, adhesion, or other deformity in the healing process. If the perforation lies in the duodenum or close to the pylorus it should be sutured transversely to avoid interference with the lumen of the part involved. Generally speaking, if it seems probable that stricture or other secondary condition will result and the patient is in good condition, gastrojejunostomy should be performed providing it can be done without spreading the already present infection. If in doubt, a living patient is better than a completed operation at a greatly enhanced risk.

Hemorrhage from the stomach occurs in two forms, acute and chronic. Recurring acute hemorrhages are best treated by opening up the stomach or duodenum, locating the bleeding point and suturing the part firmly with catgut on a curved needle from the inner (mucous) side. The outer surface is then exposed and protected over this area by a few mattress musculo-peritoneal sutures of linen (modified Andrews method). We have not found gastrojejunostomy for acute hemorrhage a reliable procedure. Out of 6 cases we have had 1 patient bleed to death within two weeks after gastrojejunostomy and a second saved by opening up and securing the bleeding vessel, while 5 who have been treated by primary operation on the bleeding point with or without excision of the ulcer; all recovered.

* Read in the joint session of the Sections on Practice of Medicine and Surgery and Anatomy of the American Medical Association, at the Fifty-seventh Annual Session, June, 1906.