the City of Westminster, St. Marylebone, and Stoke Newington; and the highest rates in Islington, Finsbury, Poplar, Bermondsey, Deptford, and Woolwich. The prevalence of scarlet fever showed a marked decline from that recorded in the previous month; among the several metropolitan boroughs this disease was proportionally most prevalent in St. Pancras, Poplar, Bermondsey, Battersea, Deptford, Lewisham, and Woolwich. The number of scarlet fever patients under treatment in the Metropolitan Asylums Hospitals at the end of the month was 1727, against 1439, 1630, and 1825 at the end of the three preceding months; the weekly admissions averaged 183, against 160, 207, and 229 in the three preceding months. Diphtheria was rather less prevalent than it had been in October; the greatest proportional prevalence of this disease was recorded in Hammersmith, Fulham, Chelsea, Islington, Finsbury, the City of London, and Bermondsey. The Metropolitan Asylums Hospitals contained 866 diphtheria patients at the end of last month, against 580, 746, and 852 at the end of the three preceding months; the weekly admissions averaged 106, against 76, 108, and 120 in the three preceding months. The prevalence of enteric fever showed a considerable decrease from the somewhat excessive figures for the two preceding months; among the several metropolitan boroughs this disease was proportionally most prevalent in Hackney, Finsbury, Bethnal Green, Stepney, Southwark, Bermondsey, and Lambeth. There were 134 enteric fever patients under treatment in the Metropolitan Asylums Hospitals at the end of November, against 58, 141, and 169 at the end of the three preceding months; the weekly admissions averaged 16, against 32 and 25 in the two preceding months. Erysipelas was proportionally most prevalent last month in Hackney, Finsbury, Bethnal Green, Stepney, Poplar, and Deptford. The 29 cases of puerperal fever notified during the month included 3 in Hackney, 3 in Stepney, 3 in Lambeth, and 2 each in Kensington, Poplar, Battersea, and Wandsworth. The 4 cases of cerebro-spinal meningitis notified during the month belonged respectively to Paddington, St. Marylebone, Shoreditch, and Wandsworth.

The mortality statistics in the table relate to the deaths of persons actually belonging to the several boroughs, the deaths occurring in institutions having been distributed among the boroughs in which the deceased persons had previously resided; the death-rates are further corrected for variations in the sex and age constitution of the populations. During the five weeks ending Dec. 3rd the deaths of 7742 London residents were registered, equal to a corrected annual death-rate of 17.4 per 1000, the rates in the three preceding months having been 10.3, 11.9, and 12.6 per 1000 respectively. The death-rates last month ranged from 11.3 in Hampstead, 12.1 in Lewisham, 12.8 in Wandsworth, 13.1 in Greenwich, 13.4 in the City of London, 13.6 in Chelsea, and 13.7 in Woolwich, to 20.5 in Poplar, 21.2 in Shoreditch, 22.1 in Bethnal Green, 23.3 in Southwark, 23.7 in Finsbury, and 24.9 in Bermondsey. The 7742 deaths from all causes included 672 which were referred to the principal infectious diseases; of these, 409 resulted from measles, 21 from scarlet fever, 56 from diphtheria, 45 from whooping-cough, 23 from enteric fever, and 118 from diarrhoea. No death from any of these infectious diseases was recorded last month in the City of London; among the metropolitan boroughs they caused the lowest rates in Chelsea, the City of Westminster, Hampstead, Wandsworth, and Lewisham; and the highest rates in Hammersmith, Islington, Finsbury, Shoreditch, Southwark, Bermondsey and Camberwell. The 409 deaths from measles were more than three times the average number for the corresponding period of the five preceding years; this disease was proportionally most fatal in Hammersmith, St. Pancras, Islington, Finsbury, Shoreditch, Southwark, Bermondsey, and Camberwell. The 21 fatal cases of scarlet fever showed a decline of 36 from the corrected average number, and included 3 in St. Pancras, 3 in Poplar, 2 in Stepney, 2 in Battersea, and 2 in Lewisham. The 56 deaths from diphtheria were 25 below the corrected average number; the greatest proportional mortality from this disease was recorded in Hammersmith, Finsbury, Bethnal Green, Southwark, and Bermondsey. The 45 fatal cases of whooping-cough showed a decrease of 29 from the corrected average number; the highest death-rates from this disease last month were recorded in Hammersmith, Fulham, St. Pancras, Finsbury, Shoreditch, Bethnal

Green, Bermondsey, Battersea, and Woolwich. The 23 deaths from enteric fever were 8 below the corrected average number; of these 23 deaths, 4 belonged to Lambeth and 2 each to the City of Westminster, Islington, Hackney, Bethnal Green, Battersea, and Wandsworth. The 118 deaths from diarrhoea were 33 fewer than the corrected average number; this disease was proportionally most fatal in Paddington, Stoke Newington, Finsbury, Bethnal Green, and Bermondsey. In conclusion, it may be stated that the aggregate mortality in London last month from these principal infectious diseases was 27.4 per cent. above the average.

Correspondence.

"Audi alteram partem."

LAVATORY ACCOMMODATION FOR WOMEN.

To the Editor of THE LANCET.

SIR,—I desire to bring to your notice the accompanying resolutions which were passed at a conference of representatives of the following bodies on Nov. 21st:—The Royal Sanitary Institute, the National Health Society, and the Women's Imperial Health Association. Mrs. Oxley Grabham, who has conducted an extensive investigation into the sanitary accommodation provided by railway companies for women, was also present.

With regard to resolution No. 1 it may be said that although the system now so largely adopted by the railway companies of providing lavatory carriages on long-distance trains, has proved an enormous benefit to both men and women travellers, still there is considerable evidence that the accommodation for women at railway stations all over the country is often insufficient and insanitary, whilst the necessity for payment, which is still universal for women at railway stations, operates harshly on the travelling public of the more necessitous classes.

Appropriate communications have been addressed on the subject of the aforesaid resolutions to the Local Government Board, the Board of Trade, the London County Council, and the principal railway companies, and the conference desires to enlist the powerful aid of THE LANCET in making the resolutions known to its readers, or in any other way helping a reform which is so intimately associated with the health and comfort of the female members of the community.

I am, Sir, yours faithfully,

LOUIS C. PARKES,

Chairman of Council, the Royal Sanitary Institute.
London, S.W., Dec. 16th, 1910.

[RESOLUTIONS.]

1. That in the opinion of this meeting sufficient sanitary accommodation for women's use should be provided free of charge by sanitary authorities, railway companies, and in places of public entertainment and resort.
2. That in the opinion of this meeting it is essential in the interests of public health and decency that there should be an attendant in charge of the sanitary convenience provided for women's use in public places.
3. That in the opinion of this meeting all sanitary conveniences provided for the use of women should be subject to periodical inspection by women sanitary inspectors appointed by the local sanitary authorities.
4. That this meeting desires to call the attention of the railway companies, and managers of theatres and other places of public resort, to the unsatisfactory character of much of the sanitary accommodation provided for women, and to the necessity for having only sanitary conveniences of good modern type and construction.

THE POSITION OF HOMŒOPATHY.

To the Editor of THE LANCET.

SIR,—The appreciative and sympathetic references recently made by the press to the death of my old and valued friend, Dr. Dyce Brown, whose ability, amiability, and keen artistic sense had won for him the affection and respect of a wide circle, must have made others besides myself think whether it is not time to reconsider our attitude towards homœopathy. The objections to that method of treatment, so far as I understand them, are based on the belief that its professors are practising what they do not believe, and are therefore guilty of a grave breach of the moral law, in consequence of which they are scorned

and rejected of men, boycotted in medical circles, excluded from hospital appointments, and practically from social intercourse with their brethren. It is not likely that respectable people honestly trying to do their duty according to their lights should feel anything but sore when they are thus branded with the mark of Cain, and a bitter feeling has thus grown up between the two equally qualified sections of medical and surgical science, which, I venture to think, is little creditable to ourselves. For what great crime, after all, have the homœopathists committed? The absurd doctrines of Hahnemann have long since been abandoned, and all that they now do is to treat disease on a much more scientific principle than ours, by often-repeated doses of strong tinctures instead of the nauseating tablespoonfuls and wineglassfuls which too often nauseated the patient and retarded his chances of recovery. I have no doubt that Sir Richard Quain lost the Presidency of the Royal College of Physicians of London, which he had fairly earned by earnest and successful work, because he met Dr. Kidd in consultation over Lord Beaconsfield's case, and I well remember the storm in a tea-cup which ruffled the calm waters of Pall Mall, when an attempt was made to "run in," or rather "run out," Dr. Murrell and Professor Ringer for an alleged therapeutic heresy, an attempt which properly and ignominiously failed. I suppose my professional brethren are too much beaten down by the authority of the Royal Colleges to air their views.

I am, Sir, yours faithfully,

Finzean, Dec. 19th, 1910.

R. FARQUHARSON.

THE MOTOR FUNCTIONS OF THE STOMACH.

To the Editor of THE LANCET.

SIR,—In his letter of Dec. 17th Mr. Gray excuses himself for ascribing Bayliss and Starling's discovery of secretin to Wertheimer, who had nothing to do with it, by saying that in the same paragraph "an extensive quotation, duly ascribed to Starling, appears." However gratifying this must have been to Professor Starling, a quotation from a discussion by Starling on some observations made by Edkins (who is not mentioned by Mr. Gray) hardly compensates for giving the credit of Bayliss and Starling's epoch-making discovery of secretin to "Wertheimer and others."

With regard to Mr. Gray's "recent radiographers and others," who "assert that the pylorus during gastric digestion descends naturally below and to the left of the umbilicus," I think that Mr. Gray must have misunderstood Dr. Barclay's references to the work of Pfahler and Holzknacht instead of referring to their original papers. Dr. Barclay will doubtless reply to Mr. Gray himself, so that I shall content myself with quoting from Pfahler and Holzknacht. According to Pfahler,¹ the pylorus is "usually about an inch above the umbilicus," and "the pyloric portion extends from one to two inches beyond the median line to the right," and Holzknacht² states that the pylorus is above the umbilicus and "lies in front of or to the right of the spinal column." The figures given in Pfahler and Holzknacht's papers as types of what they regard as normal stomachs bear out this description. The disagreement between various radiographers as to what is the "normal" stomach is simply a matter of terminology. Barclay, Rieder, Holzknacht, and Pfahler all agree as to the shape of the stomach most commonly seen in healthy people, but while Barclay and Rieder think that this shape should be regarded as the normal, Holzknacht and Pfahler think that the "ideal" form, in which the greater curvature does not reach below the umbilicus and the pylorus is on a level with the lowest point of the organ, should be regarded as normal, although they admit that it is comparatively rarely seen.

Our disagreement as to the occurrence of peristalsis in the "cardiac end" of the stomach is, I expect, simply due to a misunderstanding as to nomenclature. Peristaltic waves generally begin near the junction between the upper and middle thirds of the stomach; this point lies in what Cunningham, following His, called the "body" of the stomach, and corresponds probably to what Mr. Gray calls "the distal part of the cardiac portion," which seems to me

to be a less convenient term. Mr. Gray apparently believes that these waves at first pass no farther than his "third sphincter," whilst physiologists and radiographers agree that they pass without interruption to the pylorus.

Why does Mr. Gray think that food may pass directly into the duodenum in infants, but only after a considerable delay in adults? I can only imagine that it is because it is easier to observe the process through the small, and therefore more transparent, bodies of infants than through the more opaque bodies of adults. I have watched food (and by food I mean *food*, and not water or albumin water) enter the duodenum immediately after a meal has been taken in a very large number of cases, and any afternoon that Mr. Gray calls at Guy's Hospital I am sure that Dr. A. C. Jordan, our medical radiographer, will be pleased to demonstrate this to him in as many cases as he may desire.

If Mr. Gray thinks that the pylorus requires "at least a little free hydrochloric acid before its sphincter action is unloosened," how does he account for the fact that, according to all writers on this subject, the stomach empties itself with undue rapidity in primary achylia gastrica? In this condition, as well as in almost all cases of carcinoma of the stomach, in nearly all cases of pernicious anæmia, and in many of simple gastritis, free hydrochloric acid is present at no period during digestion, so that the pylorus should never relax and its contents never leave the stomach, if Mr. Gray were correct.

Even if Dr. Barclay disagrees with me as to the position of the pylorus, which I doubt, by what process of reasoning does Mr. Gray conclude from this that Barclay's observations on duodenal ulcer, which are in complete agreement with mine, should not be accepted as confirmation of mine? Our observations prove that the late onset of pain is not due to food being held back by the pylorus or by Mr. Gray's "third sphincter," as we both have found that the food passes into the duodenum with unusual rapidity in cases of duodenal ulcer. The true cause of hunger pain hardly comes within the scope of a discussion on the "motor functions of the stomach."

I am, Sir, yours faithfully,

Weymouth-street, W., Dec. 17th, 1910.

ARTHUR F. HERTZ.

ORGANIC PREPARATIONS OF ARSENIC.

To the Editor of THE LANCET.

SIR,—It would be a striking confirmation of the truth of Goethe's admonition, "Warum in die Ferne schweifen, sieh das Gute liegt so nah," were it now to be shown that what is practically our oldest organic preparation of arsenic, introduced into therapeutics 46 years ago by a now forgotten German physician, exhibits the same action on spirochætae as the recently evolved complicated compound of another compatriot.

Cadet in 1760 obtained a fluid by distilling a mixture of arsenic and potassium acetate; this was known under the designation of Cadet's fluid, alkarsin, or fluid pyrophore. Bunsen, in 1837, first analysed this fluid, and found that it consisted principally of cacodylic acid and cacodylic oxide, which could be converted into cacodylic acid by oxidation. The first pharmacological experiments with this preparation were undertaken by Bunsen himself on frogs, and by Kürschner on rabbits. The latter found that a dose of 0.42 gramme could be given intravenously without harm; indeed, one animal bore this dose following the administration per os of a dose of 0.36 gramme. In 1843 it was therefore already established that cacodylic acid was non-toxic. In 1860 Schmidt and Chomse confirmed the above statement regarding the non-poisonous character of cacodylic acid, but they found that under certain circumstances it was reduced in the stomach. Lebahn, in 1868, stated that it possessed the action of arsenic, but that it was considerably less toxic than the latter or its salts. Schulz (1879), Marshall and Green (1886) were also able to confirm the above statements; in other words, as far back as 1886 no less than eight authors had published the results of their physiological and pharmacological investigations of cacodylic acid. Jochheim was the first to try the use of cacodylic acid in his practice in 1862, and he published at the time a report of the excellent results he had obtained from its use in the treatment of chronic skin diseases and pulmonary tuberculosis. I recently came across a now forgotten publication of his which bears the date 1865 and is in the possession of the Grand Ducal Court Library here. In this work he says that in his opinion

¹ Journal of the American Medical Association, xlix., 2070, 1907.

² Mitteil. aus den Labor. für radiol. Diagn. u. Ther., Vienna, 1906.