a post-mortem, but the progress of the paraplegia will be well worth your careful study. I have repeatedly witnessed this form of paraplegia in cases of recurrent cancer after excision of the breast, and in two or three cases other bones besides the vertebræ were affected. In two ladies some years ago, on whom I had operated within a few weeks of each other for cancer of the breast, almost precisely the same series of events followed. Both were partially paraplegic, when in each spontaneous fracture of the femur from cancer of the bone occurred. In some cases the disease of the spine stops short of paraplegia, severe pain in the back being the only symptom. In these probably the cord escapes any implication in inflammation. Modern observers are much inclined to believe that myelitis is almost a necessary link in the chain of which paraplegia is the final result, and that pressure alone rarely produces that symptom. Since I spoke to you last week on this case I have received from Dr. James Hutchinson, the distinguished physician to the Pennsylvania Hospital, a copy of a valuable paper on this subject. He has recorded two cases with post-mortems, and in each myelitis was found. He insists on the fact that pain usually precedes symptoms of paralysis, and that the advent of the latter is usually definitely due to myelitis. His concluding sentence is:—"Symptoms belonging properly to myelitis are, as I have attempted to show, paraplegia, impairment or loss of the different varieties of sensation (anæsthesia and analgesia), loss of power over the bladder and sphincter, and permanent contraction of the limbs and bedsores."¹

XII. Recovery from Paraplegia possibly of Syphilitic origin.

We sometimes, in going over the past life-histories of our patients, come upon interesting fragments of evidence, always, however, to be taken with a certain amount of hesitation on account of the sources of fallacy. A gentleman who consulted me a few days ago for a syphilitic gumma in his tongue told me that he had formerly suffered from paralysis, and had wholly recovered. The facts were these. Fifteen years ago he had syphilis, with eruption, &c., rather severely, and took mercury. Almost as soon as his treatment was over, he began rather suddenly to lose the use of his legs. The weakness began at his heels, and crept upwards, and his lower extremities wholly failed. He was on the couch for three or four months. His arms were weak, but not wholly paralysed. After four or five months' treatment he wholly recovered, and has never since had any reminder of it. He was married at the time, and married a second time some years afterwards. He has now enjoyed ten years of good health, and has during that time had nothing of a specific character.

FURTHER REMARKS ON THE CONTAGIOUSNESS OF YELLOW FEVER.

By ROBERT LAWSON,
INSPECTOR-GENERAL OF HOSPITALS.

(Concluded from p. 589.)

THE important points in the foregoing details are—first, that extensive and repeated importations of malignant yellow fever were not followed by an extension of the disease to any of the susceptible individuals who were exposed to the emanations from the sick; secondly, that at the end of October yellow fever appeared among the patients in hospital who had not been in communication with those labouring under yellow fever, apparently from exposure to the emanations from the marshy ground in its vicinity, and the hospital sergeant and orderlies of the 33rd, who had all come into contact with the imported cases for months previously, without being affected, were then attacked, and six of them died; thirdly, that, after excluding the night air, the 47th had no further attack in the hospital, and only a single case in the middle of December, which came from barracks; fourthly, the immunity of the men on guard at the hospital,

who were outside the usual range of the emanations carried by the wind from the swampy ground. These facts leave no alternative but to conclude that the yellow fever among the residents in the hospital, in this instance, was due to emanations proceeding from the marshy spot mentioned. Nor is this the only occasion yellow fever has been observed to spring up here under similar circumstances. In 1839, after heavy rains alternating with hot sun, which is described to have rendered the low ground close to the hospital "a complete swamp," the first cases of yellow fever in the garrison were the hospital sergeant and one of the orderlies of the 52nd, who were attacked here on the 29th of October. The disease then appeared among the soldiers' families of that corps, occupying huts near the Brick barracks, rather less than half a mile away. It then affected the officers, and finally the men. Altogether there were 93 cases, and 27 deaths, in the regiment.¹ So long ago as 1816 the liability of persons in the military hospital to fever was remarked. During a severe epidemic that year 11 of the servants here contracted fever, while of those at what was designated the naval hospital, in a more elevated and dry position, nearly half a mile to windward, there were only 2 attacked (both of whom had frequently to visit the former), though a large proportion of the fever cases among the troops, and all the sick of the navy, comprehending many of the worst cases of yellow fever, were treated in it. Even at the military hospital, where the fever cases were placed on the upper floor, the orderlies who attended them, and who invariably slept on the same floor, were less frequently attacked than those employed otherwise, who had not this advantage of elevation during the night to place them above the denser portion of the emanations from the ground.²

The following is another well-marked example of the harmlessness of yellow fever cases of the most virulent description when imported into the same locality when no local cause was in operation. H.M.S. *Dauntless* arrived at Barbadoes on Nov. 16th, 1852, from St. Thomas, having had 41 cases of fever and 11 deaths during the passage. The sick were sent to the military hospital the following day. Part of the crew were landed and encamped, but as it was deemed impossible to land the whole of them at one time, those retained on board continued to be exposed to the causes of the disease in the ship herself, and new cases were in consequence of almost daily occurrence. Between Nov. 17th and Jan. 5th, 1853, 22 officers and 135 men were admitted to the military hospital, and 13 of the former and 52 of the latter died. Mr. Denny, surgeon of the 34th Regiment, states: "Although 61 men of the *Dauntless* have been treated in the wards of the 34th hospital, indiscriminately intermingled with the soldiers of that corps affected with various complaints, in no instance has any individual been attacked under such circumstances, nor has any hospital attendant suffered. I have now (Jan. 5th) 18 cases of fever from the *Dauntless* in hospital, and 37 men of the 34th Regiment, but not one affected with any febrile disorder."³

Proceeding now to Jamaica, we have Port Royal, the naval station, with a dockyard and naval hospital, besides a small garrison consisting usually of a company of European artillery and a detachment of black troops. The inhabitants of the town had suffered a good deal from intermittent and remittent fever during the summer of 1857, but the local medical practitioner, as well as the military and naval surgeons, had not met anything like yellow fever among them, and the only case which had occurred at this station was in a white artilleryman who had been at Spanish Town on the 24th July, and on his return was confined in the cells from the 26th to the 31st. That afternoon he was discharged, and after drinking freely at the canteen, went to hospital in the evening with fever, which proved to be yellow fever, and he died on the 2nd August with black vomit. Having been at Spanish Town so recently, where he had been drinking and exposing himself, it is impossible to say whether this man's attack was attributable to that locality or to Port Royal.

There was no other case here until October, when a very sharp outbreak commenced, which was at first confined to a line of three buildings 225 feet in length, parallel to the line wall, and distant from it some fifty feet. The wall is not above ten or twelve feet high, and does not prevent the free access of the sea breeze during the day. The space between

¹ See Transactions of College of Physicians of Philadelphia, Third Series, vol. iii. The paper is accompanied by a bibliography of the subject.

² Quarterly Report of Medical Officer in charge, to Dec. 31st, 1839.

³ Bancroft's Sequel to an Essay on Yellow Fever, p. 448-9.

⁴ Milroy in THE LANCET, April 13th, 1861, p. 358.

these houses and the wall is covered with a bed of loose open gravel, and a foot beneath the surface of this the soil was moist. The houses were of two floors; the upper floor of the end ones was occupied as officers' quarters, and of the centre one as mess-room, and the ground floors were employed as offices, library, stores, &c. In the southernmost of these houses an officer's wife, who was nursing an infant about two months old, was attacked with fever on October 16th; on the 17th the mess waiter, who slept on the ground-floor of the mess-room, was attacked; on the 19th the husband of the first case, and the same day a child of a soldier who acted as librarian, and who passed much of its time with him in the library on the lower floor of the southernmost house. This man and his family resided in another house in the same line with, but 400 feet distant from, the library, with seven other families, all of whom remained healthy, though the case last mentioned and the following one was treated there. On the 23rd another child of this man's was attacked, who had also been much with him in the library, and on the 24th the man himself. On the 26th a second mess waiter was seized who had slept in the same place as the previous one. On the 31st an officer's wife, who lived at a distance, but had attended to the first case very assiduously, contracted fever. On Nov. 7th, a young officer living in the northernmost house was seized, and on the 10th another in the same house. All these were well-marked cases of yellow fever, and five of them died with black vomit. The white artillery, with a very few exceptions, were removed to the mountain station of Newcastle on Oct. 28th. Of these three were attacked at Newcastle, and one died; one of these had been much in the library at Port Royal, but not near any fever case, the other two had been in attendance on fever cases. Two white artillerymen who remained at Port Royal were seized on Nov. 15th and Dec. 3rd; both died yellow and with black vomit. A black soldier was attacked on Dec. 17th; he had yellow conjunctivæ, hæmorrhage from gums, and albumen and tube-casts in urine, but recovered. This man occupied a bed close to a window on the ground floor of the men's barrack, which is 400 ft. to the west of the southernmost of the three houses mentioned above; the ground outside the window was removed to some depth, to improve the surface drainage, just before his attack. Two European artillerymen, who were sent from Port Royal to the Kingston Penitentiary on Oct. 23rd, had fever there on Oct. 28th and Nov. 11th respectively; the former died, the latter recovered. The medical officer who had attended most of the above cases at Port Royal embarked for England on Dec. 12th, was attacked the same day, and died at St. Thomas on the 16th, yellow and with black vomit. No case of yellow fever had been brought to Port Royal shortly before the outbreak commenced, nor had any of the subjects of the first eight cases, with the exception perhaps of the third, been away from it for some time before; neither had the librarian and his children, or the mess waiter, any communication with the first case, or with each other, after the fever commenced in them. We must conclude, then, that these persons contracted the fever from exposure to emanations originating in the locality, as they have been shown to do above, both at Sierra Leone and Barbadoes. What occurred at this particular spot may of course have taken place at others in the neighbourhood also, or elsewhere, as, for instance, in the penitentiary at Kingston; and until this possibility of the disease originating from such local causes in any given instance be excluded satisfactorily, we can never be justified in concluding that it arose from communication with imported cases.

The following striking case of importation took place a few months after the outbreak just detailed. The U.S. steamship *Susquehanna* was lying at Greytown, in the middle of March, 1858, when the first case of fever appeared in her. On April 5th she arrived at Port Royal, with her crew suffering from yellow fever; and within the next three days eighty-five of these were landed at the naval hospital, where twenty-three died with the characteristic features of the disease. Some of the medical attendants and visitors had never had yellow fever; and, though no seclusion was practised, not a single case occurred, either within the hospital or in its neighbourhood. One man of the *Imaum* receiving-ship, who attended the funeral of some of the *Susquehanna's* people, but in a separate boat, became affected with yellow fever within a day or two; the disease, however, ceased with him, and, as sporadic cases had occurred in Kingston both before the arrival and after the departure of the *Susquehanna*, nothing can be made of this

single case. After leaving Jamaica the *Susquehanna* proceeded to New York. During the passage there were several fresh attacks and some deaths; and when in quarantine at New York all the crew who were sent on shore continued in good health, while of the small number left on board six sickened and two died. At last everyone was removed from the ship, and she remained unoccupied for nearly three months, at the termination of which period her stores were discharged, when several persons employed in landing them were attacked with yellow fever. White troops had returned to Port Royal in March, 1858, and remained quite healthy, though the officers occupied some of the quarters which had proved so sickly in the previous October; and the town itself was healthy. The cause of the disease was evidently in the *Susquehanna* herself; and, having arrived at Port Royal when there was no cause of fever active there, her people, like those of the *Hecla*, *Dauntless*, and others at Barbadoes, could not communicate the disease, though it was of a virulent form. Had her arrival taken place shortly before the outbreak among the troops six months previously, many would have overlooked all the circumstances detailed above, and have concluded that she had imported the disease and communicated it to them.

A CASE OF BULLET-WOUND OF THE ANTERIOR LOBES OF THE BRAIN.

WITH REMARKS.

BY THOMAS SMITH,

SURGEON TO ST. BARTHOLOMEW'S HOSPITAL.

THE following case should, I think, be put on record, as the patient made a good recovery, without loss of mental or physical power, after shooting himself clean through the head from side to side, the bullet passing in at one temple and out at the other.

A person of education and some acquirements, aged thirty-five, was brought into St. Bartholomew's Hospital under my care within half an hour of having made an attempt on his own life with a revolver.

There was a wound in the fore part of the right temple, from which blood was flowing freely; blood was also running from the nostrils, and there was great and rapidly-increasing swelling beneath the scalp, which extended all over the frontal and orbital regions to the opposite side of the head, where there was a large effusion of blood beneath the scalp; and here, in the left temple, a hard body was to be felt, which was taken to be the bullet; the soft parts about the fore part of the head were emphysematous. The cellular tissue of the eyelids, orbits, and brows was distended with blood, the right eyeball being protruded and the ocular conjunctiva being pushed forward. The bleeding from the wound, which was very free, was at first controlled by pressure on the temporal artery, but after a while blood began to flow from a deeper source. The pupils were natural, pulse feeble, respiration natural.

The patient was quite conscious, and answered questions correctly concerning his name, age, address &c. of his own accord. He spoke in German, but when addressed in either French or English he would reply in the corresponding language. He showed no sign of mental incapacity, nor was there any loss of motor power. Being blinded by the swelling of his eyelids, and annoyed by the quantity of blood about his face, he searched methodically for his handkerchief in various pockets, and having found it, used it in the ordinary way. Whilst lying on the stretcher in the surgery he vomited several times, the vomit consisting of blood mixed with food, the straining in each attack of sickness forcing blood and cerebral substance from the wound in the right temple. A probe could be passed its whole length into the wound and across the head without meeting with the slightest resistance. On being asked if he felt it, the patient replied, "in the ear." A piece of lint soaked in carbolised oil, with a pad and a light bandage, being applied, the patient was removed to bed, when it was found that he had passed both fæces and urine under him. The quantity of