ture, the vaccination is merely the *occasion*, and not the cause. Erysipelas may be prevalent as an epidemic, or some other infectious disease, at the time of vaccination; or there may be an unfavorable state of health in the person, or a predisposition to scrofulous complaints which have always lain dormant in the system, awaiting any exciting cause for development. Several witnesses refer to such a state of things. Dr. Cabot refers to two cases vaccinated with the same matter—one his own father, who nearly lost his life, and upon the other it had no effect. The fault, in the first case, arose from constitutional peculiarities. Dr. Cabot relates several cases in which erysipelas followed vaccination, but he could easily account for it.

Dr. Homans mentioned two cases of erysipelas following vaccination, which he could not well explain unless there was some predisposition to it. He says that "vaccination may *educe*, but not *create* erysipelas: also develop other constitutional diseases." Dr. Bigelow says, "I have known three cases in which erysipelas occurred in a vaccination patient, beginning about the eighth or ninth day: two of these proved fatal." But he does not attempt any explanation of them.

Dr. John Ware states that he never had a case of erysipelas following vaccination, but thinks it possible, though of very rare occurrence. Several of the medical witnesses pronounce the cases at Westford as *phlegmonous* erysipelas, and two others describe them the same as *dissection* wounds. The inquiry here suggested is, does not an *animal* poison produce a *specific* inflammation—having a run, and affecting the system, unlike any kind of erysipelas? Would not a different cause, entirely unlike any other, produce a distinct and specific disease?

In the seven worst cases at Westford, there was a remarkable resemblance as to all the leading symptoms and effects, both local and general.

There are two inferences that naturally follow the perusal of this report. *First*, The matter of vaccination should assume far more importance and attention on the part of the medical profession than it now receives; and, *secondly*, the public should be enlightened and set right in reference to the subject of disease being communicated by vaccination.

SURGICAL CASES.

BY A. F. SAWYER, M.D.

[Communicated for the Boston Medical and Surgical Journal.]

CASE I.—*Wound of the Profunda Femoris near the Linea Aspera; Ligature of both Extremities.*

A young Spaniard, aged about 23 years, in a quarrel with one of his countrymen, received a severe knife-wound on the outside of

the left thigh, extending from a little below the trochanter, downward, a distance of about seven inches.

I was called suddenly to see the patient, about a fortnight after the receipt of the wound, on account of a severe secondary hæmorrhage which had supervened. I found him nearly pulseless and speechless, with a large amount of blood oozing and dripping from the dressings of the thigh. Firm compression was made at once over the femoral artery, where it escapes from the pelvis, which immediately checked the bleeding.

On inquiry, I learned that the *primary* hæmorrhage had been of a most dangerous character—the patient bleeding to syncope. Still no efforts had been made to secure the divided vessels by ligature; the wound being stuffed with Monsel's salt, the edges approximated and retained by sutures, then adhesive strap, and a simple roller applied to the limb. Recurring hæmorrhages took place at intervals of three or four days, the patient each time bleeding nearly to syncope, which afforded a temporary check to the drain of blood, to be renewed again when reaction supervened, and the force of the circulation became once more established. At each hæmorrhage the wound was re-opened, then stuffed with Monsel's salt in its pure state, and the compression renewed—this being applied only locally to the wound.

On removal of the dressings, the edges of the wound were found widely gaping and sloughy, enclosing also a large amount of fresh and half decomposed coagula, which were carefully washed out. The deepest part of the wound was at the lower third of the thigh, having its direction behind the femur. The *linea aspera* was denuded for about two inches, and just to the inside of the *linea aspera* the pulsations of the femoral could be distinctly felt, where this artery is seeking its posterior position with reference to the thigh, before it becomes the popliteal artery. We now had but little doubt that the femoral had been pricked, or partially divided, by the point of the knife which had made the wound. As the hæmorrhage appeared to be equally controlled by pressure on the superficial femoral, at the inside of the sartorius muscle, it was concluded that none of the larger branches of the profunda femoris were divided, as at first suspected.

On account of the sloughy condition of the wound, it was deemed preferable to tie the superficial femoral at the usual place of election, rather than to search for the bleeding orifice in the wound itself.

This operation was scarcely concluded, when, to our surprise, the hæmorrhage set in as profusely as before the ligature of the artery, proving conclusively our error, and that the hæmorrhage really came from some one of the larger branches of the profunda femoris.

The ligature which had been placed on the femoral artery was at once removed, and, without loss of time, we proceeded to make

search for the true source of the hæmorrhage, by dilating the original wound, resolving, if we failed, which seemed quite probable, that we would, as a last resort, tie the external iliac.

Fortunately, however, we were able to find the bleeding vessel, resting upon the linea aspera, and which was evidently the terminal branch of the profunda femoris. Both extremities of the vessel required ligatures.

From this period the case progressed favorably. The circulation of the femoral was at once re-established after the removal of the ligature, and the wound healed by first intention. The edges of the original wound were kept gently supported by adhesive plaster, over which emollient applications were applied for a day or two, until a healthy granulating surface was indicated. Firm dressings were then continued, with adhesive plaster and roller, until the wound was completely cicatrized, which occurred in about five weeks after cessation of the hæmorrhage.

CASE II.—*Intra-Capsular Fracture of the Os Femoris in a young Man.*

Mr. F., a painter, aged 24 years, six feet two inches in height, and muscular in proportion to his size, while engaged in work, accidentally fell from a stage which was suspended by ropes, about twenty-five feet from the ground, and struck upon the edge of the curb-stone of the side walk—the force of the fall, and the weight of the body, being received directly upon the left trochanter.

On examination, a short time after the accident, there was found considerable swelling, with discoloration and tenderness about the upper portion of the thigh. The patient had no power of voluntary motion over the limb, and any forcible movement, as flexion, extension, &c., or pushing the head of the femur against the acetabulum, was attended with acute pain, which was referred to the hip-joint. Crepitus was perfectly distinct, especially on outward rotation of the thigh; still there was no inversion or eversion of the foot, no shortening of the limb, and the patient suffered little or no pain when the limb was allowed to rest quietly. These latter indications seemed rather to point to a fracture of the pelvis, involving the acetabulum, than to any fracture of the head or neck of the femur.

We found it necessary to etherize the patient, to enable us to make a more detailed survey of his injuries. We then got the same crepitation as before the patient was insensible, though more strongly marked on outward rotation of the femur. Crepitus was most perceptible when passive motion was given to the limb, by grasping the trochanter firmly with one hand, while the fingers of the opposite hand were pressed deeply towards the joint on the inner side of the thigh. We were unable to detect any grating or unnatural mobility about the pelvic bones, after a very careful examination both externally and by the rectum. The crest of the

ilium, the pubes, pubic arches and ischiatic tuberosities were firm, and in situ, and manipulation was unattended with any pain.

It is difficult to conceive of a fracture of the acetabulum, without fractures involving also the surrounding bone, sufficiently extensive to yield motion and crepitation. There have been rare instances where the acetabulum has been driven inward by the head of the femur, thus producing a shortening of the corresponding limb; but there was no shortening in this case. Here, if there had been any fracture of the acetabulum, the unusual crepitation that existed must have shown the fracture to have been a very extensive one, in which event it is obvious that grating could have been detected by the finger introduced within the rectum, which was not the case.

There was no pain in urinating, nor appearance of blood in the urine; in short, the pelvic viscera had escaped all injury.

We were thus forced to the conclusion that the head or the neck of the os femoris had sustained a fracture—especially when it is borne in mind that the nature of the fall was technically such as to produce this rare injury, the shock being immediately applied to the great trochanter, and transmitted directly in the axis of the cervix femoris. The absence of shortening, or of any distortion in the appearance of the limb, or of any tendency to eversion or inversion of the foot, were peculiar features in the case. The trochanter also appeared to occupy its natural place.

Many explanations have been attempted, by Syme and others, to account for the marked crepitation on rotation, an important diagnostic sign in this class of injuries. For crepitus *from rotation* is much more distinct than can be obtained from any other form of passive motion that can be given to the extremity. However ingenious these explanations may appear, we think, on the whole, they are to be rejected as unsatisfactory.

Was there any likelihood, in this instance, of the head of the bone being involved in the fracture? The absence of shortening of the limb and of other anatomical lesions, characteristic of fracture of the neck of the femur, whether extra- or intra-capsular, makes it probable that there was an oblique fracture of the head of the bone, all displacement being prevented by the integrity of the capsular ligament, and by the acetabulum acting as a closely-adapted splint to confine the fragments in close apposition with each other. We can hardly expect a fracture of the cervix, without a more or less complete laceration of the capsular ligament; and it is in part the laceration of this ligament which permits these displacements to take place, so important in the diagnosis of intra-capsular fractures of the cervix. We know also that fractures of the head of the bone may occur; and we have at the present time a specimen of this fracture in our possession. The injury was produced by a fall from a horse, when the subject of it was about fourteen years of age. The father informed us that the

nature of the injury was never distinctly made out by the surgeons in attendance, which is not to be wondered at, since the accident occurred at a time when the existence of intra-capsular fractures in young subjects was denied.

Treatment.—The patient was placed upon a firm, hard mattress, with his shoulders slightly raised, and both knees flexed and supported upon a pair of pillows—one placed upon the other, thus giving him the advantages without the discomforts of the ordinary mechanical appliances of the double-inclined plane. At the end of two months the joint was sufficiently firm to enable him to bear his weight on the limb without complaint of pain; and there was no pain or embarrassment in the articulation on motion. The fracture was considered to be sufficiently consolidated to allow him to move freely about on crutches.

San Francisco, Cal., April 4th, 1860.

Bibliographical Notices.

Seventeenth Report to the Legislature of Massachusetts relating to the Registry and Return of Births, Marriages and Deaths in the Commonwealth, for the year ending Dec. 31st, 1858.

THE State Registration Report for the year 1858, comes to us with full evidence of the care and labor exercised in putting the facts together and drawing inferences from them, but also with the customary and well-founded regret that these facts are so incomplete.

During the year there were registered, in a population estimated at something less than one and a quarter million for the middle of the year, 34,491 live births, 10,527 marriages, and 20,776 deaths; being a decrease of 829 births, of 1212 marriages, and 504 deaths, as compared with those registered in 1857. Assuming that all the births, marriages and deaths, which actually occurred, were registered, this would give a rate of one birth to 35 persons living, one marriage to 116 persons living, and one death to 59 persons living, and shows an average, for each day in the year, of 94 births, 29 marriages, and 57 deaths. These numbers are from the outline which precedes the main report, and are followed by some just observations from Dr. Curtis on the importance of statistical inquiry, as applied to this class of facts, and upon the injury which is done by the dissemination of erroneous results, drawn from incomplete data.

From the article on population we learn that the rate of increase, from the excess of births over deaths, was 1.13 per cent.; and that the whole rate of increase, from this and other causes, was 3.02 per cent. On page 9 is an interesting table, showing the distribution of the population by ages, in which Massachusetts is compared with England and France, with two of the New England States, and with two of the Southern. It appears that France has in every 100 persons of the population 61 over 20 years, Massachusetts has 58, England has 55, and Kentucky 44. The number over 50 years, is 19 in the 100 in France, 13 in Massachusetts, 14 in England, and 8 in Kentucky. Between the ages of 20 and 50, France has 42 in the 100, Massachusetts