STEREOCHEMISTRY.

DURING the past year but few modifications have been suggested in the current views regarding the arrangement of atoms in space; the main activity in this direction has been in connection with the configuration of quinquivalent nitrogen compounds, a subject in which it can scarcely be claimed that any general agreement of opinion has been reached.

E. Erlenmeyer, jun., and A. Arnold have expressed the view¹ that in addition to *cis*- and *trans*-isomerism of the well-recognised fumaroid and maleinoid types a third kind of stereoisomerism is sometimes to be observed; they derive this type theoretically by supposing one of the groups CR'R'' in the ethylene derivative, CR'R'' : CR'R'', to be rotated at 90° to the other (I or II) so as to give rise to the two enantiomorphously related configurations, III and IV. They consider

that the plane configurations I and II represent respectively *iso*-cinnamic and cinnamic acids, whilst *allocinnamic* acid, which forms hemihedral crystals, is the externally compensated mixture having the solid configurations III and IV. P. Pfeiffer has also discussed the apparent discrepancies between the ordinary views as to the configuration of carbon compounds and certain facts relating to the changes of maleinoid and fumaroid compounds.² He suggests that the configuration of saturated ethane derivatives is one in which the two ethane carbon atoms occupy the centre of an octahedron, at the six apices of which are situated the six groups attached to the pair of carbon atoms; he thus discards the idea that valency acts in definite directions, and also gives up the ordinary conception of double and triple bonds. The configurations at which Pfeiffer arrives by starting from the above assumption are not very dissimilar from those which Barlow has derived on purely crystallographic grounds, and those which Werner has suggested for explaining the isomerism of many inorganic compounds.

¹ *Annalen*, 1904, **337**, 329.

² *Zeit. physikal. Chem.*, 1904, **48**, 40.

The question of the primeval origin of optically active substances has been reopened in a new way by A. Byk,¹ who recalls an old experiment of Jamin's, which indicates that terrestrial magnetism causes the partially plane polarised light reflected from such surfaces as that of the ocean to become partially circularly polarised. In the circularly polarised light thus produced, one component—say, *d*-light—predominates on the earth's surface. Byk then confirms the observation of Cotton² that *d* circularly polarised light is differently absorbed by copper ammonium *d*- and *l*-tartrates. The preponderance of *d*-light on the earth's surface must therefore be more favourable to the persistence of one optically active component of racemic acid than to the other.

A result of great interest has been obtained by W. Marckwald³ in connection with the problem of "asymmetric synthesis." On heating the acid brucine salt of the potentially inactive methylethylmalonic acid, $\text{CMeEt}(\text{CO}_2\text{H})_2$, until the evolution of carbon dioxide is complete, and liberating the valeric acid thus produced, the latter was found to contain a considerable excess of the *l*-isomeride, $\text{CHMeEt}\cdot\text{CO}_2\text{H}$. This is claimed as being the first recorded case of a true "asymmetric synthesis"; to this view, however, Cohen and Patterson have objected that the brucine may separate from the original solution in methylethylmalonic acid with one optically active ion, $\text{CMeEt}(\text{CO}_2\text{H})\cdot\text{CO}_2^-$, of the latter, and that Marckwald's synthesis should then be regarded merely as an application of one of the three Pasteur methods of resolution. To this objection, Marckwald replies, and apparently effectively, that the whole of the methylethylmalonic acid used was converted into the brucine salt.

In connection with the question of asymmetric synthesis, A. McKenzie shows⁴ that on treating menthyl benzoylformate, $\text{PhCO}\cdot\text{CO}\cdot\text{O}\cdot\text{C}_{10}\text{H}_{19}$, successively with magnesium methyl iodide, water, and acid, a mixture of externally compensated and *l*-phenylmethylglycollic acids, $\text{CMePh}(\text{OH})\cdot\text{CO}_2\text{H}$, is obtained; the production of an asymmetric carbon atom in the acidic part of the menthyl benzoylformate molecule is so influenced by the presence of the optically active menthyl group that the acidic group becomes optically active. A similar result is obtained on treating menthyl benzoylformate with magnesium ethyl bromide. Further, on reducing *l*-menthyl benzoylformate, slightly more *l*-menthyl *l*-mandelate than *d*-mandelate is produced.

M. A. Ragusin has made the important observation that the distillates from both Russian and American petroleum are lævorotatory;

¹ *Ber.*, 1904, **37**, 4696; *Zeit. physikal. Chem.*, 1904, **49**, 641.

² *Ann. Chim. Phys.*, 1904, [vii], **8**, 373.

³ *Ber.*, 1904, **37**, 349.

⁴ *Trans.*, 1904, **85**, 1249.

the rotatory powers recorded are small, and the highest values are obtained with the less volatile fractions. These results recall a remark made by P. Walden,¹ some years ago to the effect that if optically active petroleum could be found the geological dispute as to the origin of petroleum would be ended; the optical activity seems to indicate that petroleum is necessarily of animal or vegetable origin.

E. Jungfleisch has recorded² the results of a number of experiments from which he concludes that *l*-lactic acid undergoes hydrolysis more readily than its enantiomorphously related isomeride. The contradiction of theory thus introduced is perhaps counterbalanced by the observations of A. Piutti³ that the recorded statement that the crystalline *d*- and *l*-β-asparagines have different densities is erroneous.

A series of determinations of the speed of inversion of cane-sugar by *d*- and *l*-camphorsulphonic acids has been made by R. J. Caldwell⁴ with the object of ascertaining whether these enantiomorphously related acids effect the hydrolysis of the optically active sugar at the same or at different speeds. Neither in the case of cane sugar nor of milk sugar could any difference in the speed of hydrolysis be detected.

An interesting discussion as to the optical functions of the asymmetric carbon atoms in the closely related alkaloids *l*-ecgonine, *d*-ψ-ecgonine, and anhydroecgonine has been contributed by J. Gadamer and T. Amenomiya;⁵ they conclude that the three alkaloids have the following configurations:

This result follows mainly because (1) the same anhydroecgonine is produced from both *l*-ecgonine and *d*-ψ-ecgonine, and (2) on subjecting anhydroecgonine to treatment with caustic potash, which converts *l*-ecgonine into *d*-ψ-ecgonine, its optical activity remains unchanged.

During the past year, no methods of a novel character have been devised for resolving externally compensated substances into their enantiomorphously related components. Those variations of the second Pasteur method of resolution which have been introduced during

¹ *Naturwissenschaftliches Rundschau*, 1900, **15**, 15.

² *Compt. rend.*, 1904, **139**, 203.

³ *Gazzetta*, 1904, **34**, [ii], 36.

⁴ *Proc. Roy. Soc.*, 1904, **74**, 184.

⁵ *Arch. Pharm.*, 1904, **242**, 1.

recent years have, however, been somewhat extended. Thus, McKenzie, continuing his former work with Marckwald¹ on the esterification method of resolving externally compensated acids, shows that when *dl*-mandelic acid is heated with *l*-borneol the residue of unesterified acid contains an excess of the *l*-acid;² the mixture of esters also yields a preponderance of *l*-mandelic acid. This result is interpreted as meaning that *d*-mandelic acid is esterified more rapidly than the *l*-isomeride by *l*-borneol, since the *l*-acid remains unesterified; the fact that the mixture of esters yields an excess of *l*-mandelic acid on hydrolysis is attributed to the acid, in *l*-bornyl *d*-mandelate, undergoing optical inversion during the formation of the ester more rapidly than is the case with *l*-bornyl *l*-mandelate. On fractionally hydrolysing *l*-bornyl *dl*-mandelate with caustic potash, the first fraction hydrolysed contains an excess of *d*- over *l*-mandelic acid, whilst the last fraction which undergoes hydrolysis gives *dl*- or inactive mandelic acid; *l*-bornyl *d*-mandelate thus appears to be more readily hydrolysed than *l*-bornyl *l*-mandelate, although the latter is the more easily formed.

On hydrolysing *l*-bornyl *dl*-mandelate with more than one equivalent of alcoholic potash, inactive acid is produced; by treating the ester with much less than one-half an equivalent of potash, *d*-mandelic acid is formed, whilst on using more than one equivalent of caustic potash a salt of *l*-mandelic acid is produced. *l*-Mandelic acid undergoes complete optical inversion when heated with 13 per cent. caustic potash during several hours on the water-bath; this fact is interesting in connection with the observation of Kipping³ that *d*-mandelic acid sometimes yields externally compensated phenylchloroacetic chloride with phosphorus pentachloride, and of Easterfield⁴ that *l*-mandelic acid gives the same externally compensated acid with fuming hydrochloric acid at 100°. *l*-Menthyl *dl*-mandelate cannot be resolved into its component esters by crystallisation from light petroleum, and is therefore described as partially racemic.

The resolution of externally compensated primary bases by combination with an optically active aldehyde has been further studied by E. Erlenmeyer, jun., and A. Arnold.⁵ On condensing *dl*-isodiphenylhydroxyethylamine with *l*-helicin, two substances of the constitution

are obtained; these have $[\alpha]_D - 6.43^\circ$ and -43.60° , and on hydrolysis yield *d*- and *l*-isodiphenylhydroxyethylamines having $[\alpha]_D + 109.72^\circ$ and -108.33° respectively in alcoholic solution.

Neville and Pickard⁶ have prepared *l*-menthylcarbimide and Forster

¹ *Ber.*, 1899, **32**, 2130.

² *Ibid.*, 1904, **37**, 378.

³ *Trans.*, 1903, **83**, 1005.

⁴ *Ibid.*, 1891, **59**, 72.

⁵ *Annalen*, 1904, **337**, 307.

⁶ *Trans.*, 1904, **85**, 685.

and Atwell's *d*-bornylcarbimide; they suggest the application of these two substances to the resolution of externally compensated hydroxy-compounds and bases. Forster and Atwell, however, indicate¹ that bornylcarbimide reacts too sluggishly with hydroxy- and amino-compounds to render it useful in the suggested connection. The writer may perhaps be allowed to point out that *d*-butylcarbimide, which forms the main constituent of oil of spoonwort, could certainly be substituted with success for the bornyl- and menthyl-carbimides mentioned above.

A number of externally compensated acids have been resolved into their optically active components during the past year. A new method for resolving fermentation lactic acid has been devised by E. Jungfleisch² depending on the crystallisation of the quinine salt under special conditions; although the carrying out of the new method suggests several points of theoretical interest, it does not appear to offer any advantages over the method devised by Purdie and Walker.³

Morrell and Bellars⁴ have resolved β -methylglyceric acid,

by crystallisation with quinidine; the salt *L*-B *L*-A is the least soluble in water, and the crystalline *L*-acid separated therefrom gives $[\alpha]_D - 13.51^\circ$ in aqueous solution. The salt *L*-B *d*-A could not be obtained in a pure state, but by applying the fact that the barium salt of the inactive acid is insoluble in alcohol, whilst that of the active acid is somewhat soluble, the pure barium salt of the *d*-acid was isolated.

Externally compensated *p*-methoxymandelic acid can be readily resolved into its *d*- and *L*-components ($[\alpha]_D$, in water, $+146.1^\circ$ and -146.2°) by crystallisation with cinchonine;⁵ the specific rotatory powers are but slightly less than that of mandelic acid ($[\alpha]_D +$ or -156.4°).

H. B. Hill and F. W. Russe⁶ have also resolved β -dihydrofurfurane- $\alpha\alpha'$ -dicarboxylic acid by the crystallisation of its cinchonine salts; in one-half per cent. aqueous solutions, the acids have the specific rotatory powers of $+$ or -500° . Owing to the possibility of this resolution, the authors are able to prove that the β -dihydro-acid is externally compensated, and that the so-called α -acid is the potentially inactive isomeride.

Phenylisoparaconic acid has been resolved by crystallisation with strychnine;⁷ the optical antipodes give $[\alpha]_D + 14.72^\circ$ and -14.51° in

¹ *Trans.*, 1904, **85**, 1188.

² *Compt. rend.*, 1904, **139**, 56.

³ *Trans.*, 1895, **67**, 616.

⁴ *Ibid.*, 1904, **85**, 197.

⁵ E. Knorr, *Ber.*, 1904, **37**, 3172.

⁶ *Ber.*, 1904, **37**, 2538.

⁷ R. Fittig and P. Jehl, *Annalen*, 1904, **330**, 292.

alcoholic solution, whilst in acetic acid solutions the signs of the rotatory powers are reversed. *Penicillium glaucum* destroys the *l*-acid more rapidly than its isomeride, so that the latter can be readily isolated by aid of the mould. Phenylparaconic acid,

has also been resolved by crystallisation with strychnine; the component acids give $[\alpha]_D + 64.33^\circ$ and $- 65.33^\circ$ in alcoholic solution.

C. Neuberg and M. Silbermann¹ have set at rest the doubts which prevailed as to the optical constants of the *d*- and *l*-glyceric acids $\text{OH} \cdot \text{CH}_2 \cdot \text{CH}(\text{OH}) \cdot \text{CO}_2\text{H}$, by resolving the externally compensated acid by crystallisation with brucine; brucine *l*-glycerate separates as the least soluble salt, and from it barium *l*-glycerate of $[\alpha]_D - 17.34^\circ$ was obtained. The barium *d*-glycerate from *d*-glucuronic acid gave $[\alpha]_D + 17.1^\circ$.

E. E. Blaise and L. Marcilly² state that $\alpha\alpha$ -methylethylhydracrylic acid, $\text{OH} \cdot \text{CH}_2 \cdot \text{CMeEt} \cdot \text{CO}_2\text{H}$, cannot be resolved into its optically active components by crystallisation with quinine or cinchonine.

S. Condelli³ finds that the optimum temperature for the decomposition of racemic acid by *Aspergillus niger* is 35° ; he shows that *d*-tartaric acid is most readily destroyed by the organism at low temperatures, whilst the *l*-isomeride is the more readily attacked at higher temperatures.

A. Pictet and A. Rotschy,⁴ having prepared synthetically tetrahydronicotyrine and shown it to be identical in constitution with nicotine of natural origin, resolved it into its optically active components by crystallisation with *d*-tartaric acid. The salt *l*-B *d*-A is the least soluble, and from it *l*-nicotine ($[\alpha]_D - 160.93^\circ$), identical with the naturally occurring alkaloid ($[\alpha]_D - 166.39^\circ$), is separated. The hitherto unknown enantiomorphously related isomeride, *d*-nicotine ($[\alpha]_D + 163.17^\circ$), was also isolated. Physiological experiments made with the two isomerides lead to the interesting result that *l*-nicotine is about twice as poisonous as *d*-nicotine; the injection of the *l*-base is extremely painful to the animal, whilst that of the *d*-isomeride seems to cause no pain.

Optically active homologues of benzene have been prepared by A. Klages and R. Sautter.⁵ Magnesium *d*-amyl iodide acts on benzaldehyde yielding *d*-hexenylbenzene, $\text{CHPh} \cdot \text{CH} \cdot \text{CHMeEt}$ ($[\alpha]_D + 50.3^\circ$), which, on reduction, gives *d*-hexylbenzene, $\text{CH}_2\text{Ph} \cdot \text{CH}_2 \cdot \text{CHMeEt}$

¹ *Ber.*, 1904, **37**, 339.

² *Bull. Soc. chim.*, 1904, [iii], **31**, 317.

³ *Gazzetta*, 1904, **34**, 86.

⁴ *Ber.*, 1904, **37**, 1225.

⁵ *Ibid.*, 649.

($[\alpha]_D + 17.2^\circ$); the latter substance could not be caused to undergo optical inversion. The authors found difficulty in preparing *d*-amyl alcohol by the method of Marckwald and McKenzie, and the former has therefore¹ given further directions for applying the method. He also remarks that the difficulty of preparing pure *d*-amyl alcohol is partly due to the interesting fact that all the known crystalline derivatives of *d*-amyl and *iso*-amyl alcohols form solid solutions; thinking that this tendency towards the formation of solid solutions might be sufficiently diminished if large substituting groups were introduced into the molecules of the mixture of alcohols, he has prepared the urethanes. *iso*-Amylurethane and *d*-amylurethane, however, still form solid solutions, although the tendency to do so is less than with the previously investigated solid *iso*- and *d*-amyl compounds. Thus the *d*-urethane is insoluble in the *iso*-compound, but the *isourethane* is fairly soluble in the *d*-isomeride; the two isomerides therefore cannot be separated by fractional crystallisation.

Marckwald finds that the solubility curve of mixtures of the barium salts of *d*-amyl- and *iso*-amyl-sulphuric acids is quite continuous, and has found an analytical expression representing the curve. He also gives a good method of preparing *d*-valeric acid ($[\alpha]_D + 8.75^\circ$) by the oxidation of *d*-amyl alcohol with chromic acid, and has prepared pure *d*-amyl iodide ($[\alpha]_D + 5.64^\circ$) and bromide ($[\alpha]_D + 3.68^\circ$). *d*-Amylamine ($[\alpha]_D - 5.86^\circ$), prepared from *d*-amylphthalimide, is remarkable in that its hydrochloride is optically inactive to sodium light. Marckwald has also prepared the simplest possible optically active paraffin, namely, *d*-methylethylpropylmethane, CHMeEtPr, in a pure state ($[\alpha]_D + 9.5^\circ$) by the interaction of *d*-amyl iodide, ethyl iodide, and sodium.

W. Meyerhoffer discusses² the equilibrium conditions attending the separation of the components of an externally compensated acid by Pasteur's method of crystallisation with an optically active base. Since, in such a crystallisation, three salts are capable of formation, namely, the salts, *d*-B *d*-A, *d*-B *l*-A, and the partially racemic salt, 2*d*-B *dl*-A, there should be three ranges of temperature distinguishable, corresponding with the separation from the solution, side by side, of the two solid salts, (a) *d*-B *d*-A, and *d*-B *l*-A, (b) of the less soluble of the two latter and the partially racemic salt, and (c) of the partially racemic salt alone.

Power and Tutin³ have separated a *l*-quercitol,

having $[\alpha]_D - 73.9^\circ$, from the leaves of *Gymnema sylvestre*. Eight optically active isomerides of this constitution should exist, and the

¹ *Ber.*, 1904, **37**, 1038.

² *Ibid.*, 2604.

³ *Trans.*, 1904, **85**, 624.

one now described is not enantiomorphously related to the *d*-quercitol already known, because this has $[\alpha]_D +24.16^\circ$. The two sugars fucose and rhodose have been shown to be enantiomorphously related by E. Votoček.¹

A good deal of work has been done on the actual determination of rotatory powers and on the deductions to be drawn therefrom. A. Rosenheim and H. Aron find that the rotatory power of *d*-tartaric acid is quadrupled by addition of a caustic potash solution of stannic chloride, and attribute this to the formation of a complex acid ion containing tin and the tartaric acid radicle; the stannitartrates thus formed were isolated by Henderson, Orr, and Whitehead.² H. Grossmann and H. Pötter have examined³ the effect of change of temperature on the rotatory powers of ammonium molybdanylmalate, $\text{MoO}_2(\text{C}_4\text{H}_5\text{O}_5 \cdot \text{NH}_4)_2$, in aqueous solution. In 3—12 per cent. solutions of malic acid to which the corresponding amount of ammonium molybdate has been added, the specific rotatory power attains a maximum value at 35° ; above and below that temperature, the rotatory power diminishes. The same kind of change of rotatory power until a maximum value is reached during continuous rise of temperature has been previously observed to occur by P. F. Frankland and F. M. Wharton in the case of diethyl dibenzoyltartrate.⁴

Patterson has determined⁵ the rotatory powers of methyl, ethyl, and *n*-propyl *d*-tartrates at temperatures from 10 — 100° , and finds that the comparison of the data at the same temperatures indicates but little relationship. By extrapolation, however, it is shown that the esters should be optically inactive for the D-line at 0° , -34° , and -60° respectively, and it is suggested that these are "corresponding temperatures" for the optical properties of the substances. Generally, also, the temperatures T° , $T^\circ - 34^\circ$, and $T^\circ - 60^\circ$, are corresponding temperatures for the three esters, T° being the temperature on the absolute scale. This view is supported by the result that the ratios of the molecular rotatory powers of the esters are nearly constant at corresponding temperatures. At corresponding temperatures, the ethyl *d*-tartrate gives $[\text{M}]_D$ about 2.1 times that of the methyl ester, and the $[\text{M}]_D$ of *n*-propyl tartrate is 1.41 times that of the ethyl salt.

The same worker has also⁶ contributed a mass of valuable experimental data including values of the rotatory powers of sodium tartrate, potassium tartrate, potassium methyl tartrate, potassium ethyl tartrate, potassium *n*-propyl tartrate, and of methyl, ethyl, and *n*-propyl tartrates in aqueous solutions at series of temperatures between 10° and 100° , and values of the molecular solution volumes

¹ *Ber.*, 1904, **37**, 2859.

² *Trans.*, 1899, **75**, 555.

³ *Ber.*, 1904, **37**, 84.

⁴ *Trans.*, 1896, **69**, 1583.

⁵ *Ibid.*, 1904, **85**, 765.

⁶ *Ibid.*, 1116 and 1153.

of methyl, ethyl, *n*-propyl, potassium methyl, potassium ethyl, and potassium *n*-propyl tartrates. A distinct connection is shown to exist between the gram-molecular contractions on solution and the change in molecular rotatory power set up by such solution. The rotatory powers of the alkyl tartrates are increased by solution in water, and a rapid diminution occurs on heating the dilute solution; in the solvent-free state, the rotatory powers of the esters increase as the temperature rises.

P. F. Frankland and J. Harger have studied¹ the rotatory powers of the methyl and ethyl esters of the di-*o*-, -*m*-, and -*p*-nitrobenzoyl-tartaric acids at temperatures between 15° and 180°. The introduction of nitro-groups into the dibenzoyltartrates in para- or meta-positions greatly increases the molecular rotatory power; the introduction of nitro-groups into the ortho-position diminishes the molecular rotatory power above 130°, but as the temperature falls the rotatory power increases so rapidly that ethyl *o*-nitrobenzoyl-tartrate has a higher rotatory power at 15° than has its para-isomeride.

The rotatory powers and dispersions have been determined by E. Wassmer and P. A. Guye² of a large number of acidic derivatives of ethyl malate. In this series, the specific rotatory power diminishes continuously whilst passing from ethyl propionylmalate to ethyl decoylmalate, but the molecular rotatory power attains a maximum in ethyl butyrylmalate. A number of alkyl lactates were also examined, and in the latter series the specific rotatory power attains a maximum value in octyl lactate.

As a supplement to the work of Briggs and Cohen³ on the *l*-menthyl esters of the six dichlorobenzoic acids, J. B. Cohen and H. S. Raper have examined the *l*-menthyl esters of the ten chlorobromobenzoic acids. The values of $[M]_D$ for the corresponding chlorobromo- and the dichloro-esters are of the same order of magnitude. The molecular rotatory power of *l*-menthyl benzoate ($[M]_D - 236.3^\circ$) decreases most when the substituting halogen atoms are ortho- to the carboxyl group, and decreases least when the halogen atoms are meta- to the carboxyl. The rotatory powers of the *l*-menthyl *o*-, *m*-, and *p*-iodobenzoates ($[M]_D - 246^\circ$, -234.5° , and -241.2° respectively) are not in the same order as those of the corresponding chloro- and bromo-benzoates given by Cohen and Briggs (*loc. cit.*) and by Tschugaeff.⁴

W. Urban,⁵ has prepared and determined the rotatory powers of a long series of alkylated *d*-butylthiocarbamides and *d*-butylcarbamides by the action of primary and secondary amines on *d*-butylthiocarb-

¹ *Trans.*, 1904, **85**, 1571.

² *J. Chim. phys.*, 1903, **1**, 257.

³ *Trans.*, 1903, **83**, 1213.

⁴ *Abstr.*, 1903, **84**, ii, 2.

⁵ *Arch. Pharm.*, 1904, **242**, 51.

imide, $\text{CS}\cdot\text{N}\cdot\text{CHMeEt}$. The substances examined were of the types $\text{NHR}\cdot\text{CS}\cdot\text{NH}\cdot\text{CHMeEt}$, $\text{NR}_2\cdot\text{CS}\cdot\text{NH}\cdot\text{CHMeEt}$, and

and the corresponding types in which sulphur is replaced by oxygen. In spite of the large changes which are caused throughout the series in the mass of the group attached to the butyl radicle, the molecular rotatory powers do not differ greatly; large variations in the rotatory power attend the change of solvent from alcohol to chloroform.

J. W. Walker found¹ that the complete hydrolysis of amygdalinic acid yields externally compensated mandelic acid; H. D. Dakin now shows² that, on fractional hydrolysis with baryta, *d*-mandelic acid is first liberated, whilst *l*-mandelic acid preponderates in the hydrolysis product of the final fraction. Amygdalin is a maltoside of *l*-mandelic acid, and on partial hydrolysis yields amygdalinic acid, which is a mixed glucoside of *d*- and *l*-mandelic acids. By agitating amygdalin with baryta solution, Dakin has converted it into an isomeride, *iso*amygdalin ($[\alpha]_D - 47.6^\circ$), which yields both *d*- and *l*-mandelic acids on complete hydrolysis. Walker had previously suggested that some such partially racemic substance as this is an intermediate product in the conversion of amygdalin into amygdalinic acid.

Bowack and Lapworth³ find that *l*-menthyl cyanoacetate,

and several of its derivatives do not exhibit mutarotation; the specific rotatory power of these substances does not afford evidence of changing molecular structure, although such evidence is found in the mutarotation of derivatives of *l*-menthyl acetoacetate.⁴ Only slight mutarotation was observed by Hann and Lapworth⁵ to occur amongst the *l*-menthyl alkylideneacetoacetates.

Lowry has shown⁶ that the equilibrium proportion in which two dynamic isomerides exist in a solution may be ascertained by saturating a solvent with the stable form, and at once determining how much (*a*) has gone into solution, then allowing the solution to remain in contact with the stable form until equilibrium has been established between the dynamic isomerides present in the solution, and again determining the composition (*b*) of the solution; the ratio $a/(b-a)$ represents approximately the proportion in which the isomerides are present in the equilibrium solution. In the case of solutions of nitro- and π -bromo-camphors, the normal and ψ -forms are in equilibrium when present in the solution in the ratio 5:1 approximately. On applying a similar method to glucose and galactose, Lowry found⁷ that the

¹ *Trans.*, 1903, **83**, 472.

³ *Ibid.*, 42.

⁵ *Ibid.*, 1904, **85**, 49.

⁷ *Ibid.*, 1551.

² *Ibid.*, 1904, **85**, 1512.

⁴ *Ibid.*, 1903, **83**, 1117.

⁶ *Ibid.*, 1541.

proportion of α -sugar present in the equilibrium solution decreases as the amount of water in the solvent increases; thus in methyl-alcoholic solution one-half of the sugar is present as the α -form, whilst in the mixture $\text{EtOH}, \text{H}_2\text{O}$, only 40 per cent. is of the α -form. It is suggested that the α - and β -forms are present in about equal proportion in the solutions, but that in the aqueous solutions a third hydrated and intermediate form is produced, thus:

E. F. Armstrong and Arup¹ have made careful determinations of the velocities of hydrolysis of α - and β -glucose pentacetates, β -galactose pentacetate, and sucrose pentacetate by potash. The results obtained are very concordant, and indicate that the velocity and course of the hydrolysis of α - and β -glucose acetates are identical, but that galactose pentacetate hydrolyses considerably more rapidly. The values obtained for octacetylsucrose are of peculiar interest, in that they indicate no distinction between the behaviour of the acetyl groups in the sucrose derivative and the glucose from which it is derived. The velocity coefficient in each case diminishes as the time increases, indicating that the acetyl groups do not all separate at the same rate. Determinations of the velocity of hydrolysis of the tetracetates of α - and β -methylglucosides and of α - and β -methylgalactosides show that all four substances are hydrolysed at the same rate; this is important, in view of the fact that in these isomerides the acetyl groups occupy similar positions in the molecules. The interconversion of the *d*-glucose α - and β -pentacetates has been studied by C. L. Jungius.²

T. Purdie and J. C. Irvine have shown³ that on methylating tetramethylglucose, tetramethyl- β -methylglucoside (m. p. 42—43°; $[\alpha]_D$ in water, -11.6°) is the main product, a liquid stereoisomeric tetramethyl- α -methylglucoside ($[\alpha]_D$ in water, $+147.4^\circ$) being formed in small quantity. The β -glucoside contrasts with the α -isomeride in its more rapid hydrolysis with acid and by being hydrolysed by emulsin.

Tetramethylglucose exhibits mutarotation; the α -form (m. p. 88—89°; $[\alpha]_D + 100.8^\circ$) is produced by repeated crystallisation from petroleum,

¹ *Trans.*, 1904, **85**, 1043.

² *Proc. K. Akad. Wetensch. Amsterdam*, 1904, **6**, 779.

³ *Trans.*, 1904, **85**, 1049.

and on heating above its melting point yields the β -form ($[\alpha]_D + 73.1^\circ$), which on preservation reverts to the α -modification. The form stable in solution ($[\alpha]_D + 83.3^\circ$) appears to be an equilibrium mixture of the two. Since mutarotation is also observed in benzene and carbon tetrachloride solutions, the assumption of ionisation or the formation of hydrates is not essential for explaining the phenomena.

R. Behrend and P. Roth have observed also that *d*-glucose shows¹ mutarotation in pyridine solutions, and state that the specific rotatory power of *d*-glucose in pyridine is at first $[\alpha]_D + 138.88^\circ$, but that during the lapse of 24 hours this value falls to $+71.17^\circ$; they conclude that this fact excludes the possibility of the mutarotation being due to the formation of hydrates.

J. C. Irvine and A. Cameron² obtain a liquid tetramethyl- α -methylgalactoside ($[\alpha]_D$, in water, $+143.4^\circ$) by the silver oxide method of alkylating α -methylgalactoside; on hydrolysis with hydrochloric acid, it yields a liquid tetramethylgalactose, which exhibits mutarotation in aqueous, alcoholic, and benzene solutions. On methylation, tetramethylgalactose yields stereoisomeric tetramethyl- α - and - β -methylgalactosides; the α -isomeride is the main product when methyl alcohol and hydrogen chloride are the methylating agents, whilst alkylation with silver oxide and methyl iodide gives mainly the β -isomeride. The latter is crystalline (m. p. $44-45^\circ$) and is more readily hydrolysed than the α -isomeride with dilute acids or emulsin.

Tutin and Kipping³ demonstrate the probability that the so-called *d*- and *l*-menthones produced from menthol are stereoisomeric, but not enantiomorphously related. They are interconvertible by passage through an enolic form, thus:

From *l*-menthone, by reduction of the oxime or by heating with ammonium formate, a mixture is obtained of the four possible stereoisomeric bases containing three asymmetric carbon atoms, one of which remains of fixed sign. The bases are named *l*-menthylamine ($[\alpha]_D - 61.9^\circ$), *l*-neomenthylamine ($[\alpha]_D - 17.4^\circ$), *l*-isomenthylamine

¹ *Annalen*, 1904, **331**, 359.² *Trans.*, 1904, **85**, 1071.³ *Ibid.*, 65.

($[\alpha]_D + 22.7^\circ$), and *l*-isoneomenthylamine ($[\alpha]_D - 3.8^\circ$), the numbers quoted being the specific rotatory powers of the corresponding benzoyl derivatives. The separation of the components of the basic mixture was effected by crystallisation of the hydrochlorides, bromocamphorsulphonates, camphorsulphonates, and the benzoyl and formyl derivatives.

Considerable activity has prevailed during the past year in connection with the stereoisomerism of derivatives containing tervalent nitrogen and of quaternary ammonium derivatives. H. O. Jones has published¹ an exhaustive summary of our previous knowledge of the stereochemistry of nitrogen.

A. Ladenburg² has continued his study of the new kind of stereoisomerism of tervalent nitrogen to which he has previously referred.³

He resolves synthetic stilbazoline, , into its

optically active components by the aid of *d*- and *l*-tartaric acids, and obtains the *d*- and *l*-stilbazolines, having $[\alpha]_D + 12.16^\circ$ and -11.5° at 20° . He then heats *l*-stilbazoline at 300° , and finds that it is converted into a mixture of *d*- and *l*-stilbazolines and a new isomeride, *l*-isostilbazoline. The resolution of the mixture is effected by first removing the *l*-stilbazoline as its *d*-hydrogen tartrate, and converting the base recovered from the residue into its neutral *l*-tartrate, when *l*-isostilbazoline *l*-tartrate separates. Salts of *l*-isostilbazoline were also isolated by another method of separation depending on the use of *d*-camphorsulphonic acid, and in each case the purified salt yielded *l*-isostilbazoline of $[\alpha]_D - 5.7^\circ$. From this fact and from differences observed between the behaviour of the corresponding salts of the stilbazolines and *l*-isostilbazoline, Ladenburg concludes that the latter is not a mixture, but a true stereoisomeride of stilbazoline.

Kipping and Salway⁴ refer to the discrepancies between the theory of Hantzsch and Werner,⁵ which explains the isomerism of the oximes by assuming that the three valency directions in tervalent nitrogen do not lie in one plane, and the failure of previous attempts to resolve bases in which three different groups are attached to the tervalent nitrogen atom; they again point out that the latter experiments are indecisive owing to the possibility of a change in the valency directions attending the conversion of ter- into quinqué-valent nitrogen and *vice*

¹ *Brit. Assoc. Reports*, 1904.

² *Ber.*, 1904, **37**, 3688.

³ *Ibid.*, 1895, **28**, 854.

⁴ *Trans.*, 1904, **85**, 438.

⁵ *Ber.*, 1890, **23**, 11.

versâ. They treated *d*-hydrindamine, *l*-methylhydrindamine, *l*-menthylamine, and *l*-phenethylamine with *d*-benzylmethylacetic chloride, and in each case obtained but one acetyl derivative of the primary or secondary base; if the three nitrogen valency directions do not lie in one plane, the production of stereoisomerides would be anticipated. It is therefore concluded that the three valency directions of tervalent nitrogen lie in the same plane.

It is pointed out that on treating an optically inactive base with an externally compensated acid chloride, only one acidic derivative should be formed if the original base is potentially inactive, whilst two isomerides are capable of formation if the base is externally compensated. Thus, *dl*-hydrindamine yields two isomeric acidic derivatives with *dl*-benzylmethylacetic chloride, whilst methylaniline, *p*-toluidine, benzylaniline, and phenylhydrazine yield but one product in each case. Some months later, E. Mohr described¹ the same method for ascertaining whether a base is potentially inactive or whether it is externally compensated. In studying the following reaction,

using externally compensated materials, he found that two isomeric compounds were formed; this is, of course, due to the potential optical activity of both the reacting substances.

Externally compensated hydrindamine may be resolved by treatment with *d*-benzylmethylacetic chloride; the least soluble of the two products of the reaction is *d*-benzylmethylaceto-*d*-hydrindamide. Similarly, *dl*- α -phenethylamine is resolvable by the aid of the same acid chloride, and *d*-benzylmethylaceto-*l*-phenethylimide is readily separable from the product. In these two cases, it was found impossible to regenerate the optically active bases by the hydrolysis of the acidic derivative.

The question as to the distribution in space of the three valency directions of a tervalent nitrogen atom has been also dealt with by H. O. Jones and J. P. Millington.² These workers were unable to resolve either phenylbenzylhydrazine, $\text{CH}_2\text{Ph}\cdot\text{NPh}\cdot\text{NH}_2$, by the aid of *d*-camphorsulphonic acid, or methylaniline-*p*-sulphonic acid by crystallisation with cinchonine.

H. O. Jones³ prepared phenylbenzylmethylethylammonium iodide, introducing the alkyl groups in three different orders, namely, by combining benzyl, methyl, and ethyl iodides respectively with tertiary base, but obtained no evidence bearing on the occurrence of Wedekind's α - and β -isomerism. He resolved the quaternary iodide by Pope and Harvey's method,⁴ applying the silver salt of *d*- and

¹ *Ber.*, 1904, **37**, 2702.

² *Proc. Camb. Phil. Soc.*, 1904, **12**, 489.

³ *Trans.*, 1904, **85**, 223.

⁴ *Ibid.*, 1901, **79**, 76.

l-camphorsulphonic acids in a water-free solvent. The sparingly soluble salts are *d*-B *d*-A and *l*-B *l*-A, and give $[M]_D + 71.0^\circ$ and -71.2° in aqueous solution. The basic ions therefore have $[M]_D +$ or -19.5° . The optically active iodides give $[\alpha]_D + 8.3^\circ$ and -8.4° in alcoholic solution; they undergo autoracemisation in chloroform solution, much as do the *d*- and *l*-phenylbenzylmethylallylammonium iodides. Crystallographic measurements showed that the inactive iodide is either pseudoracemic or a mechanical mixture. The corresponding *l*-bromide gives $[\alpha]_D - 13.4^\circ$ in alcoholic solution; its rotatory power diminishes on recrystallisation from alcohol. Thomas and Jones¹ have also resolved phenylbenzylmethylisopropyl- and *iso*amyl-ammonium iodides into their enantiomorphously related components.

E. Wedekind and F. Oberheide² have prepared the following quaternary ammonium iodides containing asymmetric nitrogen atoms: *p*-tolylmethylallylammonium iodide, benzyl-*p*-tolylmethylallylammonium iodide, benzyl-*p*-tolylethylallylammonium iodide, and benzyl-*o*-tolylmethylallylammonium iodide, but have not resolved them into optically active components.

A. W. Harvey³ has prepared phenyldimethylallylammonium iodide from dimethylaniline, but was unable to resolve the corresponding *d*-camphorsulphonate into fractions containing an optically active basic ion.

H. O. Jones⁴ has obtained two stereoisomeric phenylmethylallyl-*l*-amylammonium iodides by the combination of methyl-*l*-amylaniline with allyl iodide; one of the products is dextrorotatory and the other levorotatory. In chloroform solution, the rotatory powers of the salts gradually change until a constant value is reached owing to dissociation, which causes optical inversion of the asymmetric nitrogen atom. More recently, M. Scholtz has extended⁵ Jones's study of compounds containing both an asymmetric carbon and nitrogen atom. The combination of *n*-ethylconiine with benzyl iodide leads to the formation of two isomeric benzylethylconinium iodides of different melting points and rotatory powers. Similar results were obtained on combining *n*-iso-amylconiine with benzyl iodide and *n*-benzylconiine with methyl iodide. Scholtz, however, has not observed any change of rotatory power of the isomerides such as is recorded by Jones.

On treating *dl*-dihydro- α -methylindole with *d*-bromocamphorsulphonic acid, W. J. Pope and G. Clarke⁶ obtained *l*-dihydro- α -methylindole *d*-bromocamphorsulphonate ($[M]_D + 278^\circ$ in water) as the least soluble salt; the basic ion therefore has $[M]_D + 5^\circ$, and on treating the salt with alkali, *l*-dihydro- α -methylindole was obtained. A second

¹ *Proc. Camb. Phil. Soc.*, 1904, **13**, 33.

² *Ber.*, 1904, **37**, 2712 and 3894.

³ *Trans.*, 1904, **85**, 412.

⁴ *Proc. Camb. Phil. Soc.*, 1904, **12**, 466.

⁵ *Ber.*, 1904, **37**, 3627.

⁶ *Trans.*, 1904, **85**, 1330.

salt of $[M]_D + 242.7^\circ$ was isolated, but this does not give active base with alkali despite the low value of the molecular rotatory power; it is suggested that this salt really contains the active base, but that during the action of the alkali tautomeric change takes place in such a way that the optical activity of the base is lost.

Tattersall finds¹ that *dl*-hydrindamine is conveniently resolved by first crystallising it with *d*-tartaric acid, when *d*-hydrindamine hydrogen *d*-tartrate is obtained pure, and crystallising the base recovered from the mother liquors with *d*-bromocamphorsulphonic acid; the latter process yields *l*-hydrindamine *d*-bromocamphorsulphonate in a state of purity. On crystallising *d*- and *l*-hydrindamines with *d*-chlorocamphorsulphonic acid, indications of the production of more than one salt from each base are obtained, just as when *d*-bromocamphorsulphonic acid is used.²

Lastly, it should be noted that Kipping has prepared the asymmetric silicon compound, phenylethylpropylsilicon chloride, PhEtPrSiCl , but has not yet announced its resolution into components owing optical activity to the presence of an asymmetric silicon atom.³

W. J. POPE.

¹ *Trans.*, 1904, **85**, 169.

² Kipping and Tattersall, *ibid.*, 1903, **83**, 918.

³ *Proc.*, 1904, **20**, 15.