nouncement of Koch's discoveries, was a lamentable misfortune. The length of this address already reached compels a simple reference to the present state of information concerning the other infectious diseases.

It is conceded that the bacillus of diphtheria of Loeffler is the cause of that disease. Failures to find it have arisen from a mistake in diagnosis, or lack of skill in technique, or failure to search for it in the proper position. The practical suggestions are made, that as it may develop in cases of simple pharyngitis, and may be found in the mouth days after the malady is cured, great care in local antiseptic measures is indicated in all cases of pharyngitis.

Special interest is aroused concerning tetanus, from the fact of Kitasato's success in making pure cultures of the bacillus and placing it among bacillary diseases beyond question; the conclusions of Bosaud are that almost all soil contains the bacillus tetanus, and that its presence depends upon defective drainage, hygienic conditions and degree of cultivation of the soil.

In regard to hydrophobia two things are to be said; one, that the best judges believe it due to one or more microorganisms not demonstrated, the other that Pasteur's protective inoculations do protect under favorable conditions.

Klein's review of Koch's cholera report evidently shows Koch mistaken in claiming the comma-bacillus always present in cholera, and establishes the claim that this bacillus is both found in other cases and has not been demonstrated.

The Cause of Cholera.—Woodhead reviews the ground carefully in a most judicious manner, and concludes that it is extremely probable that the comma-bacillus is the cause of cholera.

It is practically conceded that typhoid fever is due to the presence and growth of the bacillus of Eberth and Klebs. No progress has been made in preventive inoculation. It grows in water and more freely in milk.

The ray fungus is the accepted cause of actinomycosis, and specific bacilli are claimed for syphilis, leprosy and glanders, the remaining members of the infective tumors, so-called, the class in which tuberculosis belongs.

The relation of bacteria to wound infection, and to puerperal fever, is considered so firmly established, that a mere repetition of the surgical aphorism, that "he who has suppuration in plastic surgery is either a poor or a careless surgeon," is sufficient in this connection. The amount of work which has been done, and the literature which represents it, are very large. It is the direction in which at present the greatest hope lies for progress in preventive medicine. This is the appropriate time for oratorical pyrotechnics in summing up the benefits, but you are spared more than a mere mention of the names of those to whom the greatest debt of gratitude is due. There names are Semmelweiss, Cohn, Pasteur, Lister and Koch.

POST PARTUM HÆMORRHAGE.

Read before the Milwaukee Medical Society, May 10, 1892.

BY S. W. FRENCH, M.D.,

OF MILWAUKEE, WIS.

Perhaps I should apologize to the society for bringing before them a subject which at first sight seems so simple; but post partem hæmorrhage does not come under our hands, fortunately, very often, but

when we do encounter it, it is a condition that needs immediate attention, and no delaying or expectant plan whatsoever. I therefore bring up the subject more for discussion than for any intrinsic value that the paper may have to the society. It has been my fortune during a practice of thirteen years in this city to meet with but few cases of post partem hæmorrhage. Of course we often meet with cases that are of a slight character, which after a thorough massage of the uterus, or a good free injection of hot water, stop within a short time. But I refer to cases where the hæmorrhage is particularly severe. The only really serious case of post partem hæmorrhage that I ever had, occurred in my practice within the last month. I say serious, because it was at least eight hours after delivery before I felt that my patient was in a safe condition. The delivery was simple. It was her second child. The labor was very light indeed, and the expulsion was entailed with but little difficulty, although there was some slight laceration of the perineum, enough, however, in my belief, to put in a few sutures. She had had no anæsthetic and it was therefore a little difficult to bring the two parts in apposition. After the birth the uterus seemed to have contracted well, and the placenta was delivered. I thought things were then in a safe condition, and in order to dam back a little flow that was taking place I put a small tampon of cotton into the vagina. I then introduced my sutures, and after considerable difficulty in the case, I finished the operation. The cotton was extracted and the pulse examined, but instead of finding it reduced, I found it ranged above 120. I immediately felt for the uterus and found the fundus very near to the epigastrium. I then began massage, and I think I squeezed out in the vicinity of about two quarts of clots. The woman complained a great deal of faintness, but soon as I let go of the uterus, away it bulged up again to the self-same locality as it was before, and by continuous massage again I expressed out the same amount of clots; and it began to be a question in my mind as to how long this thing was going on. I suddenly remembered having read in a journal some few weeks before of perhaps a similar condition, in which the physician did exactly the same thing as I had done, but to the extent that he attempted to keep the uterus down. The consequence was that in keeping it down, however, there was still a considerable hæmorrhage, and eventually the patient died. He drew the conclusion that it would have been better if the uterus had been left as it was and the clots had not been expressed. This suddenly came to mind and I resolved to act upon it; and the third time when the uterus bulged up above the umbilicus, I concluded to let the thing alone and not try to express it. However, I did give a teaspoonful of ergot and some whiskey. The pulse, however, kept up above 120, and was at times intermittent. After that there seemed to be no active hæmorrhage. The woman became collapsed. All of this happened at 9 o'clock in the morning. At about between 11 and 12 I thought I detected some change for the better, and left the case to see my other patients; but I had not been absent more than three quarters of an hour before I was telephoned for to come immediately, as the patient had fainted. But as usual in those cases, when I arrived, I found that the patient had recovered from the faint, but that the pulse was still in a very uncertain and irregular condition. I

then gave the patient $\frac{1}{30}$ grain hypodermically, of strychnia, followed by $\frac{1}{120}$ of atropia. I also gave hypodermically a number of injections of whiskey. This was at 2 o'clock, and by 5 the pulse had assumed a normal beat. She went on to convalescence, but it was somewhat retarded on account of the extreme anæmia that was present. In this case I used a hot water injection, and it was the first time in my practice in which hot water did not have the effect to produce a good, firm contraction of the uterus. We know there are many things that have the power of contracting the uterus. We know that there are some who believe entirely in the use of vinegar passed into the uterus on a swab. We know that the same effect is sometimes produced by squeezing a lemon into the cavity of the uterus. We know that there are some who believe entirely in the use of the fluid extract of ergot, or in ergotine injected hypodermically. We know that there are others who believe entirely in the use of hot water. But I think that the method that I pursued in this case is a method that is not generally laid down in text books. I think that the general rule laid down in text books is to clear the uterus and keep it well contracted down. Under the condition existing in my case it was impossible to keep that uterus contracted down, and it flashed through my mind that the very best way to stop that hæmorrhage was to bring about the condition of things that nature herself was attempting to do, and that was to produce a large clot in the uterus, and by the presence of that large clot a contraction would take place. I will say this, that the contraction was slow, that it was not until the next day at noon that the uterus was down in the condition that you would have wished to have found it after delivery.

Post partum hæmorrhage necessitates not only a careful but also a rapid survey of the conditions. The question is what is the cause of post partum hæmorrhage, or especially what is the cause of the post partum hæmorrhage in your own individual case. It is a question that I think you cannot always answer, because if you could answer immediately, you would then probably be better able to select your remedy for stopping it. It more often comes like a thief in the night and takes you entirely off your guard. Now, in respect to this woman, she was the last person that you would ever expect a post partum hæmorrhage in. The labor had been unusually rapid, the contraction of the uterus had been good during the whole labor. There was no delay, the blood clotted, which does not always occur. I think it has been brought up by some authors that in these severe post partum hæmorrhages the blood does not always clot. I think it was fortunate in this case, that the blood did clot, and that that clot in the uterus did act as a tampon, so to speak, to close up the uterine sinuses. That word, tampon, brings to mind that it has been considered a good plan in severe post partum hæmorrhage, where hasty means are used, in the way of hot water and vinegar, and the hæmorrhage seems to keep up, to tampon the whole uterus with antiseptic gauze, and in other words attempt to bring about a clot by this self-same pressure. Of course, the use of the gauze, being introduced into a freshly parturient uterus in that way, is a method attended with a great deal of danger, especially of producing sepsis. Another point that came to my mind was that if you did not produce the sep-

sis, you might produce an absorption of your antiseptic, if you used an antiseptic gauze. If a sublimate gauze was introduced into a cavity of the uterus with a freshly denuded surface, it might have the effect of producing sublimate poisoning. The idea is to pack the uterus firmly, as if you were going to tampon the vagina. The matter of post partum hæmorrhage is a serious thing and requires the most rapid and heroic treatment possible, and it is a little difficult sometimes to select the most æsthetic means at your hands, and possibly oftentimes difficult to select a method that would be free of danger in itself, more particularly a perfectly septic one.

DISCUSSION.

Dr. W. H. Washburn: There are two classes of patients in whom we are most apt to have post partum hæmorrhage of a serious character. First, those cases in which labor is unusually rapid and precipitate, and second, those in which the labor is protracted and difficult. In the first case the uterus seems to be, as it were, paralyzed by its heroic efforts to empty itself, and in the second case the hæmorrhage is caused by prolonged efforts in the same direction. The case that Dr. French has detailed seems to be one of those that belong to the first class, where the uterus is paralyzed by a sudden and great effort. So far as I am individually concerned, I should dislike very much to leave a patient with the uterus dilated and filled with clotted blood; I could not feel safe for the future of the patient, and it does not seem to me as though that is very good treatment. To be sure, the uterus fills up with a clot, but that does not prevent hæmorrhage from taking place from the uterine sinuses, and I should think it would be a precarious thing to do. I have had one case of this kind where hæmorrhage occurred in a woman in which the labor was extremely rapid. In fact, the child was born before my arrival, and the woman was in a condition of syncope from loss of blood when I reached the house, and everything around about was perfectly deluged with blood. In that case I remember very distinctly that she had had a previous experience of the same kind, and her husband when he engaged me to attend her, told me the circumstances of the case. In that case I remained with the woman all night. I gave her stimulants immediately on arrival, and succeeded in emptying the uterus partly by pressure and partly by introducing the hand into the uterus and breaking up the clots. I kept my hands on the uterus all night. I stayed there from nine o'clock at night until daylight next morning, and that was in the winter. The woman recovered rapidly.

Another case occurred after a prolonged and difficult labor in which instruments were used. I was present as consulting physician and soon after the child was born and while I was in the kitchen with the child attempting to resuscitate it, the physician who was in attendance called me, and he was so frightened that he could not tell me what he wanted me for. When I got into the room the perspiration was dropping off his forehead. I never saw such a picture of terror; and the woman exclaimed, "For God's sake, Dr. Washburn, don't let me bleed to death;" and certainly she would have bled to death in a very few minutes, for when I turned back the bed clothes there was a steam of blood coming from her as large as my two fingers. I at once introduced my left hand into the uterus and made pressure over the abdominal walls with the right hand, and got her husband to go out and get a piece of ice, which I introduced into the uterine cavity. The irritation of my hand in the uterus caused strong contraction, and immediately after the ice was introduced the hæmorrhage ceased, and did not recur. The woman made a good recovery.

Of course there are some cases that resist all kinds of treatment. As to the injection into the uterine cavity of such substances as perchloride of iron and any substance that causes such a firm clot as that, I should not feel satisfied to try them myself. I think that in the majority of cases the introduction of the hand into the uterus, massage over the abdomen, the introduction of ice into the uterus, or hot water injections, which at the present time perhaps are more preferable and least liable to be followed by septic trouble, would be the most appropriate treatment, and treatment which would be successful in the largest number of cases.

Dr. E. Copeland: I would agree with Dr. Washburn as

regards the use of substances which cause such a firm clot as perchloride of iron would. I would not feel comfortable or satisfied to adopt such a procedure. I certainly believe in the hot water injections in a great number of cases, if not all, using also strong pressure on the uterus of the patient, continued for hours if necessary, which one physician cannot do if it is necessary to keep it up for any great length of time. A man's hands will certainly tire out and he cannot do that and must call help; but those are very rare cases; I never met but one case and in that case pressure was kept up for several hours. I think in the great majority of cases of post partum hæmorrhages, and they are not very common, hot water injections constantly kept up for a long time if necessary, will control the hæmorrhage. So far as the introduction of vinegar, lemon juice and things of that kind into the uterus is concerned, I would not think of it for a moment.

Dr. Puls: Post partum hæmorrhage is an accident which happens not so very seldom after all. I remember of having had a number of cases. Of course the causes differ, but in most cases the hæmorrhage is due to atony of the uterus, and simple massage and hot water injections will suffice generally to stop the hæmorrhage. Of course, if you have a case of retention of parts of the placenta, the only course is to remove that part of the placenta to stop the hæmorrhage. Another source of the hæmorrhage is tearing of the cervix or perineum which must not be overlooked; if it cannot be remedied by sutures, the tampon will probably be the only way; a hot water douche would not stop the hæmorrhage. Kuestner lately reported a case of post partum hæmorrhage due to aneurysmatic changes of the artery and also to arterio-sclerosis as being a cause of hæmorrhage. The blood would not clot. Now, in most of these cases, wherever we have a rent of the cervix, or where we have no clotting of the blood in the uterus to stop the hæmorrhage, it is natural to tampon the uterus; and if Dr. French succeeded in his case in saving the patient, it was due to the clots. As Dr. Washburn well remarked, by introducing his hand into the uterus he induced a contraction of it to take place, and by applying the ice he got firm contraction. I do not advise the use of ice, but I advise the use of iodoform gauze to pack the uterus. It is a method adopted by Duehrssen about three years ago, and since then probably two hundred cases have been reported altogether treated in that way, with no case of septicæmia resulting. Nor has any case of iodoform poisoning followed. In my practice I have used that method probably three or four times with no septicæmia following and without any fever. The best way if you have no iodoform gauze on hand, is to take ordinary cheese cloth prepared in carbolic solution and apply that. I remember a case only two months ago where the blood came oozing out of the uterus as thick as one finger, and I felt that the woman was dying right under my hands. It was a very rapid delivery; I had nothing at all to introduce into the uterus except my hand, and I called for the basin in which I had the cheese cloth ready, and I took one of those cloths and packed it into the uterus and the hæmorrhage stopped, and I extracted the cloth the next day. I certainly would advise this method in cases of natural birth or after an abortion, not waiting to employ ergot.

Dr. Burgess: When a hæmorrhage occurs anywhere, when it has been of sufficient extent, the patient faints. In that condition the energy of the heart relaxes, and the treatment followed by the laity as long as whiskey has been known, is to pour in whiskey and other stimulants to start the heart up to pump more blood out. It struck me when Dr. French spoke of using whiskey in that case, that it was not good treatment for the reason that you only stimulate the heart to greater exertion, and perhaps, in that way increase the hæmorrhage. If a person faints it does no harm, but does good. Why not? It is the exact condition you want to bring about.

In regard to tamponing the uterus, that may be well enough under certain conditions in proper hands; but I think if you set everybody who attends to midwifery cases to tamponing the uterus to stop hæmorrhage, he will do it with his fingers and it will not be half done, and if he uses instruments he will be very likely to stick them through the fundus of the uterus occasionally. It is a valuable measure perhaps, but must be used with circumspection.

Dr. Shimonek: I have had quite a number of cases in my practice. I have always succeeded very well in stopping hæmorrhage by massage, and stimulation of the uterus internally by the hand. I had one severe case among those, that very nearly bled to death, a case in the country that I was called in to see. The woman finally recovered. I found

the uterus filled. I expressed the clots, contraction took place and the woman recovered. But I certainly should not advise the treatment proposed by Dr. French in regard to allowing the clots to remain in the uterus. I do not think that is very good treatment. I should not rely upon it. I do not know what effect the doctor got from the injection of strychnia and atropine, but it seems to me that it would be likely to produce more serious hæmorrhage from increasing the circulation. As soon as I can I always inject ergotin hypodermically.

Dr. French: In respect to the injection of strychnine and atropine, that was some hours after the hæmorrhage had ceased, and also the use of the whiskey was at the same time. This woman was in a state of collapse, and it was a great question as to whether she was going to pull through from one minute to another. Her extremities were cold. She was ice cold all over, and although the hæmorrhage had stopped, still with the pulse missing a beat about once every third beat, and when I could not tell whether the pulse was in her wrist or in my fingers, it seemed to me that it was necessary to use some stimulants to keep the action of the heart up. I readily agree with Dr. Burgess in the matter of the use of stimulants under the fainting condition where the hæmorrhage is in its active period; but there is another point to be considered in that respect; that although the whiskey may produce an increased action of the circulation, there comes a question as to whether it will not also stimulate the worn out condition of the uterine muscles. I well remember an incident related by old Dr. Buckingham, the professor of obstetrics, where he gave a woman ten teaspoonfuls of laudanum and produced a contraction of the uterus immediately, or within a very short time. I would like to ask Dr. Puls how long he would keep the gauze in?

Dr. Puls: Twenty-four hours, if necessary.

Dr. French: As I say, I see that the gentlemen would not agree with me in this method of treating this case. As I said in my paper, it was directly contrary to rules I had followed in other cases, but it seemed to me that this case was one that it seemed necessary to treat in this way if it would not yield to other methods of treatment. Dr. Copeland has spoken about sending for assistance. When you get caught in a case of this kind, unless assistance is next door or around the corner, it is pretty difficult to get the assistance there to help you.

THE INFLUENCE OF BODILY EXERCISES UPON LENGTH OF LIFE.

Read at the Annual Meeting of the Association for the Advancement of Physical Education, April 12, 1892.

BY J. MADISON TAYLOR, A.M., M.D.,
OF PHILADELPHIA.

It is not possible to estimate statistically just how much length of days is enhanced by physical exercises, but it is feasible to establish two propositions: (1) That judicious activities of the body tend to maintain and increase its efficiency, and (2) That the hurtful effects of violent athletic competitions are popularly overrated.

It would be most agreeable to me to consider the first of these, and drawing from history and experience, illustrate amply what few will be disposed to deny. Concerning the second there may be differences of opinion, and here it is well to apply the test of carefully marshalled facts and suffer inferences to evolve themselves. I hope to escape the rôle of advocate, but propose reviewing such meagre data as I have been able as yet to collect, and deduce therefrom only warrantable conclusions. My study so far is most imperfect, but my hope is to elicit discussion and comment so that the truth may come ultimately forth. Within the coming year I shall hope for much light and added knowledge. I have reason to expect large additions to the data yet collected, and these may have a value of which you shall judge.¹

¹ Dr. Taylor will be gratified for any information connected with this subject.