tiary in an official capacity and the result of my observation is that 70 per cent. of 1,940 convicts come there directly or indirectly from the use of alcohol. I mention that as regarding the moral side of the question which I have heretofore avoided discussing. One gentleman used the burning of an alcohol lamp as an illustration; on the same principle why not use gasolin or make a decoction of coal products and other things we use outside of the body, for oxidization. I do not believe alcohol can be oxidized in the system. It can be burned outside. We would not think of inhaling natural gas because we use it for illuminating purposes. A few inhalations would satisfy us, I think. I have followed it with much interest, but I yet believe it is a bad thing to use alcohol in the treatment of typhoid fever.

TRANSFUSION, INFUSION AND ANTO-TRANSFUSION; THEIR COMPARATIVE MERITS AND INDICATIONS.

Read before the Kentucky State Medical Society.

BY AUGUST SCHACHNER, M.D., PH.G.

DEMONSTRATOR OF ANATOMY LOUISVILLE MEDICAL COLLEGE; VISITING SURGEON LOUISVILLE CITY HOSPITAL. LOUISVILLE, KY.

Whenever a number of measures of a more or less varied and changeable character are advanced for the fulfillment of a long existing demand, it means an absence of unity in the selection of the proper measure and a general distrust in a satisfactory adjustment of the demand. This is clearly illustrated by the number of procedures that from time to time have been recommended to overcome the depressing and even fatal condition following enormous hemorrhages.

The fact that the operation of transfusion is but rarely performed at the present time, and that of infusion faring but little better, tempted me to refreshen the subject somewhat more than a month ago before one of our local societies. Since that time I have been impressed more than ever with the neglect to which these operations have been subjected and the importance of bringing them before the eyes of those engaged in operative work.

I may be pardoned for borrowing from my former paper the historic outlines in connection with the development of these minor operative procedures.

The first intimation of transfusion can be found in the 333d and 334th verses of the seventh book of Ovid's *Metamorphoses*, "*Veteremque haurite cruorem Ut replam vacuas juvenili sanguine venas.*" This carries us back to the time of Christ, and from then down to the present period; "a long but unbroken chain can be traced running through the Hebraic and Egyptian medical records." In 1492, or more than a hundred years before the circulation was understood, Pope Innocent VIII. was struggling with his last illness. He was attended by a Jewish physician who, it is supposed, was prompted by the idea noted in the verses of Ovid and performed transfusion with the blood taken from three Roman youths. In 1615 Andreas Libavious, of Halle, wrote an article upon a charlatan who is supposed to have performed transfusion. In 1628 another appeared under the authorship of Johann Colle; coupled with these came the discovery of the circulation by Harvey, which was directly responsible for the vigorous attention which the learned men at that time gave to the subject.

In 1652 an apparatus for arterio-venous transfusion was contrived by Folli, of Italy. In 1666, Richard Lower drained a good sized dog by tapping the jugu-

lar vein. When the animal was exhausted he filled the vessels with blood drawn from a cervical artery of a second dog until the animal had recovered; he then drained the same animal a second time and again filled his arterial system with blood from a third dog, thus completely changing the blood twice in the same animal without any unfavorable consequences. This represents the first well-authenticated experimental maneuver made in the direction of transfusion.

This aroused an interest in the Royal Philosophical Society, and it was then believed that a remedy had been discovered which was not only capable of curing disease but by means of which the aged could be transformed to the youthful and the immoral could be changed to the moral. So strong was the belief in the latter, that a trial was actually undertaken by Lower and King.

At that time a religious fanatic by the name of Arthur Boga, 30 years old, offered himself in consideration of a guinea as willing to undergo the experiment, which was conducted, in the presence of the Bishop of Salisbury and a large and brilliant audience, by Lower and King. Six to seven ounces were withdrawn and ten ounces of arterial blood from a sheep was injected. This operation was again successfully performed on the twelfth of December of the same year by the same operators.

It must be noted, however, that these experiments were preceded by others performed by Jean Dennis and Emmerez in Paris in the year of 1667, and to whom the honor of the first successful transfusion in a human subject is due. Dennis employed the blood of a lamb, and several times repeated the operation upon several different subjects with almost uniform success, so far as the transfusion was concerned. These operations aroused a vigorous and jealous opposition in Germany, France and Italy, and were partly terminated when Dennis narrowly escaped a trap that had been set for him, but notwithstanding his innocence, it had involved him in a criminal case.

From this time enthusiasm began to lag and matters continued with a varied interest until 1818, when James Blundell, the obstetrician, placed transfusion upon a scientific basis, at the same time giving a method for its correct performance. To Blundell is likewise due the honor of first employing human blood instead of that of lower animals for the transfusion. With this the history of transfusion can be dismissed. In parting we might add that there are but few procedures known to medicine whose histories are more replete with incidents that are as varied, interesting, pathetic and amusing as the history of transfusion.

Before entering upon the subject of transfusion, it is well to glance at the conditions which are produced by the loss of large quantities of blood.

In dangerous hemorrhages death may ensue from one of two causes, i. e., either from the absolute loss of blood itself or from a fatal reduction of the intravascular pressure. In the first instance, the amount of blood is insufficient to meet the demands necessary for the sustenance of life. This makes the case unmistakably clear, and the indications are more blood, or death must ensue. In these cases nothing short of transfusion will fulfill the requirement; fortunately, however, these constitute the minority. In the majority of cases the death following hemorrhage is not due to the direct loss of blood itself, but rather to a disturbance in the mechanism of the cir-

ulation. Let us glance at the physiology of the circulation.

Two factors must be considered: First and foremost, the force of the heart, and second, the elasticity and contractibility of the arteries. If the heart has an insufficient volume of blood, or perhaps more properly speaking, volume of fluid, for blood is not absolutely necessary for this feature, there is an irregularity in its contractions and a serious crippling in the proper working of its valves, and in consequence we have a condition at once produced which for the time being, can be compared to the worst variety of valve lesion, so that the heart not only fails in sending out the required amount for nutrition, but also for the closure of the valves. Aside from this, the amount of blood sent out is not sufficient to produce the necessary dilatation by means of which the elastic nature of the large arteries are enabled to propel the force of the heart to the more distant parts of the body.

When this state of affairs exists, the indication for an increase of the intravascular pressure is equally plain and the requirement then is more fluid; whether that be blood or salt solution is immaterial.

TRANSFUSION.

By transfusion we mean the injection of the blood of one individual into the vessels of another. Although this operation has been variously designated by different writers as both safe and unsafe, there can be no doubt that in the hands of a careful operator, all possible elements of danger can readily be eliminated.

Indications.—From the history of transfusion it is apparent that formerly its range of application was far more extensive than its merits justified, and in view of this state of affairs the disappointments naturally were quite frequent. This largely aided in relegating it to the disuse to which it was subjected. At present any factor may serve as an indication for transfusion, which reduces the quantity of the blood so that the remaining volume is unable either by reason of the reduction, or incapable by reason of any alteration to which it may have been subjected, to carry on the functions necessary for the sustenance of life. The principal conditions which such a statement would comprise, would be either an acute anemia, dependent upon an extensive hemorrhage, or a paralyzation of the oxygen carrying power of the red corpuscles, carbonic oxid, or other similarly toxic gases. We would still scarcely regard such conditions as morphin or atropin poisoning, leukemia, chlorosis and a host of other pathologic states as justifiable indications for the employment of transfusion.

Modus operandi.—There are a few practical features in connection with the operation of transfusion which deserves special attention. During the cholera epidemic of 1866, Von Graefe raised the question as to whether the centrifugal should be given the preference over the centripetal method? Whether it would be better to inject the blood into the arteries rather than into the veins? The advantage urged in support of the centrifugal method was that since the capillaries intervened between the arteries and veins, the blood in its passage through the capillaries was subjected, as it were, to the influence of a strainer which safely withheld any air or emboli which might be present in the transfused blood. The arteries which were employed were either the radial, tibial or

even perhaps the brachial. Although this method has the endorsement of Billroth, Landois, Heuter and others, it has likewise many very able opponents who have pointed to the force necessary for the centrifugal method which not infrequently resulted in a rupture of the capillaries with accompanying gangrenous conditions.

Another question to be decided is whether or not the blood should be defibrinated and whether we should select the venous or the arterial blood? Referring to the former of these two questions, we are safe to say that the verdict is almost unanimously in favor of the defibrinated blood. It has been said that the integrity of the corpuscle is considerably affected by the defibrination of the blood, but this is somewhat problematic. However, the dangers of a partial or an entire coagulation which attended the use of non-defibrinated blood, fully justifies the defibrination.

As to the second question, there are reasons why the venous blood is to be preferred to the arterial, namely, the veins are more accessible than the arteries. The tapping of a vein is simpler and is attended with less subsequent dangers than accompanies the opening of an artery, besides the blood, although venous in character, rapidly becomes arterialized during the act of defibrination. In addition to these we might ask, should the transfusion be direct or indirect, i. e., should the blood pass directly from the vessels of the donor into the vessels of the receiver without any exposure to air? To this we are inclined to say that we believe it to be far safer to employ the indirect method.

The extreme tendency on the part of the blood to coagulation, and the grave result which attends the introduction of even a minute embolus, not to speak of the chances of the introduction of air, even at the hands of a careful operator, makes us unhesitatingly declare in favor of the indirect method.

Should the transfusion be undertaken for the relief of a poisoned condition of the blood, such as results from the exposure to carbonic oxid gas, the individual should be subjected to venesection before the transfusion is undertaken. In the performance of the operation no elaborate outlay of instruments is necessary. Two or three bowls, a reversible aspirator, a glass rod and a scalpel, all of which being in a perfectly aseptic state, will fill the requirements. Everything in connection with the operation must not only be perfectly aseptic but must be heated to 105 or 110 degrees F. When all is in readiness, the blood is withdrawn from the arms of one or two donors into one of the aseptic bowls, the amount varying from five to fifteen ounces, according to the circumstances of the case. It is rapidly defibrinated by whipping it with a glass rod and then straining through a piece of sterile gauze into an aseptic bowl. This whipping may be carried out for a minute and a half before straining, but still better is to whip for a minute and strain whipping, whipping it a second time and again straining through a fresh piece of gauze. The actual operation must be performed within two or three minutes, and the greatest care must be exercised that all of the air has been forced out of the syringe and tube before the injection is commenced.

Dangers of transfusion.—The possible dangers that may attend the operation of transfusion are: Phlebitis, sepsis, embolus and the introduction of air into the veins; all of these can be readily eliminated if the operator exercises care in the transfusion.

INFUSION.

By infusion we understand the introduction of non-sanguineous fluids into the circulation. Although the popularity of this measure is but of recent date its real history, however, can be traced as far back as the year 1677, when Johannes de Muralto of Zurich practiced the injection of milk into the vessels of one of the lower animals. The term infusion, however, as applied to-day carries with it the idea of a salt solution having the same strength as the serum of the blood. The advantages of this operation over that of transfusion are manifold, especially since death, in the majority of instances, is not due as much to the insufficiency of the remaining quantity of blood as it is to a disturbance of the mechanism of the circulation. By the use of the salt solution the dangers common to transfusion are all minimized and especially the dangers from emboli are entirely absent. This widens the field of its usefulness so that the indications for its use are not wholly confined to conditions following alarming hemorrhages, but include any pathologic state attended with a feeble pulse which is dependent upon a diminution of the intravascular pressure which makes it one of our most valuable measures for combating profound shock.

This property of restoring the tone of the circulation in a condition of shock is not entirely due to the increased intravascular pressure, but also due to the stimulating influence which the salt solution has upon the heart. This fact, however, has not received the recognition which it deserved, for it has been but a little more than a decade since it has practically received any attention.

In the year 1881, transfusion lost most of its esteem as a life saving measure in alarming hemorrhages. At this time E. Schwartz published his paper, "Ueber den Werth der Infusion Alkalischer Kochsalz Lösung in das Gefass System bei Acutes Anemie." From this time the adherents to transfusion have been abandoning it in favor of infusion, not that the latter can ever completely replace the former in every case, but because it can successfully replace transfusion in most cases, and where such is possible it is always given the preference, owing to its greater safety and convenience. In view of these advantages, it is proper that we employ the saline infusion in all alarming hemorrhages.

This operation is uniformly followed by an improvement in the circulation. Should, however, the improvement in the circulation last but a brief interval, it is plain that the amount of blood remaining is incapable of carrying on the condition necessary for life and then it is evident that transfusion must be performed in addition to infusion.

In addition, it has been pointed out by other authors that the saline infusion protects the internal organs from a too rapid and extensive abstraction of their parenchymatous fluids, which nearly always occurs after an extensive hemorrhage.

By saline infusion we mean the injection of a solution of sodium chlorid having the same strength as the serum of the blood (0.6 per cent.) into the veins of the bloodless subject. Some add to this a trace of sodium hydrate or sodium carbonate, while others contend that this addition is unnecessary and that should either the carbonate or hydrate exist in proportion of more than a trace, their presence would become positively injurious. Again, others dissolve the salt

in a weak saccharin solution, or the infusion has been successfully carried out by Thomas and others with pure fresh milk.

As for the proper performance of infusion, there are several features to be observed. First of all, the solution should have a temperature of about 100 degrees F. The infusion must be made with a slow, steady and not too strong a stream. Ordinarily, sufficient force is obtained by raising the funnel or the vessel containing the solution to the height of an ordinary arm's length; should this convey the solution with too much force, the latter can easily be regulated by lowering the container to the desired level. The quantity to be employed must depend upon the nature of the case. It is not necessary in every instance to employ as much solution as the amount of blood lost; all that is required is to inject just enough of the solution to restore the tone of the circulation. Ordinarily twelve to fifteen ounces can be considered as the minimum quantity to be used for ordinary purposes, but from twenty to twenty-five ounces may be used in alarming hemorrhages.

The operation itself can be divided into three stages: First, the exposure of the vein, preferably at the bend of the elbow. In the second stage there is one feature that deserves emphasis, the transfusion tip should only be introduced into the vein while the fluid is running. The observance of this precaution not only washes apart the lips of the opening in the vein, but affords an absolute safeguard against the introduction of any air, which is one of the chief dangers associated either with transfusion or infusion. The third step consists in ligating the punctured vein and applying an aseptic compress.

The principal points in connection with the operation are: To have all steps performed in an aseptic manner, to be careful that the infusion is not made too rapidly nor with too much force, otherwise there will be danger of over distending and paralyzing the already much enfeebled heart. Care should also be exercised that too much fluid is not employed which might give rise to a two-fold danger. 1. By the unfavorable influence upon the corpuscles by the dilution of the blood. 2. By raising the intravascular pressure to such a degree as to occasion a rupture of one of the smaller vessels in some vital part of the economy. It is exceedingly important that great care should be exercised to have the salt solution absolutely free from any minute floating bodies, which if present might act as emboli producing death or perhaps a gangrene of one of the extremities.

Before dismissing the subject of transfusion and infusion we desire to recall the fact that owing to the extreme cerebral anemia which exists after dangerous hemorrhages, either of these operations can and should be performed without the use of an anesthetic. In this condition all manipulations are practically of a painless character and owing to the enfeebled heart an anesthetic would only be adding an additional danger.

ANTO-TRANSFUSION.

By anto-transfusion we understand the forcing of the blood by means of elastic bandages from the extremities to the more vital centers. The indications for this can be summed up as follows: Shock or any condition attended with relaxation and diminished vascular pressure. Again, this is very useful in guarding against accidents in anemic subjects during chloroform narcosis.

SUMMARY.

In conclusion I beg to submit the following:

1. In enormous hemorrhages the resulting danger are more frequently due to the reduced intravascular pressure than to the actual loss of blood.

2. In view of this the indications point more decidedly toward infusion than transfusion.

3. That transfusion has not received the attention which its merits justify.

4. In transfusion we possess a measure which in the severest hemorrhages is the only agent capable of restoring the vital functions.

5. The indication for transfusion includes any condition which reduces the total quantity of blood to a fatal degree or which alters the character of the blood to such an extent as to render it incapable of sustaining life.

6. When the transfusion is performed for the relief of a poisoned condition of the blood it should be preceded by venesection.

7. Centripetal is to be preferred to centrifugal transfusion.

8. In centripetal transfusion the injection should be made with a slow steady stream, carefully avoiding undue force.

9. In withdrawing the blood from the donor the veins afford an easier, safer and better source than the arteries.

10. Indirect transfusion with defibrinated blood is safer than direct transfusion with non-defibrinated blood.

11. In alarming hemorrhages infusion should be performed before transfusion; should however, the improvement be transient in its nature, the infusion must be supplemented with transfusion.

12. In addition to hemorrhages the indications for infusion include any pathologic state attended with a feeble pulse which is dependent upon a relaxed condition and a diminished intravascular blood pressure, namely, shock.

13. Restoring the tone of the circulation by infusion is not wholly dependent upon the increase of the intravascular pressure, but is in part due to the stimulating influence which the salt solution has upon the heart.

14. In performing transfusion or infusion after an enormous hemorrhage, the use of an anesthetic is not only unnecessary but absolutely dangerous.

15. In the ante-transfusion we have a valuable measure for combating shock and preventing accidents in anemic subjects during chloroform narcosis.

OVARIAN TUMORS COMPLICATING PREGNANCY; WITH REPORT OF A CASE.

Read before the Chicago Medical Society, June 15, 1896.

BY C. S. BACON, M.D.

Professor of Obstetrics, Chicago Polyclinic.
CHICAGO, ILL.

Within the last ten to fifteen years the management of this interesting and important, if somewhat uncommon, complication of pregnancy has become much more uniform, with a corresponding improvement in the prognosis. While formerly the let alone policy, the induction of premature labor and the puncture of the tumor were methods quite commonly employed, now the much better results obtained from ovariectomy are generally known and recognized, and this mode of treatment has been substituted for the

others. Two questions of much interest are still undecided. One is, Which is the better route to reach the tumor, the abdomen or the vagina? The other is, What is the best time to operate? The following case is presented because it may have some bearing on these mooted questions and because the casuistic contributions are not yet too great:

Mrs. C., aged 32 years, III-para. Family history good. Nothing abnormal in menstrual history. First pregnancy, four years ago, was without complications. Labor was slow and forceps were applied, probably unnecessarily. Laceration into the rectum with severe hemorrhage resulted. A few months later I was first called in and repaired the laceration. During the examination a small tumor was noticed on one side. The differential diagnosis was not made between a small cyst of the ovary, a parovarian cyst and a pedicled subserous myoma of the uterus.

Soon after the operation the patient again became pregnant. This pregnancy was also undisturbed. I attended her confinement, which was quite normal. After labor the adnexa tumor was again noticed as a rather hard mass, to the side of the uterus, about one inch in diameter.

May 4, 1896, the patient came to me, to arrange for her next confinement. She had removed from the city to a suburb. She had no suspicion of anything abnormal and complained only of feeling unusually weak with much distension of the abdomen. Her last menstrual period occurred Nov. 1 to 4, 1895. Fetal movements were first felt March 17, 1896, i.e., in the twentieth week. She therefore would be in the beginning of the twenty-seventh week of pregnancy.

Examination disclosed the following condition: The uterus lay in the left side of the abdomen, the fundus being but a finger's breadth below the ribs. The child lay with the head presenting, freely movable above the pelvis and corresponding in size to the supposed age of the fetus. In the right side of the abdomen, extending about two inches above the navel, was a tumor which could be distinguished from the uterus by a groove which ran along between it and the uterus one to two inches to the right of the median line. Through the vagina the pelvis was found filled with a tumor apparently in connection with the tumor in the right side of the abdomen. The cervix was reached above the symphysis a little to the left of the middle line.

May 12, in the twenty-eighth week of pregnancy, I operated at the Polyclinic Hospital, with the valuable assistance of Professor Henrotin. On opening the abdomen the large veins of the pampiniform plexus, feeling like coils of intestine, were found lying in front of and to the left side of the tumor, which was not adherent. Its contents were evacuated through a trocar and an attempt made to follow the sac to its pedicle. Then it was found that the tumor arose from the left broad ligament, a fact that could have been suspected from the location of the pampiniform plexus on the left side of the tumor. With considerable difficulty the sac was pushed under the gravid uterus, its broad pedicle ligated and the sac removed. The abdominal wound was closed with deep sutures through the skin and muscular layers, and two rows of continued sutures through the peritoneum and the muscular fascia. A small accidental incision of the peritoneum covering the uterus had been closed with three sutures.

The tumor sac, which held nearly two quarts of brownish transparent fluid, was plainly of ovarian origin.

For thirty-six hours after the operation the patient had considerable pain, for which she received three or four hypodermic injections of one-fourth grain morphia. There was also considerable nausea for two days, but after free action of the bowels she became easy and made a good convalescence. Except for a temperature of 101.2 degrees the evening after the operation there was no fever. As a special precaution against premature labor she was kept in bed till the twenty-third day. She then got up and improved rapidly in strength, and on June 7 felt perfectly well and planned to leave the hospital the next day. During the night, however, labor pains came on, resulting in the delivery of a living child June 8, 9 A.M., not quite twenty-eight days after the operation. The labor was not unusually painful and the abdominal wound, which had perfectly healed, was not disturbed. The placenta was delivered shortly afterward by the Duncan mechanism. Its uterine surface contained several old hemorrhagic fibrinous patches and was quite friable. The child was 43.5 cm. long and weighed about three pounds. It was not very vigorous, and was kept in an improvised couveuse and fed every hour. In spite of this attempt to save it the child became cyanosed and died about eight hours after birth. The patient was