however, it is necessary to cleanse the general cavity, the 0.6 per cent. salt solution at blood-heat should be employed. Drainage will be necessary if a noxious material is left in the peritoneal cavity, or if it is expected that an extensive effusion will follow. Objection is made to the glass drainage-tube on account of its rigidity. The fenestrated rubber tube meets all indications, and is free from this objection; occasionally the use of strips of iodoform gauze will best secure the necessary drainage.

In conclusion, the author remarks that the surgeon's position with regard to peritonitis is, at the present time, similar to that which obtained some thirty years ago in regard to external inflammations. His efforts are directed against the effects of damage already done, rather than toward measures of prevention.

OBSTETRICS.

UNDER THE CHARGE OF

EDWARD P. DAVIS, A.M., M.D.,

PROFESSOR OF OBSTETRICS AND DISEASES OF INFANCY IN THE PHILADELPHIA POLYCLINIC;

CLINICAL PROFESSOR OF OBSTETRICS IN THE JEFFERSON MEDICAL COLLEGE; CLINICAL

PROFESSOR OF DISEASES OF CHILDREN IN THE WOMAN'S MEDICAL COLLEGE;

VISITING OBSTETRICIAN TO THE PHILADELPHIA HOSPITAL, ETC.;

ASSISTED BY

WILLIAM H. WELLS, M.D.,

ASSISTANT DEMONSTRATOR OF CLINICAL OBSTETRICS IN THE JEFFERSON MEDICAL COLLEGE;

CLINICAL ASSISTANT TO THE CHAIR OF OBSTETRICS AND DISEASES OF

INFANCY IN THE PHILADELPHIA POLYCLINIC.

TWO CÆSAREAN SECTIONS FOR CICATRICAL STENOSIS OF THE SOFT PARTS.

KAYSER (*Zeitschrift für Geburtshilfe und Gynäkologie*, Band xxvii., Heft 2) reports in detail the histories of two cases operated on at the University Frauenklinik at Giessen. The first patient was twenty-nine years old and at the end of her second pregnancy. In her first labor the child had apparently been extracted by forceps after cephalotripsy had been performed, and six weeks later the patient had been operated on for vesico-vaginal fistula by Prof. Löhlein, with favorable result. On vaginal examination it was found that the vagina was narrowed, and stretching from the cervix backward and on all sides were strong bands of cicatricial tissue. On the anterior vaginal wall the cicatrix of the old operation for vesico-vaginal fistula existed as a hard swelling. A true portio was not present, but in place of it there existed only a flat projection, in which, high up toward the promontory, the external os uteri existed as a small shallow cavity. A double promontory with some sacral lordosis were present. Induction of labor by means of hot and cold vaginal donches, tampons, etc., was tried without success, the cervix being only capable of a small amount of dilatation, owing to the existing cicatricial tissue. As the patient earnestly desired a living child, and it was feared that deep cervical incision might lead to unfavorable results for mother

and fœtus, the Cæsarean operation was decided on and performed on the fifth day after the patient's admission, the child being safely delivered. No unfavorable symptoms occurred to the mother except slight secondary hemorrhage from the vagina, which was easily controlled by an ice-bladder and ergot. Both mother and child left the hospital in good condition, though in the former the old scar tissue still remained and no portion could be felt.

The second case was a II-gravida, who gave a history of having been delivered of a dead child by forceps after a labor of three days. From this a vesico-vaginal fistula developed. The vaginal entrance, the canal itself, and in particular the inferior portion of the recto-vaginal septum were very much narrowed by cicatricial tissue. A urethro-vaginal fistula also existed. Both fistulae were closed by operation, but the vagina remained in a state of incomplete atresia. The patient's pelvic measurements were as follows: Sp., 22.3; Cr., 25.5; Tr., 29; Conj. ext., 18.5. In the hope of helping dilatation of the soft parts and of exciting uterine contractions, warm vaginal injections of carbolic acid solution and tampons were tried without success. As it seemed impossible to bring a living child through the unyielding cervix and vagina, it was decided to do Cæsarean section. Considerable hemorrhage was encountered during the operation, but was successfully controlled. The child weighed 1850 gms., and was living. The mother, after a somewhat variable puerperal period, died in coma on the eighth day, and at the autopsy the following conditions were found: Fatty degeneration of the heart with anæmia of all the organs. Spleen enlarged, and in the gall-bladder, which was considerably enlarged, were found many gall-stones. In pieces of the uterine tissue subjected to bacteriological examination there were found in the endometrium strepto- and staphylococci with other bacteria, while in the muscular tissue only small oval forms were present. A guinea-pig inoculated from the endometrium died in twenty-four to thirty-six hours, whilst another into which a solution of the muscular tissue was injected remained healthy. The child of the second patient died of icterus.

SIX CASES OF BRIGHT'S DISEASE IN PREGNANCY AND LABOR.

HERMAN, at a meeting of the Obstetrical Society of London (*Lancet*, January 13, 1894), presented the histories of six cases of Bright's disease occurring during pregnancy and labor. From the comparisons of the histories of the above and others published in the Society's *Transactions*, he concludes that there are at least two kinds of renal disease to which a pregnant woman is especially liable. One of these is a very acute disease in which premonitory symptoms are either absent or of duration measurable by hours or days. It attacks chiefly primigravida and often causes intra-uterine death of the child. The above form of the disease is also attended with extreme diminution of the quantity of uria; the small amount that is passed being greatly deficient in urea, but containing enough albumin to make it solid on boiling. This form of disease is accompanied with rapidly occurring fits; but if it runs a favorable course, the fits cease and the urine increases in amount, the percentage of urea rising therein. If the excretion of urea is not re-established the case quickly ends fatally. Such cases seldom, if ever, pass into chronic Bright's disease. The other is a disease which attacks older subjects, chiefly