erable benefit to the institution as well as to each attendant who takes interest in it. It is my object to give a sufficiently thorough course of training in the second year to be able to train our own nurses for the positions of head nurses and all superior positions.

I am certain, however, that with uniform scale of wages no such results can be expected and the institution will drift along way behind the times, simply because of lack of efficiency in the force of attendants and nurses.

V. H. PODSTAT, M.D.,
General Superintendent Cook County Institutions.

"Congresses on Tuberculosis."

NEW YORK CITY, Dec. 14, 1903.

To the Editor:—The American Congress on Tuberculosis was founded by the Medicolegal Society in 1900, and has held its annual meetings and elected its officers each year since that time at the City of New York. This congress is not, nor was it ever, for a moment, considered as a strictly medical organization. It was composed of legal and medical men, scientists, and publicists, because the medical profession, as such, in this country failed to organize such a congress, although all the leading powers of christendom had moved and acted. Sweden, Italy, Germany, France and England had done so.

The Medicolegal Society, because of the inaction of the medical profession of America, considered that if consumption was a communicable disease, as their medical members believed and advised, that the highest questions in forensic medicine ever submitted to the two great professions of law and medicine were:

(1) "What legislation could be adopted by Congress or by the legislatures of the American states that could possibly arrest its ravages, prevent its spread, and reduce the mortality among our people that had reached such alarming and amazing proportions?" as the first question, and,

(2) "How could the masses of our people be educated up to be willing to vote for and aid in securing such legislation as a commencement; and to form and place on solid foundations a public opinion in America, among all classes of our people, that could be relied on to enforce such laws, after they were enacted?"

The Medicolegal Society then considered that, because of the exigencies of the questions, the supineness and inactivity of the medical profession in our country to grapple with the subject, it became an urgent, an immediate and the paramount duty of lawyers, jurists, medical men, scientists, legislators and the general public in such a struggle to inquire into all sides and phases of this question.

If the medical profession had in 1900 organized a national movement or congress on tuberculosis, as that profession had done in almost all continental countries, it would have been a creditable work, and . . . it is only fair to say that the present congress would not have been organized.

If the medical profession should to-day, through its great national organizations, decide to call and hold a medical congress on tuberculosis, either on broad or restricted lines of labor, as to the medical questions involved, or on larger and broader lines, it would be a creditable work, and every member of this organization would aid it in every way.

As this Congress is not a medical congress, *per se*, and as men of all professions are on its list of officers and members, its faults ought not to, nor could they ever, be laid at the door of the medical profession.

(Signed) CLARK BELL,
Chairman Committee on Organization,
International Congress on Tuberculosis.

[Because of lack of space for them, we have omitted from Mr. Bell's very long letter a number of paragraphs, some of which are a repetition of a considerable part of the letter from the Government (which has already appeared in THE JOURNAL, December 12). We print the above portions, which are pertinent to the matter discussed. Mr. Bell discusses the

Lewis-Brown Congress, which discussion we omit, as we do not wish to enter into the controversy between the two congresses. We note that Mr. Bell relieves the medical profession from responsibility of any sort for the congress, and it is well for physicians to know this fact.—ED.]

The First Hip-Joint Amputation in America.

WINFIELD, ARK., Dec. 18, 1903.

To the Editor:—In THE JOURNAL, December 5, page 1379, in his address on "The Progress of American Medicine," etc., Dr. W. W. Grant of Denver, speaking of great surgical operations of the past, gives the late Dr. V. Matt the credit of the first "hip-joint amputation" in America, in 1824. This is not correct, as we find, on pages 4 and 5 of Dr. P. S. Townsend's preface to volume III of Velpeau's Operative Surgery, 1847, that he gives Dr. Walter Brasher of Bardstown, Ky., the credit of the first hip-joint amputation, which was performed in August, 1806. In 1891, this question was discussed by Dr. Will F. Arnold of the U. S. N. and myself, and our communications were published in the *Medical Record*. In one number of that journal I had a copy of Dr. Brasher's letter to Dr. Townsend published.

Dr. Wyeth of New York, in the third edition of his "Text-Book of Surgery," page 218, gives Dr. Brasher credit for the first operation of the kind in America, and cites Dr. Yandell of Kentucky as authority. Dr. Eve, in his "Surgical Cases," gives the proper author of the operation, as Townsend did.

I write this simply to place the credit on the right man, and on the right date.

CHEVES BEVILL.

Queries and Minor Notes.

ANONYMOUS COMMUNICATIONS will not be noticed. Queries for this column must be accompanied by the writer's name and address, but the request of the writer not to publish his name will be faithfully observed.

DIFFERENTIAL DIAGNOSIS OF TYPHOID FROM MEASLES AND RHEUMATISM.

RIDGELAND, WIS., Dec. 25, 1903.

To the Editor:—CASE 1.—A patient, 4½ years old, came down with a high fever and aches and pains all over her body. She had been exposed to measles. By the end of the first week her temperature was 104. At this time also there was pain in her abdomen and it was also tympanitic and covered with roseola. Her bowel movements were like pea soup and there was some blood. Spleen was enlarged. She gradually grew better until the end of the second week, when her temperature was 100. During the second week there was also bronchitis. Beginning with the third week there was a rise of temperature and the second day she broke out with measles. She then rapidly recovered. I diagnosed the case typhoid fever. Was I right or was it measles? She had two good sweats at the first.

CASE 2.—A woman, 38 years old, after three days of being unwell and having a very sore throat, came down with pain and swelling in her joints, mostly in wrists and elbows, ankles and knees; there was much vomiting and there was enlargement of spleen; temperature was from 100 to 101; there also was a marked roseola on abdomen. Tongue covered with white coating, breath bad. She was constipated, but there was no sweating. I diagnosed the case rheumatic fever. I opened the bowels with colchicin, and just at this time another physician was called, who pronounced it typhoid fever. Under typhoid treatment her temperature would run up to 101, while under rheumatism treatment her temperature would remain at 99.6. She finally recovered with stiffness in the joints. Who was right?

— C. C.

ANSWER.—CASE 1.—There are two or three possibilities in this case. (A) The patient had typhoid which was followed by measles. The early fever, abdominal symptoms, roseolæ, stools, splenic tumor, hemorrhage from the bowels, etc., speak as strongly for typhoid as any symptoms could, in the absence of a Widal test. (B) In some cases, in young children at this age, the typhoid spots, usually small, few and confined to the trunk, may come out very diffusely, abundantly and with unusual distribution, as over the face, arms, hands, etc. These cases may resemble measles, and careful observers have recorded such cases, although relatively infrequent. We have seen typhoid in which the diagnosis of measles was made. The Widal test, finding the typhoid bacilli in the blood and urine, etc., removed all doubt of the diagnosis. Your case was, then, in all probability typhoid with, as an equal choice, measles as a sequel or an unusually abundant typhoid eruption.