Original Articles.

THE HUMAN FOOT IN ART.¹

BY E. H. BRADFORD, M.D., BOSTON.

WHEN the sculptor Hiram Powers pointed with pride to the anatomical excellence (in his eyes) of his "Greek Slave" in contrast to the anomalies in the classic masterpieces, he argued himself lacking in true artistic spirit.

Correct commonplace is not preferable to what is beautiful even though the latter is defective in minor details.

It is certainly hypercriticism to lay undue stress upon small trivialities where the whole commands admiration.

No true Romanist kissing the bronze toe of St. Peter pauses to examine the modelling of the foot; and any one who looks at the anatomical defects of Fra Angelico's "Coronation of the Virgin," or Botticelli's "Madonna" at the Louvre, robs himself of time which might be more enjoyably employed.

And yet a study of great Homer nodding has its interest and value, and need not detract from our true admiration.

Any one whose attention has been called to the subject of the shape of the human foot will find it of interest to examine the feet in a collection of statues. While there is some variation in the work of different sculptors, it will be found that the types can be classed under three heads: first, Barbaric and Egyptian type; second, the Classical type; third, the neo-Classical and Modern.

The first may be considered the type of the barefooted or sandal-wearing people; the second, sandal- and cothurnus-wearing people; and the third, those of the races that have worn shoes.

The Barbaric and Egyptian type is, as a rule, badly modelled, with the exception of some of the best of Egyptian sculptures, where modelling in some instances is carefully done.² The various statues of Rameses II, B. C. 882, the famous wooden statues of the Fourth Dynasty, B. C. 3700, are all excellent types of this class. Under this head can also be placed Assyrian specimens and the works of early Indian, Burmese, Siamese, Chinese and Japanese art, as well as the few crude figures from Africa, Mexico and Alaska, and also the archaic Etruscan, Cyprian, and the early Greek statues.³ This class is characterized by the lack of divergence of the axes of the feet in standing figures, that is to say, by the parallel position of the axes of the feet, by the straight inner edge of the foot, and by the straight and uncrumpled condition of all the toes. In the Egyptian and early Greek art, where there is any attempt at careful modelling of the toes, the first toe is separated from the second, and in most instances the second toe is represented as somewhat longer than the first.

The second, or Classical, type is best represented by

the famous foot of the Hermes, supposed to have been made by Praxiteles. The characteristics are well marked. The line of the inner edge of the foot is always straight, except that there is a slight curve between the head of the metatarsal and the ankle. There is a separation between the first and second toes; the second, third, and fourth toes frequently diverge from the axis of the foot at an angle of at least thirty degrees and are parallel to each other; the fifth toe is represented as short and crumpled, and is placed back of the base of the other toe; the outer edge of the foot curves behind the fifth toe to the middle of the foot. The angle of divergence of the standing foot is slight. This type is seen in all Greek and Roman statues, with but little variation, the straight direction of the first toe being universal as well as the crumpled condition of the small toe. The curling of the small toe appears in Greek art after the sixth century B. C., and may be said to be universal since in all art influenced by Greek tradition.

The foot in modern sculpture as well as in that of the Renaissance is copied largely from the classic. The peculiar distortion of the small toe is universal; and, in addition, the fifth toe is often placed even farther back on the outer edge than in the best Greek type.

Wherever the sculptor has endeavored to leave classic models and to copy nature, he has produced a curious medley in distortions. This is particularly the case in recent sculpture, where deviation of the greater toe is extremely common. Modern sculptors, neglecting the classic tradition, copy imperfect models. The outward divergence of the second, third and fourth toes from the median line is copied from the Greek and exaggerated, representing the fourth toe as straight and uncrumpled—impossible in modern shoe-wearing models. The modern sculptor copies also the curved line on the outer side of the foot, which is seen in Greek and in Roman art, especially exaggerating the classic curve from a desire to present a curve of beauty.⁴

Certain modern sculptors place the feet diverging from each other at a greater angle than is seen in the Egyptian and Classical periods.

The Greek showed such a mastery in surface anatomy, the Barbaric type is so crude, and the Modern type so impossible, that the first and readiest impulse would be to accept at once the foot of the Hermes of Praxiteles and show it as a model for all sculptors, as is done in many art schools; but any inquiry as to which of these types is the most correct should rest on a firmer basis than a hasty conclusion.

To form an opinion in the matter it is necessary to bear in mind the deformities of the feet caused by the footwear in the different periods referred to, and to study the foot unaffected by shoes. Of most importance is the common distortion of the great toe, which is known as hallux valgus, which results from the yielding of the great toe to outward pressure at its extremity. The corresponding inward distortion of the small toe from shoes is also often seen. A

⁴ The number of modern sculptors who give to their statues hallux valgus is too great to be enumerated in a short paper. Even as careful an artistic anatomist as Gérôme, in his "Bellona" gives this defect. The sculptors in the earlier part of the century were more careful than the modern French school, but they frequently modelled the deviating great toe (Thorwaldsen, Canova). This is also true of the brilliant school at the time of Louis XIV and of the Renaissance (Donatello, Michael Angelo).

¹ Read before the American Orthopedic Association, Washington, D. C., June, 1897.

² A mummy may be seen in the Museum of the Louvre, illustrating the care with which the straightness of the toes was preserved; each toe, including the fifth, was separately wrapped. In the British Museum an unwrapped mummy of an Egyptian official of the Eleventh Dynasty showed the straight first toe seen in the statues of the Egyptian type.

³ In the Greek statues on the Sacred Way at Branchidial Caria, Asia Minor, B. C. 550, the fifth toe is uncrumpled; in all subsequent Greek statues the peculiarities mentioned of the Classical type are found. In the Museum of Northern Antiquities in Copenhagen a round beach-stone may be seen on which is cut two footprints; the inner edge is straight; the toes, including the fifth, are straight. It is of unknown antiquity.

crumpled and crowded condition of the toes known as hammer-toe, or curling downward of the toe, results from too short shoes or from the backward pressure exerted by the front of the shoe on the extremity of the smaller toes. Another affection of the toe is extremely common in persons whose feet have been compressed by shoes, namely, a weak or powerless condition of the muscles which prevents active muscular use at the phalangeal joints.

Some distortion of the toes is seen at a comparatively early age. The examination of the feet of a number of children indicates that in communities where shoes are worn the feet, even in young children, differ in shape from the normal shape of the foot; the separation of the great toe from the second toe is lost at an early age; the fifth toe is crowded towards the fourth and the muscular strength diminishes even at the age of three. Some distortion of the foot in modern cities may be said to be universal, and sculptors seeking perfect models for the feet find no model with uninjured feet.

SHAPE OF THE FOOT UNAFFECTED BY SHOES.

If the foot of a young infant is examined, it will be seen that there is muscular power in the movement of all of the toes. The great toe can voluntarily be drawn to the inner side, and the fifth toe can be drawn to the outer side by voluntary muscular exertion; the toes can be flexed readily. The second toe is, when stretched to its full length, frequently longer than the first; the third is of the same length of the first; the fourth is somewhat shorter; and the fifth, though shorter, is but slightly so, and is readily moved, flexed, extended as well as abducted, by muscular action. None of the toes remain permanently curled, though when in a relaxed condition the terminal phalanx drops somewhat and the smaller toes curl. A separation between the first and second toes is normal when the muscles are active, and the great toe is drawn to the inner side frequently. The line of the extremity of the toes presents a gradual curve, the greatest convexity being at the tip of the second toe. There is no twist inward of the fifth or small toe. The curved line connecting the proximal end of the toes is a curve similar to that formed by the extremities of the toes. The line of the inner edge of the foot is always straight, except when there is contraction of the muscles when the great toe diverges to the inside.

Adult feet that have not worn shoes present shapes similar to that of infants, except that in the ordinary wear and tear of locomotion the skin is thickened, and some curling downwards of the smaller toes is frequently seen, owing to the strength of the plantar muscles. For this reason the second toe is not usually seen longer than the first, as the position which it assumes is usually less straight, it being less strong. The inner edge of the foot is always straight; but in people who walk actively, the great toe and its metatarsal becomes developed in strength and size more than is seen in infants. The foot is longer relatively to its width; the fifth toe, being the weakest, is very susceptible to slight changes from lateral pressure, and where slippers have been worn or loose shoes, a slight twist of the terminal phalanx of the fifth toe is frequently seen. The widest part of the foot is at the cross-line drawn across the toes and including the fifth toe, as in the Egyptian foot. The foot somewhat re-

sembles that depicted in the Egyptian sculptures, except the toes are parallel, and the second, third and fourth do not diverge from the long axis of the foot, and in the position of rest the smaller toes are more flexed than in the Egyptian statues.

An investigation of the deformities of the foot due to footwear leads to an inquiry into the different kinds of foot-covering. No proof is needed to show that the modern shoe distorts the foot, and an examination of any collection of shoes (as the excellent one at the Museum of the Cluny and the remarkable historical exhibit of shoes at the Chicago World's Fair) shows that shoes of the Middle Ages distorted the feet nearly as much as the modern ones.⁵ Distorting shoes were probably not as universally worn as at present, and were perhaps used chiefly by the leisure classes. But pointed-toed shoes were fashionable even among the Egyptians. In fact, it may be said that as soon as man ceases to need the free use of the foot in hunting, bread-earning, marching or locomotion, he attempts to use the shoe as a means of ornamentation, according to his own fancy; as an evidence of distinction and a mark of freedom from base toil — like the compressed foot of the Chinese lady.

Footwear can be classified as follows: boots or shoes, where the whole of the foot is covered; slippers, where the front of the foot is covered, both holding the front by the upper, covering the dorsum of the foot; and sandals, where an inter-toe strap connected with cross-straps holds the sole.

Although boots, shoes and slippers vary in shape according to period and fashion, with the exception of the boots of laboring men and jack-boots, ornamental shoes were almost always pointed, though at the period of the Thirty Years' War, and at a time in the seventeenth century, square-toed shoes were occasionally seen for men. Medieval and Oriental slippers do not retain the heel, and allow the foot to slip backward without crowding the front of the foot forward.

Sandals furnish a protection to the sole, and are secured to the foot by some form of thong or strap. The oldest form of sandal seen in sculpture is the Assyrian,⁶ which was retained by a strap around the great toe connected with straps from the sides of the sole and behind the heel. In the Egyptian sandal a thong passing between the first and second toes, from the middle of the front of the sandal over the dorsum of the foot, was connected with thongs from the sides of the sandal. This sandal is the most common type, and resembles the modern Japanese and the modern Mexican and South American sandal. The ancient Toltec sandal was furnished with two thongs, one passing between the first and second toe, and the other between the third and fourth. The modern Somalis wear a sandal with a thong split so as to admit the second toe, leaving the other toes free. The early Etruscans wore a sandal with cross-straps, the front one passing from the sole of the sandal, between the first and second toes, over the second, third and fourth toes, where it is secured to the sandal, leaving the first and fifth toes free, the other toes being secured under a strap.⁷

In all these forms of sandals, no pressure is exerted upon the small toe; but in a form of sandal with a

⁵ This is also seen in the sculptured tomb monuments of the Middle Ages.

⁶ Collection at the Peabody Museum, Harvard University.

⁷ This is seen in the rare terra-cotta tomb effigy of the early Etruscan period (uninfluenced by Greek art) in the British Museum — showing a sandal differing entirely from any seen elsewhere.

FIG. 7.

FIG. 8.

FIG. 9.

THE HUMAN FOOT IN ART.

THE HUMAN FOOT IN ART.

front, a cross-strap seen in the Orient for the insertion of the front of the whole foot, a certain amount of pressure is exerted on the side of the little toe, as in a slipper. This strap, however, is as loose as the cross-strap in snow-shoes. In the Greek sandal, however, a cross-strap or straps were connected by lacings to be tightened in such a way as to be more firmly held to the foot. The forms varied greatly. In most a single inter-toe strap existed to be caught in the cross-straps. In later sandals, however, the inter-toe strap was omitted, a reminder surviving conventionalized in the shape of the sole, which preserved a slight depression at the front of the sole between the first and second toes. The Greek and Roman sandal gave freedom to all the toes except the little toe, which was crowded by the tight cross-strap, as well as by oblique straps, which in some forms of sandals passed backwards from the sides of the front of the sandal to the instep, raising and crowding the little toe and the side of the first metatarsal. When this was snugly laced, an inter-toe thong became unnecessary, as the pressure on the little toe and the top of the foot prevented the foot from slipping forward.

This form of footwear can be easily imagined, if we suppose a low bicycle shoe, with the tip removed, leaving all the toes except the small toe free. A sandal of this sort was perhaps not developed in the early period of Greek history; but, judging from sculpture, the cothurnus of the late Greek and Roman period differed but slightly from that of the Classical period.

The ancients wore at times beside the sandal a loose covering like a moccasin, which did not affect the shape of the foot. Footwear of this sort would develop the shape in the foot seen in the Hermes of Praxiteles, crowding the small toe, giving the full length of the fourth, third and second toes (from the free and strong development of these toes), allowing a straight first toe and the slight inward curve on the inner edge of the first metatarsal at its proximal end, from the pressure of the oblique strap, and also the outward curve on the outer edge of the foot, as the chief pressure of the oblique strap came upon the little toe and not on the metatarsal. The inter-toe strap accentuated the separation of the first and second toes and the deviation from the axis of the foot of the smaller toes. This slightly distorted foot was seen in the models studied by the Greek masters.⁸

The dominating influence of Greek art explains the fact that the shape of the foot has been copied since

then by all true sculptors who have accepted Greek art as its fountain source.

In the Renaissance, where models were used by the great masters in painting and sculpture, the deformity of the great toe following the use of the shoes which crowded the first toe outward is commonly depicted, and in sculpture is occasionally seen. In modern French sculpture, among the realists this defect, the well-known hallux valgus of the model, is added to the classic defect, namely, an exaggerated difference between the fourth and fifth toes, and exaggerated rounding of the outer edge of the foot. Modern sculptors frequently also give a greater angle divergence to the axes of the feet in the standing figure than is seen in Classical statues, and much greater than in Egyptian or Barbaric figures. In this, too, the modern artists are copying defective models to the detriment of their art.

It can, with a certain amount of justice, be argued that, when a conventional use has been sanctioned for a long period of acceptance without violating the laws of beauty, it has justified its existence by a right of pre-emption, and that it is hypercritical to object to it. The argument hardly applies to a deformity, and would at once seem inadmissible, if urged by a Chinese artist in defence of the beauty of the "golden lilies," as Chinese poetic fancy terms the deformed Chinese ladies' feet. The deformities here mentioned differ in degree, but not in kind, from that accepted by usage in China; they are also distortions of nature.

There is a healthy tendency in modern art to revolt against what seems classic convention, and to copy nature; but, to do this, the artist needs thorough knowledge to enable him to escape presenting as a standard the deformity observed in his own experience.

The taint of the corseted and high-heeled, weak-ankled grisette injures every modern art exhibition, even when the ideal is attempted. The sculptor who moulds "immortal forms" should not perpetuate deformity. There should be no Zolaism in sculpture.

Few, even among artists, realize how common it is to find defects immortalized in masterpieces. The wry neck of Alexander the Great may be allowed in a portrait statue; but it would be an error to give to the statues of all heroic generals twisted heads. Crumple-footed divinities, out-toed demigods, demi-goddesses and weak-ankled national heroes cannot but offend the fastidious, when seen in lasting marble and monumental bronze.⁹

The normal foot resembles a combination of the Greek and Egyptian sculptured foot. It is straight on the inner edge, without the slight curve along the base of the first metatarsal; the fifth toe is parallel to the other, is not crumpled or twisted, and is not placed far back of the line of the other; the outer edge is but slightly curved, and does not bulge between the fifth toe and the ankle. The toes are flexible, and when at rest do not project stiffly, as in the Egyptian statues, but are slightly flexed, especially the smaller toes.

⁸ The defects in drawing the human foot seen in pictures are not alluded to here as fully as could be done in a longer paper. Less care is given to modelling in pictures than in sculpture. In brief, it may be mentioned that the defects here mentioned are more common in the paintings, even of the old masters, than in sculpture. In the gallery at Amsterdam a picture by Goltzius, A. D. 1600, depicts a dying Adonis with the feet sole forewards in the foreground and displaying double hallux valgus of an aggravated type. Rubens has never thrust his errors so directly upon the spectator; but his faults in this particular are fully as grievous, as is shown in his book of design in the Musée Plantin in Antwerp and in almost all his paintings. Michael Angelo, Titian, Tintoretto, Raphael—all show both the crowded fifth toe, and frequently hallux valgus.

⁸ An examination of Greek and Roman footwear, as seen in sculpture, is made difficult from the fact that in many instances the sandal is conventionalized. A great variety is also found, from the moccasin-like foot-covering shown in the fragments from the Parthenon to the elaborate sandals of the Roman emperors. A sandal is to be seen in the late Greek statue of Ariadne, found in Roma Vecchia and now in the British Museum, showing side loops for a cross thong over the base of the toes and crowding the fifth toe, leaving the thong to the imagination. The sandal of the statue of Jason, at the Louvre, shows the indentation in the sole opposite the space between the first and second toes, where a thong would be attached, but not only is there no inter-toe thong, but the sandal is apparently to be kept in place without one by the side leathers over the instep. A peculiar sandal is seen in the group of the Roman period, in the Louvre, called "Mars and Venus," in which a depression is delineated, allowing the longer toes to drop to a lower level. The Roman sandals in the Museum of the Cluny (found near the Rhine) and at the Museum in Munich show a straight inner edge which would have satisfied a North American Indian; and the sandals depicted in the Greek votive tablet in the British Museum would have been acceptable to a modern Japanese. The bronze statuette of the youthful Bacchus, in the Payne King collection in the British Museum, shows the side compression of the inner edge of the foot and of the little toe without a cross toe-strap, while the statue of Hadrian, also in the British Museum, the Diana Chasseresse and the Seated Philosopher of the Louvre, as well as many other statues, show the compression by lacing including the cross toe-strap.

There is not much vitality in the contention that this foot is not as graceful as that presented by the Greek artist, in the mind of any one who has seen normal feet uncrumpled by shoes or distorted by toil; but even if there were more, although it is the province of the artist to furnish us with beauty as he sees it, he must find it within the lines of truth, and it is as true now as in Shakespeare's time that

"Above the art we add to Nature
Is the art which Nature gives."

DESCRIPTION OF CUTS.

FIG. 1. Diagram representing the Greek (a), Egyptian (b), and modern type (c) of sculptured foot.

FIG. 2. (a) Foot changed by modern shoe. (b) Infant's foot, toes extended. (c) Infant's foot, toes at rest. (d) Child's foot, showing beginning changes from shoes.

FIG. 3. Types of sandals: (a) Mediterranean single-thong sandal. (b) Chaldean sandal secured to first toe. (c) Somali sandal secured to second toe. (d) Toltec sandal secured to second and third toes. (e) Early Etruscan sandal secured to second, third and fourth toes.

FIG. 4. Greek sandal with inter-toe thong, causing divergence of toes; cross straps crowding the little toe; and an oblique strap pressing on the inner edge of the first metatarsal and binding the foot at the mid-tarsal joint.

FIG. 5. Drawn from the fragment of the statue of Apollo from Phigaleia (in the British Museum), showing the compression of the small toe.

FIG. 6. Drawn from the Hermes of Praxiteles found at Olympia.

FIG. 7. Egyptian seated statue (British Museum).

FIG. 8. Japanese feet, showing the lack of crowding of the fifth toe seen in Greek and modern art.

FIG. 9. From a picture of Sir Joshua Reynolds showing the hallux valgus distortion of the great toe in addition to the crowded little toe.

HEMORRHAGE ATTENDING THE EXTRACTION OF CATARACT.¹

BY O. F. WADSWORTH, M.D., BOSTON.

THE exact cause of hemorrhage attending the operation for extraction of cataract has remained an unsolved riddle. But our very ignorance as to the cause makes such cases interesting. The condition of the eyes, and the age and health of the individuals who have been the subjects of this accident, have presented very decided differences; and probably the cause is not always the same. The cases observed in my own practice do not throw much light on the nature of the affection, yet they may serve with others as material to lead eventually to a satisfactory solution.

CASE I. A man eighty years of age had been under observation for two or three years with slowly maturing cataract. During this time he had been repeatedly examined, and, aside from the lenticular opacity, no defect found in the eyes. After an interval of some months he came again to the hospital; and a somewhat hasty examination showing the cataract, in the left eye at least, mature, he was admitted and etherized. When the speculum was introduced, it was noticed that the perforating veins of that eye were larger than usual, but the tension was not specially abnormal, although perhaps somewhat increased. Puncture and counter-puncture were in the scleral edge, the summit of the flap in the cornea. Almost immediately after the section was finished a small, threadlike stream of blood was seen to flow upward across the lens from behind the lower border of the pupil, followed quickly by larger quantities which caused extensive prolapse of the iris and gaping of the wound. The eye was bandaged, and it subsequently atrophied. A few months later, when the patient desired operation on the other eye, it was found to have lost perception of light.

CASE II. A sailor, seventy-five years of age.

Cataract in both eyes. Right eye: sight failing one or two years; counts fingers at 17'. Left eye: sight failing four or five years; cataract quite dense; projection faulty; a number of superficial and several deep veins running backward from near the corneal edge, and along its upper half forming an anastomosis about $\frac{1}{2}$ " from the limbus; anterior chamber of good depth; tension normal.

October 8, 1877. Extraction, left. Puncture and counter-puncture in sclera, the summit of the flap in clear cornea. Large iridectomy before division of capsule, a small amount of cortical following spontaneously the withdrawal of the cystotome. Brief and unsuccessful attempt to remove a portion of capsule by forceps. The speculum was raised so as not to press on the globe and a moderate sized nucleus and some cortical was readily pressed out, but a considerable amount of cortical remained and several attempts to remove this by pressure through the lower lid were only partially successful. At a moment when no pressure was being made blood welled up from behind the iris, filled the anterior chamber and flowed out of the wound. After a few minutes the anterior chamber became partially free from blood and a bandage was applied.

The following day the anterior chamber was again filled with blood and a clot distended the wound. The clot gradually protruded until the corneal flap was folded backward by it, and the whole cornea became densely infiltrated. The man was discharged at his own request on October 27th.

CASE III. A man of forty-five years, always well and strong. Right eye, normal and with good vision. Left eye, mature cataract. He said this eye had been struck by a fork when he was a child, but he experienced little inconvenience at the time and the sight had been failing only a year. There was no scar to be perceived. Tension was normal; pupil active; he counted fingers at one to two feet and projection was good.

February 3, 1893. Simple extraction. The speculum removed as soon as the corneal section was completed. The lens was removed without difficulty, the pupil immediately became round and even. But within a few seconds, while the eye was still open, there was a sudden and great prolapse of the iris attended by severe pain. The lids were closed. Shortly there was a large gush of fluid vitreous, followed soon by a smaller gush and this by a stream of blood which trickled down the cheek. The man was told of the condition of affairs, enucleation was advised and done. There was no excessive bleeding from the enucleation and the healing was normal.

Although in this case the man's history so far as known was good, it is a significant circumstance that he is reported to have died suddenly of apoplexy within a year.

CASE IV. A man of sixty-nine. Good general health; mature cataract of two and a half years' duration in the left eye. Size and dilatability of pupil good; projection normal. Eight years before, a cataract had been removed from the right eye by an experienced oculist, followed by sloughing of the cornea and panophthalmitis.

April 8, 1895. Simple extraction under cocaine. Section smooth, wholly in the cornea. Ecchymosis of conjunctiva where grasped by fixation forceps. Speculum removed after capsulotomy. The nucleus

¹ Read before the American Ophthalmological Society.