ping this tissue? In a few lacerations of the perineum I have separated the levator ani muscle and have obtained the most excellent results. The only difficulty has been in getting enough tissue to overlap. No matter how much we draw them together, the ends will come apart, and I have felt that if we could overlap the muscles, we would get a better union. Dr. Harris' paper marks a distinct advance in gynecologic work, and I am particularly pleased that this advance comes through the anatomic and surgical department.

DR. EMIL RIES—The gynecologists have not entirely overlooked this subject of the muscles. Freund, in Strasburg, taught us, eight or nine years ago, when I was his obstetric assistant, to look for the levator ani muscle in perineal laceration. I did not hear the Doctor say how he makes his diagnosis of the relaxation of the muscle and the interposition of the scar tissue. The operations which are usually performed for lacerated perineum have not been confined to the mucous membrane as much as might appear from the discussion. Hegar's operation is very successfully used for the purpose of uniting the ends of the levator ani muscle. If, in this operation, we dissect out a flap of vaginal mucosa, with the rectocele behind it, and cut away this redundant vaginal tissue, the rule is to put the stitches in very deep so that they are entirely buried. I believe that gynecologists of to-day are not content with simply passing a few stitches through the mucous membrane, but they use several layers of suture. I use at least three, obviating the objection of Dr. Carey of having only a little loop, containing but a bit of tissue. That is a good observation and one which ought to show us the necessity for taking deep sutures. We make the buried sutures and superimpose three or four layers of sutures, thus bringing together the sides of the wound, no matter how deep it is, and also the tissues by the side of the rectum. We do not fail to unite the muscle in front of the rectum. We do not shorten the muscle by excising a piece, but by bringing these two sides together, and instead of having two overstretched muscles, they are brought together in this way with the effect of shortening them.

In Freund's operation, which is not much different from Emmett's or Martin's, we do not go down into the deep layers so well, and as Dr. Harris says, we do not dissect out the muscles, which is a step in advance. Dr. Goldspohn was the first to lay stress on the union of muscles. Freund describes the lacerations of the levator ani muscle, and in some of the pictures of fresh lacerations of the pelvic floor he draws special attention to these cases, but I believe he has never dissected out the muscles himself.

DR. A. GOLDSPOHN—I would differ in regard to the Hegar operation. I have traced the levator ani muscle with my finger, several hundred times at least, and find that the Hegar denudation does not extend to it. If it is to be done with a flap, the flap must be wider in the vagina than at the vulva. We would have to dissect up on the sides where Dr. Harris does his work. Neither Hegar's nor any of these methods lays bare the territory where we find the receded levator ani and the fascia.

DR. RIES—When you have the levator ani here, and you draw these two portions together, does the levator ani come with it, or does it remain outside with an empty space between?

DR. GOLDSPOHN—You bring the two portions nearer to each other, but not together.

DR. RIES—My idea was to unite these lateral parts without grasping the levator ani. You can approximate the muscle because the tissue contiguous to it must come together.

DR. GOLDSPOHN—You thereby bring them nearer to each other, but not in actual contact, as is necessary for actual union. If we want actual union of these bearing structures beneath the vagina, the entire thickness of the vaginal wall must be lifted off and the muscles and fascia, like joists beneath a floor dealt with individually and directly.

DR. T. J. WATKINS—The Society is under obligation to Dr. Harris for his valuable and scientific contribution. In his paper he neglected to mention a class of cases of injury of the pelvic floor where a portion of the levator ani muscle is divided at or very near its insertion into the radius of the pelvis. These cases can not be treated by resection of the torn muscle;

they can only be treated satisfactorily by suturing the severed portion of the muscle to the fascia of the anterior vaginal wall in the sulcus of the vagina. In resection of the torn muscle, it would seem, in many cases, that the muscle would not have sufficient consistency to allow the ends to be held together by sutures. Emmett, a long time ago, called attention to the injury of levator ani in relaxation of the posterior vaginal wall. He devised this operation because he learned that the principal injury in cases of relaxation was invariably in the lateral wall of the vagina. The Emmett operation must restore the injured muscle because it will properly relieve symptomatically at least, any case of relaxation of the pelvic outlet if it is properly done.

DR. M. L. HARRIS—Dr. Goldspohn thinks we may not always be able to find the muscles. That possibly may be true, rarely, though thus far I have met with no such case, but have been able to find them without much difficulty by going deep enough into the pelvis through the lateral wall of the vagina. Dr. Martin does not think the fasciæ could furnish the support without the muscles, nor could the muscles without the fasciæ. As the latter are but the sheaths, and purely inactive parts of the muscles and as we never have a muscle without a sheath, Nature would have been very remiss had she attempted the anomaly of placing muscles here without sheaths or fasciæ.

The drawings were not introduced as works of art, in competition with Dr. Kelly's, but to illustrate points which are nowhere, to my knowledge, touched on by Dr. Kelly. My reason for stating that I thought the majority of muscle lacerations were subvaginal was the fact that I had found the lacerations most frequent opposite the lateral wall of the vagina. The vagina usually tears along the sulcus, but we may have lacerations of the muscles without the vagina being torn at all.

In reply to Dr. Bevan's query as to the advisability of overlapping the muscle, Hayem's experiments on dividing and reuniting muscles in the dog show that the function of the muscle is best restored when there is an intimate end-to-end union.

Dr. Ries called attention to the Hegar operation as uniting the levator ani muscle. With all due respect to Dr. Ries, I must beg to differ from him. The Hegar and Freund attempt to bring the levator ani in front of the rectum is an erroneous one. The levator ani does not pass between the rectum and vagina, and any attempt to bring it there is wrong, as it does not draw the real support of the pelvic floor forward where it should be. The real support lies behind the rectum.

In regard to a separation of the muscle from the symphysis, mentioned by Dr. Watkins, such a separation has been described. Were I to meet with such a case I would try to advance the end of the muscle attaching it to the urogenital fascia, if necessary, or shorten it up by resection, if the anterior end had taken a new firm attachment.

As to the method of diagnosing lacerations of the puborectalis, I would refer to the points brought out in the paper, such as loss of the perineal flexure of the vagina, recession of vagina and anus from the pubic arch, diminution of the depth of the perineal furrow, relaxed muscle as felt by the finger, diminished contractile power, etc.

EXTIRPATION OF THE UTERUS, ITS APPENDAGES, TUMORS, ETC., WITHOUT LIGATURES OR RETENTION CLAMPS.

DEMONSTRATIONS OF NEW INSTRUMENT FOR HEMOSTASIS WITH REPORT OF CLINICAL CASES.*

BY HENRY PARKER NEWMAN, A.M., M.D.

Professor of Gynecology, Chicago Policlinic and the College of Physicians and Surgeons (Medical School of University of Illinois).
CHICAGO.

The prime object of the angiotribe here represented is to produce absolute hemostasis and do away with ligatures or foreign bodies in the operative field. The medical profession has been struggling with this problem for a long time. It is apparent to all that the ligature is not ideal or even desirable, constituting as it does a

*Presented to the Section on Obstetrics and Diseases of Women, at the Fiftieth Annual Meeting of the American Medical Association, held at Columbus, Ohio, June 6-9, 1899.

foreign body always in the nature of a complication in surgical wounds. The *écraseur* was one of the early manifestations of an epoch to do away with the time-honored ligature. In a modified form it seems to possess some merit. The clamp and cautery has held its place in rectal surgery for a considerable period and, in spite of some advantages claimed for the ligature in hemorrhoidal operations it is probable that the cautery is still used as often as the ligature method. Notwithstanding that it is now giving way to better devices, the retention clamp, likewise, has been a life-saving substitute for the ligature, and has been a means of revolutionizing pelvic surgery. The many devices for perfecting the absorbable suture and ligature, both in material used and manner of preparing, only go to show that the profession is fully alive to the importance of doing away with the conventional ligature and suture. The Keith-Skene electric heat and the Doyen

particular reference is made to the Keith-Skene and Doyen methods—apparently leave little room for argument: 1. Immediate and absolute hemostasis. 2. Perfect asepsis. 3. An undisturbed stump, for example, no bunching of pedicle or dragging upon neighboring structures. 4. The operative field free from foreign bodies, and 5, with the improved instruments recently constructed by the author, promptness of execution and adaptability to nearly all tissues and surgical work.

The general outlines of the instrument can be seen by the accompanying cuts. (Figs. 1, 2, 3.) The length of each blade is 34 cm., of which 18 are for the handles, 6 for the lock, and 10 for the blades or jaws of the instrument. The jaws taper slightly from 1.25 cm. in width and thickness to 1 cm. in width and .5 cm. in thickness. The bite of the jaws is serrated with a deep

Fig. 1.—Showing the instrument with the original cross-bar and screw attachment in place.

crusher are among the later methods that bid fair to solve the long sought problem.

It must not be understood that the leaving of foreign bodies in the wound is the only objection to the ligature, while it must be conceded that it is an all-important argument against its use. Other vital objections still remain: 1. Uncertainty as to slipping. 2. Uncertainty as to asepsis. 3. Shrinkage in the stump, thereby loosening the constricted vessels. 4. Bunching together tissue, and thereby dragging upon and displacing neighboring structures. 5. Leaving foreign bodies in the wound.

The author recognizes that in the individual ligation of vessels some of the above objections would not apply; but this very desirable method of using the ligature is not always possible or at times advisable. Again, the overtowering advantages of its substitutes—here par-

Fig. 2.—Showing the instrument with adjustable lever attachment.

groove running through its center, .25 cm. in width; the lock is 6 cm. in length, 4 in breadth, and 1.75 in thickness, the handles are 1.6 cm. in width and the same in thickness; at their ends are slots for receiving the cross-bar and screw, or the lever appliance. A small pin placed between the handles near the end serves to prevent too great pressure on the handles and the consequent danger of breaking.

The author's modification differs from the Thumin instrument in having longer blades, enabling one to include the entire broad ligament or large pedicle at a single application. The blades are narrower, so that less space is required in their adaptation to the part, and they can grasp shorter pedicles, or even sessile growths with greater facility. At the end of one of the blades is a slightly projecting bite preventing the grasped tissue

from extending beyond the blades, as it is apt to do in closing the Thumin instrument.

The lock is separable (Fig. 3) not only rendering it a more aseptic instrument, but allowing the individual introduction of its blades when desired. A new or modified Thumin lever has been added to the instrument as a special attachment. A combination mortise lock allows of an interchange at will of the lever in applying the compressive force to that of the cross-bar and screw.

While the cross-bar and screw will be found most desirable for general use, the lever has great power of compression and can be applied with a single motion or movement of the hand. This new or modified lever (Fig. 2) not only has the interchangeable lock in com-

Fig. 3.—Showing the instrument with handles taken apart and its attachments separated.

mon with that of the cross-bar and screw referred to above, but is simple in construction and adaption, and allows of a much wider separation of the blades during their application. The Thumin lever requires a partial closure of the blades and, necessarily their previous application to the pedicle or tissue to be clamped before adjusting the lock. This obstacle is overcome by the modification, as will be seen by the accompanying cuts. (Figs. 1 and 2.) The instrument, unlike the Bissell or modified Tuffier instrument used at the Woman's Hospital, New York, only requires an application of a minute or two, while the former requires three to five minutes. The instrument is applied to the broad ligaments or any soft tissue to be excised in the same manner as the ordinary broad ligament clamp, using the screw and

cross-bar or the lever as the compressive force. Immediately on its application the broad ligament is cut away on the uterine side and close to the clamp, and at the end of a minute it is removed by releasing the screw or lever. It is then applied on the opposite side in precisely the same manner. The uterus is now cut away, the toilet of the peritoneum accomplished and the operation completed. With two instruments at hand, both broad ligaments may be clamped and the uterus at once removed. In amputation of the uterus at its neck, the abdominal operation is accomplished in the usual manner, using the clamp in place of the ligature for the broad ligaments on either side. The flaps are made from behind and in front of the uterus in the usual manner, and subsequently united over the stump with a running catgut suture. I propose substituting a clamp for this part of the operation as well, preferring the thin ribbon-like stump in cementing the flaps together to the union by running ligature. A flat right-angled clamp, as has been suggested by Thumin, for the vaginal arteries, will, I believe, accomplish this. In ovariectomy or salpingo-oophorectomy the operation is very simple, clamping and removing; where there is no pus I use no drainage; where there is a septic condition, I have employed either the cautery on the raw amputated edge of the stump—which is insignificant—or strong carbolic acid. The translucent edge of the compressed tissue is at first quite blanched, but in time—as seen upon dogs—the hyperemic edge at the attached portion gradually extends its reddened area more and more, so that the entire stump is ultimately resuscitated, and little or no sloughing occurs, as obtains in a large, compressed and bunched tissue when the conventional ligature is used.

There being no granulating surfaces left, the adhesions so much to be dreaded by the ordinary method do not occur. There being no ligatures left behind, one is relieved of the apprehension of any complications from this source.

CASE 1.—Mrs. D., aged 62, German, housewife; history of previous illness negative; family history good, except that mother died of tuberculosis. She menstruated first at 15; regular until the age of 53, when menses ceased abruptly; no history of irregularity, excessive flow or pain; married at the age of 20; husband living; nine children, whose ages range from 40 to 20. Seven years ago she began to have pain in the back, at intervals; no other symptoms until one year ago, in January, when she began to have discharge and hemorrhage; hemorrhage irregular, at times a show of blood, then absent, then a flow for a considerable time—two, three, four or five weeks. During the past year the patient has become anemic, and has lost not a little flesh and strength. Examination revealed a tumor—enlarged uterus—in the pelvis, extending two and a half inches above the pubic bone, hard and nodular.

Diagnosis—Carcinoma of the uterus, and probable longstanding multiple fibroids.

Operation, Jan. 20, 1899: hysterectomy; the vaginal route was chosen, although the vagina was somewhat small and the uterus thrice its normal size. The outlet was enlarged by bilateral incisions. Incisions were made anterior and posterior to the cervix, into the peritoneal cavity, and the left uterine artery severed and tied. An effort was made to pull the uterus down, but this was found impossible on account of the size, friability and adhesions. The abdomen was opened and the uterus freed; the clamp or crusher placed on the broad ligament from above downward, compressed, and left for two and a half minutes.

The ligament was cut on the uterine side of the clamp; clamp loosened; no oozing; clamp placed on the opposite ligament, and this similarly dealt with. The uterus, now free, was lifted out and the abdomen closed; gauze drainage through the vagina; recovery complete; no after hemorrhage; only slight serosanguineous oozing from the vagina; patient up in four weeks, and left the hospital in six weeks.

CASE 2.—Mrs. E., age 20 years; American; family history

good; menstruated first at the age of 13; never regular; backache and pain over the right ovarian region, which is increased at the menstrual period; married one and a half years; menses have been more painful and backache and pain in the right ovarian region more severe since marriage.

Examination showed the uterus to be infantile, cervix small, elongated and eroded; ovaries large and prolapsed, and this enlargement and prolapse more marked in the right side.

Operation, Jan. 21, 1899: curettement; tracheloplasty; celiotomy; right ovary found low in the cul-de-sac, twice the normal size, with cystic degeneration, tube closed, enlarged and thickened; adhesions broken up and ovary and tube removed by means of the crusher or clamp. Left ovary slightly enlarged and cystic; tube closed, and many small cysts around the extremity. A section of cystic ovary was removed, the tube opened, and ampulla everted and held by fine catgut; cysts tied off with catgut; recovery good; no evidences of hemorrhage from ovarian stump. The patient was getting up in two weeks and left the hospital at the end of the third week.

CASE 3.—Mrs. F., aged 24, English. Previous history: patient well, except a monthly hemorrhage from the nose, each time lasting about twenty-four hours. This phenomenon first appeared when the patient was 6 years of age, and lasted until she was 12; mother died at childbirth. Her menses first appeared in the thirteenth year of the patient's life, and continued regularly and without pain until the first pregnancy; since birth of first child they have been painful; married at the age of 15 years; one child, 8 years old; twins, born six and a half years ago, lived six weeks; husband died in 1894; patient had a laceration of the cervix and probable infection with the first child. She has not been well since. Her symptoms increased after giving birth to twins; had double salpingo-oophorectomy and trachelorrhaphy four years ago; since then pains in the abdomen have increased rather than diminished; bowels have been constipated and bladder irritable.

Examination showed the uterus to be large and tender, and especially tender at the horns of the uterus in the old stumps, consequently a diagnosis of metritis and interstitial salpingitis of the remaining portions of the tube was made.

Operation, Jan. 26, 1899: vaginal hysterectomy. The usual incision was made around the cervix, and the uterus freed anteriorly and posteriorly. The fundus was then turned forward and pulled downward through the anterior incision. The clamp or crusher was placed on the broad ligament from above downward and left the usual length of time; then the opposite ligament was treated in the same manner, which completed the operation; recovery; relief of pain has been very marked. No hemorrhage followed operation. Suppuration was found in both horns of the uterus. The patient was sitting up in ten days, and left the hospital at the end of three weeks.

CASE 4.—Mrs. S., aged 29, German-American. History: Cloakfitter for eight years; on her feet many hours each day; large and well developed; family history good; first menstruated at the age of 14 years, and was regular until about two years ago. She suffered considerably during the first three years of menstrual life, then the symptoms became less severe until about seven years ago, since which time they have been growing worse. She has always had severe pain over the lower half of the abdomen during the menstrual period, but during the last year this pain has been constant. She also complained of severe pain in the chest over the cardiac region, without apparent local cause.

Examination revealed stenosis of the os uteri, a large uterus, movable mass to the right and above the uterus. The mass on the right side of the uterus had undoubtedly been taken from a floating kidney, as that diagnosis had been made by Chicago physicians.

Operation: curettement; tracheloplasty; celiotomy; left ovary found to be the size of a large hen's egg, cystic, and the capsule much thickened, as seen in a specimen; tube and ovarian ligament were much thickened and elongated, allowing of considerable movement; ovary removed by means of crusher, leaving tube. The right ovary was found to be the size of a hen's egg, cystic and degenerated, but to a less degree than the left; cyst of broad ligament the size of a turkey's egg, situated above the ovary in such a way that the two gave the characteristic kidney shape; cyst removed and also section of this ovary; recovery complete; no evidence of hemorrhage. The patient was sitting up in three weeks, and left the hospital at the end of four weeks.

CASE 5.—Mrs. H., aged 30, American. Previous history: Well until a year ago last Thanksgiving; family history good; menstruated first at the age of 15; menses regular and

painless until one year ago last November; married at the age of 21, only child 8 years of age; well after birth of child. Friday morning, after Thanksgiving of 1897, she began to have headache and pains in the back. Friday night she began to vomit. Saturday morning she had pain in the right inguinal region. The pain and vomiting ceased in a few days, but she did not leave her bed until the following March; pain and vomiting with each menstrual period since November, 1897.

Examination revealed laceration of perineum, laceration of cervix, large prolapsed ovary on the left side; prolapsed ovary on the right side; hard body the size of a hickorynut in the region of the appendix.

Operation, Jan. 26, 1899; curettement; tracheloplasty; perineorrhaphy; celiotomy, appendectomy—the appendix having been doubled on itself and adherent in such a way as to form the hard body in this region; section of right ovary removed, and on the left side salpingo-oophorectomy done with the crusher or clamp; no evidence of hemorrhage; patient made a complete recovery; was sitting up on the eighteenth, and left the hospital on the twenty-seventh day after operation.

CASE 6.—Mrs. J., aged 53; Norwegian; housewife; previous history negative; family history good; menstruated first at the age of 14 years, always regular until the last year of menstrual life; ceased to menstruate at the age of 48, five years ago; married at 21; has had seven children. The abdomen began to enlarge one year ago; patient had a feeling of pressure in the left ovarian region; bowels constipated. Four or five months ago she began to notice a tumor on the left side. Examination revealed a cystic tumor of the size of the uterus at seven months of pregnancy.

Operation, Feb. 13, 1899: removal of the tumor by means of the clamp or crusher for the pedicle; abdomen closed without drainage; recovery uninterrupted; no evidence of hemorrhage; sitting up on the eighteenth, and left the hospital on the twenty-seventh day after operation.

CASE 7.—Miss T., aged 47, Swedish; dressmaker; family history good; previous personal history negative; menstruated first at the age of 15; has been regular until last year, when it became excessive and more frequent; patient is anemic and has lost flesh; has noticed enlargement of the lower abdomen, of irregular shape, approaching near the umbilicus.

Examination showed the depth of the uterine canal to be six inches; irregular nodular tumor extending laterally into both broad ligaments and up into the umbilicus; movable, and to all appearance no adhesions or involvement of the adnexa; patient has heart murmur—regurgitant, mitral.

Operation, March 13, 1899: hysterectomy; median abdominal incision from one inch above pubis to umbilicus; tumor lifted out of pelvis; clamp applied to the entire length of left broad ligament. The uterus was now cut away from this side, and on removal of the clamp, was in two minutes aptwains; had double salpingo-oophorectomy and trachelorrhaphy to the opposite side; uterus cut away with the usual anterior and posterior peritoneal flaps. At one point there was slight bleeding, owing to imperfect closure of the instrument. A catgut ligature was applied, with no further trouble. The peritoneal wound was then rapidly whipped together by continuous catgut sutures, leaving no exposures of the raw surface, or the slightest oozing of blood. The abdomen was closed without drainage. The patient's condition had been precarious from the beginning of the anesthesia, owing to the defective heart. Notwithstanding the use of restoratives, saline, strychnia, etc., she did not rally, and died on the second morning, from shock and the effects of the heart lesion.

CASE 8.—Mrs. S., aged 48, American; housewife; menstruation began at 14 years of age; regular until the birth of her last child, fourteen years ago; since then very irregular, varying from two to six weeks; four children, youngest 18 years old; last labor severe and instrumental, patient suffering considerable trauma.

Examination revealed lacerated perineum, lacerated cervix, endometritis, retroflexion with extensive adhesions.

Operation, March 23, 1899, by Dr. C. W. Barrett, house physician Marion Sims Sanitarium: curettement, tracheloplasty, perineorrhaphy, removal of hemorrhoids, celiotomy breaking up of adhesions and double salpingo-oophorectomy; abdominal fixation of uterus; abdominal wound closed without drainage; no symptoms of hemorrhage; complete and prompt recovery; was sitting up the twenty-fourth and left the hospital on the twenty-eighth day after the operation.

CASE 9.—Mrs. W., 36 years old, German; housewife; family history good; has had three children, the youngest 8 years old; irregular menstruation, menorrhagia or metror-

rhagia, with laterally a bad-smelling leucorrheal discharge; backache and constant pain over the left inguinal-ovarian-region. Diagnosis: carcinoma of the uterus.

Operation, April 5, 1899: vaginal hysterectomy, by Drs. V. Pleth and C. W. Barrett; angiotribe applied for two and a half minutes, on the right broad ligament, after the usual circular incision and separation of bladder and rectum; re-applied on the opposite broad ligament; uterus cut away, including right ovary and tube; left ovary left *in situ* to prevent storm of the menopause; gauze packing in vaginal vault. The patient made an uninterrupted recovery, the highest temperature after the operation being 100; highest pulse, 90; absolutely no pain or bleeding following the operation. She was sitting up in two weeks, and about in the third week.

CASE 10.—Mrs. H., aged 40; American; first menstruated at the age of 15; always regular but scanty; married fifteen years; never pregnant; had severe attack of leucorrhea twelve years ago—suspected gonorrhea; was somewhat relieved by treatment, but always had a feeling of weakness, and inability to be on her feet for any length of time.

Examination revealed cystic degeneration of the cervix; endometritis, subinvolution, retrodisplacement, with extensive adhesions, pyosalpinx.

Operation, April 6, 1899, at my private hospital; curettement, tracheloplasty, celiotomy, extensive adhesions torn up, double salpingo-oophorectomy, drainage through vagina; no evidence of hemorrhage; almost immediate relief of symptoms; was up on the twenty-third and left the hospital on the twenty-seventh day after the operation.

CASE 11.—Miss M., aged 43, German; menses first appeared at 14; always regular. Three years ago she gave birth to a seven months' child; septic labor; since birth of child, pain at menstrual period, severe for one week before the flow, continuing and becoming worse until it is now constant.

Examination revealed laceration of the perineum, laceration of the cervix, endometritis, retrodisplacement, and inflammatory thickening of both broad ligaments, with extensive adhesions.

Operation, April 13, 1899: curettement, tracheloplasty, perineorrhaphy, celiotomy, adhesions broken up, right salpingo-oophorectomy with clamp; cyst removed from left ovary; suspensio-uteri operation; closure of the abdomen with silk-worm gut suture, without drainage; no evidence of hemorrhage; entirely smooth recovery; got up on the twenty-second and left the hospital on the twenty-sixth day after operation.

CASE 12.—Miss F., age 28, American; teacher; family history negative; first menstruated at the age of 15; pain first two days. One and a half years ago she had a fall, and has not been well since.

Operation, April 15, 1899: left salpingo-oophorectomy by means of the clamp; cyst removed from right ovary; suspensio-uteri operation; abdomen closed without drainage; no hemorrhage; perfectly smooth recovery; sitting up on the eighteenth day; left hospital at end of third week.

CASE 13.—Mrs. R., aged 33, German; housewife; married sixteen years; five children; two miscarriages at four and a half and eight months; first menstruated at 12 years of age; last pregnancy three and a half years ago; difficult labor; not well since; backache, dragging feeling, pain over ovaries and intestinal disturbance, vomiting, constipation. Examination disclosed laceration of the perineum and cervix with large protruding hemorrhoids.

Operation, April 24, 1899, at my private hospital: curettement, tracheloplasty, perineorrhaphy and removal of hemorrhoids by means of the angiotribe and cauterization; prompt recovery; no bleeding, and but a minimum irritation from hemorrhoidal operation; up on the sixteenth day; patient is about to leave the hospital.

CASE 14.—Mrs. K., aged 37, American; housewife; widow; first menstruated at 13; always regular; no pains; two children, 16 and 17; one miscarriage; began to have irregular menses four or five years ago, and latterly has become very profuse; menorrhagia and metrorrhagia. Examination shows large irregular tumor extending to the left and as high as the umbilicus; patient anemic.

Operation, May 7, 1899, at my private hospital: abdominal hysterectomy, with clamp; patient has to date made uninterrupted recovery; no bleeding; no rise of temperature, nor increased heart action.

CASE 15.—Mrs. N., aged 43; German; housewife; family history good; two children; normal delivery; inflammation of the bowels eight years ago; first menstruated at 15 years of age; regular; always with some pain; last ten or twelve

months has had increased pain in pelvis, and more flow, with greater frequency. A mass can readily be palpated over the right iliac and lumbar region, connected with the uterus, nodular in outline. Diagnosis: Multiple fibroids of the uterus.

Operation, April 25, 1899, in the amphitheater of the Chicago Polyclinic: combined vaginal and abdominal hysterectomy with the angiotribe. The Thurin instrument was used here—my own being out of order—and in its double application on each broad ligament the uterine artery was not engaged on one side, necessitating the use of a retention clamp; patient sitting up on the sixteenth day, and will soon leave the hospital; no bleeding or adverse symptoms whatever.

CASE 16.—Mrs. H., aged 54, American; housewife; father died of pulmonary tuberculosis; mother in good health, aged 75; brother 57, good health; all the diseases of childhood; a severe attack of typhoid fever at 12; never strong; two attacks of inflammation of the bowels, twenty and twenty-five years ago; first menstruated at 17; irregular; every two to four weeks; painful; excessive flow, lasting five to six days; leucorrhea, from early menstrual life; married at 21; six children; normal deliveries; one child is tubercular. Present condition: Menopause two years ago; leucorrheal discharge has become worse; with bad odor of late; pains in the inguinal regions and back, increasing in severity and more continuous; anemic. Diagnosis: Uterine carcinoma.

Operation, May 5, 1899, amphitheater of the Chicago Polyclinic: combined vaginal hysterectomy by means of the angiotribe; nothing unusual in the application except the vessels of the broad ligament were large and a pulsating uterine artery led me to apply a retention clamp on one side, and radical operation for hemorrhoids (clamp and cautery). As my own clamp had been used and soiled, I did the operation on the hemorrhoids by means of the Kelley clamp. Three hours after the operation I was called to the Hospital, as the patient had severe flooding. On examination I found that the bleeding was from the hemorrhoidal vessels. Clamps were applied, and no further bleeding took place. Since then she has had an uninterrupted recovery. The gauze has been removed from the vagina, and there is no evidence whatever of bleeding from the broad ligaments, or hysterectomy wound, and she is feeling well.

LA GRIPPE.*

BY J. E. GILCREEST, M.D.

GAINESVILLE, TEXAS.

Influenza, la grippe or epidemic catarrhal fever may be defined as a specific epidemic and contagious disease, caused by a specific bacillus. This disease spreads rapidly over wide districts of country, causing marked febrile symptoms, is often attended by serious complications and causes great and prolonged prostration of strength.

Various epidemics of influenza are on record, Parks tracing it back to the ninth century. The first epidemic in England was reported in 1510, and in the United States in 1647 and at various periods down to the present times. The epidemic described in 1847, in England, was very much like we now find it in America. It has been more prevalent since 1890 than during any previous decade, having spread nearly all over the country in 1890, in April and May, 1891, and in the winter of 1891-92. Only a few cases were reported during the winters of 1892 and 1893. In the early part of 1895 it prevailed epidemically, and again in 1897 and the fall of 1898; and up to this time, April 20, 1899, it is very prevalent in many portions of the country.

Etiology.—At the present time it is generally agreed by medical teachers and writers, that la grippe is a contagious, infectious disease caused by the specific bacillus discovered in the pus-cells of the tracheal mucus, by Pfeiffer, in 1892, and in the blood by Canon, the same year. The toxins developed by these micro-organ-

*Presented to the Section on Practice of Medicine, at the Fiftieth Annual Meeting of the American Medical Association, held at Columbus, Ohio, June 6-9, 1899.