meal. The interior of the kidney should then be flushed with warm, sterile salt solution, so that no portion of the stone may remain behind to furnish the nucleus for a recurrence. The patency of the ureter must always be demonstrated. The hemorrhage should be controlled by hot water and pressure, and care taken that no blood clots be allowed to remain in the pelvis, as they may form the nuclei of new stones, as reported by Tuffier.³⁴ The incision in the kidney is now closed with catgut, and the external wound closed with the exception of a small gauze drain leading down to the kidney, as a provision against leakage. Should all go well, at the end of forty-eight hours the gauze drain should be removed and the wound completely closed. This is the operation of nephrolithotomy in a kidney not the seat of bacterial invasion. When the kidney is already septic, as in secondary stones, primary suture is impossible and drainage must be established. However badly the kidney may appear to be involved, primary nephrectomy in these cases is seldom advisable. It will be impossible to enter into the details of the various operative procedures that have been instituted for the relief of the numerous complications, such as ureteral calculi, strictures of the ureter, sacculated kidneys, abscesses, etc., which may be present in these cases. When both kidneys are affected, it may be necessary to do a double operation, as mentioned by Wagner,³⁵ Pinner,³⁶ Godell and Turner (quoted by Wagner).

The mortality of operation for stone in the aseptic kidney, as given by Albarran (*l. c.*), is a little over 3 per cent.; in septic kidney over 10 per cent.

100 State Street.

BIBLIOGRAPHY.

1. Handbuch. der histor. geograph. Pathol., 2 Auf 1886, iii, 319.
2. Deutsch. Med. Woch., 1890, Nos. 7, 8 and 9.
3. Traité de Path. gén., 1899, Tome iii.
4. Nothnagel: Spec. Path., xix, 396.
5. Deutsch. Med. Woch., 1896, Bd. v, p. 86.
6. Ibid., p. 130.
7. Furbringer: Ibid., 1890, Nos. 7, 8 and 9.
8. Virchow's Arch., 148, xiv, 366.
9. Deutsch. Zeit. f. Chir., x, 126.
10. Bull. de la Soc. de Biol., 1886, pp. 116-373.
11. Jour. de Med. de Bord., 1897, No. 7.
12. Deutsch. Med. Woch., 1897, Nos. 22 and 23.
13. Traité de Chir., Tome vii, p. 914.
14. Arch. f. Exp. Path. u. Pharm., Bd. xxxvii.
15. Deutsch. Med. Woch., 1897, p. 195.
16. Ibid., p. 400.
17. Ibid., Ver Beilage 110.
18. Nothnagel: Path., xix, 394.
19. Deutsch. Med. Woch., 1894, p. 791.
20. Ebstein and Nicolaier: Ueber die Experimentelle Ezeugung von Harnsteinen, 1891.
21. Arch. gen. de Med., May, 1890, p. 572.
22. Arch. de Tocologie, December, 1893.
23. Phila. Med. Jour., Dec. 9, 1899.
24. Deutsch. Med. Woch., 1897, xxii, 337.
25. Arch. f. Klin. Chir., Bd. li, p. 827.
26. Annals of Surg., xxii, 716.
27. Cleveland Jour. of Med., October, 1899.
28. The Lancet, London, April 30, 1898.
29. Deutsch. Med. Woch., 1899, Ver Beilage No. 34.
30. Ibid., No. 6.
31. The Lancet, London, April 30, 1898.
32. Traité de Chir., Tome viii.
33. Deutsch. Med. Woch., 1899, Ver Beilage No. 26.
34. Gaz. Hebdom. de Med. et de Chir., Nov. 28, 1897, p. 1132.
35. Deutsch. Zeit. f. Chir., Bd. lvi, 447.

PATHOLOGY OF RENAL NEOPLASMS.*

BY DANIEL N. EISENDRATH, A.B., M.D.
CHICAGO.

During the past ten years great progress has been made in the pathology of renal growths. Uncertainty and confusion existed in our knowledge of their nature previous to that time. At present the diagnosis of renal tumor or cancer will not suffice. Microscopic examination must be carried out in every case, and growths should

not be reported in the incomplete manner formerly the custom. We now know that the tumors or new growths of the kidney belong to two groups: The benign, including the fibromata, the lipomata, adenomata and angiomata; and the malignant, that includes the carcinomata and sarcomata. An intermediate class is formed by the so-called Grawitz tumors, which may occur in a circumscribed form as benign or in a more diffuse manner as a malignant tumor. These various forms will be taken up in detail later.

In regard to the frequency of renal tumors, one author found only six cases of primary renal growth in 1400 autopsies. Virchow gives $\frac{1}{2}$ of 1 per cent. as the rate of frequency. The kidney is an organ but seldom involved by secondary growths. Kelynnack, in 195 cases of malignant disease, found secondary deposits in the kidney in 8 per cent. of the cases of sarcoma, and in 14.5 per cent. of those of carcinoma. These metastatic tumors do not reach any considerable size, and are generally not numerous. They might therefore be left out of consideration for surgical purposes in speaking of renal neoplasms. In regard to the frequency of the different forms, in 306 cases occurring in both sexes at all ages, there were 187 sarcomata, 142 carcinomata, 15 fibromata, and 12 adenomata. We thus see that for clinical purposes we might leave out the benign forms of new growths.

The etiology of renal neoplasms, like that of growths elsewhere, is still in darkness. Embryonic defects play a distinct rôle. The development of the kidney is very complex, and the inclusion of muscle fibers, both striped and unstriped, of bone and cartilage can be easily explained, some cells of the mesoblast being included in the primitive kidney. This explains the frequency of renal sarcomata in children. The growths arising from adrenal inclusions have been referred to as Grawitz tumors above. Heredity and traumatism seem to have but little influence in the etiology of renal new growths. A direct and not infrequently observed relation between calculi and carcinomata undoubtedly exists. They have been met with in association in a number of cases, and there is sufficient evidence to indicate a clear etiologic connection. Floating kidney has also at times been given as a cause for new growths, but without much evidence in favor of it. Race seems to have no influence. Renal tumors seem to occur more frequently in the male than in the female sex. In regard to age, Kelynnack found that the tumor occurred below the age of 10 years in over 52 per cent. of 160 cases, and of these in about 60 per cent. below the age of 5 years. In regard to size, Spencer Wells has described a tumor weighing sixteen pounds in a child of 4 years. The average weight of twenty-four growths below the age of 10 years was about eight pounds, and of nineteen above the age of 21 years, five pounds. In the average number of cases the general outline of the kidney is retained. This is not the rule, however, for in the nodular form of carcinoma the organ is enlarged at one pole only, while in the diffuse form it is symmetrically enlarged, and consequently much more difficult to diagnose. In all the varieties necrosis, fatty changes and hemorrhages are very likely to occur, and the latter clinically in the form of hematuria. Both sides are affected in equal number. The sarcomata are generally soft, although they may be cystic; the carcinomata harder and more elastic. Renal growths may spread by direct continuity of tissue, by the lymphatics, or by the veins. The growth most frequently pushes aside the organs and tissues of the abdomen, frequently causing ulceration of hollow viscera. Growth into the

*Read before the Chicago Medical Society.

pelvis of the kidney, into the perinephritic tissue and along the vessels of the hilum frequently occurs quite early. The secondary growths are generally found late, and are not extensive. They occur most often in the liver and lungs.

In regard to the duration of renal tumors, the maximum time for sarcomata is $2\frac{1}{2}$ years, the minimum six to eight months. The adrenal inclusion tumors in carcinomata may exist for years.

Pneumonia or secondary growths in the heart or lungs are the most frequent cause of death. The benign growths are of little importance, and I will simply mention the different varieties; fibromata, that occur in the form of pea to bean-sized, firm, whitish nodules; lipomata, which are either in the pelvis or below the capsule. True lipomata are extremely rare. The other varieties of benign growths are the angioma and the adenoma. The latter change very rapidly into carcinoma. The sarcomata form the majority of renal tumors. They may occur in fetal life, although most frequent in children, as stated above. They grow rapidly, and often without symptoms. They are usually of a round or spindle-cell type, and not infrequently striated and non-striated muscle-fibers occur. These latter are due to the inclusion of some of the primitive muscle plates. Myxomatous changes and cyst formation are quite common. Angiosarcomata occur, but are quite rare. Carcinomata occur either as alveolar, glandular, or as papillomatous forms of tumor. They may be diffuse or nodular. They originate from the epithelium of the tubules and are subject to the same degeneration as elsewhere. They do not generally grow as large as the sarcomata, are more frequent in men above 50 years old, although a form of this variety of new growth may occur in children. Epitheliomata may arise from the pelvis of the kidney, usually from a papilloma.

The relation of renal growths to the suprarenal capsule has attracted considerable attention of late years since Grawitz, in 1883, described a form of tumor which had always been called a lipoma, as a new growth due to the inclusion during embryonic life of portions of the suprarenal tissue within the kidney. He proposed the name aberrant adrenal tumor. These may remain quite small, and only be discovered at the autopsy, or may grow very rapidly, as in the case reported by Dr. McArthur and myself, produce metastases, and we have all of the characteristics of a malignant growth. Their structure resembles that of the outer zone of the suprarenal body in having cells arranged in rows containing drops of fat and glycogen. In closing this brief description of the tumors of the kidney, I would like to call attention to the fact that much remains to be investigated in this field. Every tumor should be carefully described and subjected to microscopic examination, and confusion will no longer exist in regard to renal neoplasms.

DISCUSSION ON PAPERS OF DRS. M'ARTHUR, HARRIS AND EISENDRATH.

DR. MAXIMILLIAN HERZOG discussed the peculiar mixed tumors of kidney. He pointed out that we are indebted to Birch-Hirschfeld for having first successfully cleared up the confusion which prevailed as to these new growths, and then quoted, at length, the views of this author regarding the histogenesis of these neoplasms, and also those of Wilms, who contested the theories of Birch-Hirschfeld. Dr. Herzog then narrated his own views as to the histogenesis of these tumors, considering it necessary to enter very briefly into a discussion of the development of the myotomes and of the Wolffian body. These structures are developed from the early mesoderm, which is first differentiated into the myotome and the lateral plate. Between these structures there is formed a

thin bridge known as the nephrotome. In the latter the transverse excretory tubules of the Wolffian duct are formed. While these tubules develop, the nephrotome becomes entirely cut off from the myotomes. This is in short the normal course of events.

With reference to the embryonal, renal adenosarcomata of childhood, he assumes that they owe their origin to an inclusion which is formed in the following manner: The nephrotome is not cut off at the normal site, but in such a manner that a part of the myotome is severed from the main mass and remains in connection with the nephrotome. The separation may take place so that only a part of the myotome proper is cut off, or a part of the sclerotome may likewise be taken along. If the former is the case, we have the matrix for striated muscle fibers only; if the latter occurs, we have also the matrix for cartilage. If, now, it is assumed that a part of the nephrotome (Wolffian body), to which tissues of the myotome have become adherent by an abnormal process of embryonic separation, becomes included in the permanent kidney, we have a matrix containing all those embryonic elements which occur in the mixed renal tumors, namely, striated muscle-fibers, cartilage, other connective-tissue elements, and epithelial glandular structures. The latter are derived from the excretory tubules of the nephrotome.

This explanation appears to Dr. Herzog the most feasible one, in that it will account for the character of these peculiar renal neoplasms without being compelled to take refuge in metaplastic processes unheard of in tumor formation.

The material forming the basis of his studies concerning mixed neoplasms of the kidney, and to which he now calls attention, consists of three tumors.

The first case was one operated on by Dr. Denslow Lewis. It was an enormous tumor weighing $5\frac{1}{2}$ pounds, removed from a girl, 16 months old. This, from a histologic standpoint, presented the most perfect type of a mixed embryonal renal tumor or the embryonal renal adenosarcoma of Birch-Hirschfeld. The glandular epithelial structures were beautifully developed and the connective-tissue elements contained a preponderance of striated embryonal muscle-fibers. In some parts the tumor looked like a pure rhabdomyoma.

For the second tumor which he showed, he was indebted to Prof. L. Hektoen, who had diagnosed it as an embryonal, renal adenosarcoma. This tumor came from a child about 2 years old. Histologically the neoplasm showed epithelial, glandular structures, and various connective-tissue elements. Striated muscle-fibers were not found, in spite of a careful search for them.

The third tumor was from a child 9 months old, who came under the observation of Dr. Frank Churchill, and was operated on by Dr. M. L. Harris. He had examined the tissues of this tumor from thirteen different places, yet he had not been able to find any epithelial glandular structures, but only connective-tissue elements of very heterologous types, yet without any striated muscle-fibers. This third tumor, like the two preceding ones, presented those clinical features as they were enumerated above, as characteristic of the embryonal mixed tumors of early childhood. It therefore clinically belonged to this group; yet histologically it presented that type the possible occurrence of which Birch-Hirschfeld predicted in his article on the embryonal adenosarcomata of the kidney, a type characterized by a most extensive embryonal connective-tissue proliferation, yet without epithelial glandular structure, which would therefore have to be classified as an embryonal renal pure sarcoma. The third case presented a tumor of this type.

DR. WELLER VAN HOOK said that before an operation is undertaken on a kidney, the normal one, and the supposedly abnormal one also, are to be considered. For a knowledge of the abnormal one and its condition, the surgeon depends partly on palpation, but very largely on the urinary findings. In the latter he spoke of hemorrhage, which is of particular importance. The urine should be frequently examined for weeks. Having established the presence of blood in the urine, even in microscopic quantities, it is very important to determine its origin. Any examination of this kind is not complete without a cystoscopic one. As illustrating the every-day importance of the use of the cystoscope, he detailed two instructive cases.

In 1892, without having heard of the work of Fenger, he incised the skin and aponeurosis, separating the muscles over the left kidney, which proved to be normal, in a case in which the right kidney was tuberculous. In 1893, Dr. Fenger published the report of a case on which he had operated in 1890 by this method, determining the presence, size, and the proper consistency of a supposedly normal kidney. He spoke of this because in all cases since then, in which he has thought it desirable to remove the kidney, he has palpated the supposedly normal kidney in this manner. An incision two inches in length in a patient who is not more than ordinarily fat can be made, and the kidney easily palpated without running the risk of subsequent hernia. The risk of infection is slight.

DR. E. WYLLYS ANDREWS said that neoplasms of the kidney of the type under discussion were essentially insidious in their development. They have been repeatedly mistaken in diagnosis for other forms of abdominal tumor. The cachexia characteristic of these growths in other locations is frequently absent in kidney neoplasms. The tumor is not always easily palpable, and in many instances it is overlooked simply because a careful examination is not made. When such a tumor is found, it is usually rounded, and somewhat easily palpable. Pain is a symptom of value in the diagnosis. It is of a dull, aching character, and not increased by walking. It may be absent, however, in some cases. The treatment of these renal neoplasms is surgical. He thinks one can hardly overestimate the value of the Harris segregator. It is an instrument easy to use, and it gives positive results in a large proportion of cases, with very little risk to the patient.

DR. GUSTAV KOLISCHER disagreed with Dr. Harris that we can rely for diagnostic purposes on pain and dullness in the region of the kidney, also the examination of the urine, collected and separated from each kidney by ureteral catheterization, or by the use of the Harris instrument. Many cases are met with in which the urine can not be collected satisfactorily from each kidney, hence the surgeon is unable to draw any definite conclusion from the small amount of urine passed. The greatest value of ureteral catheterization, or, better still, ureteral sounding, lies in whether the ureter is permeable or not, or whether there be a stone in the ureter or in the pelvis of the kidney. Cases occur in which it is impossible to locate the exact seat of the pain, for the reason that it radiates all over the chest, stomach and abdomen; at the same time, the patient may be in such a abdominal condition as to call for immediate interference. Sounding of the ureters in such cases is a valuable diagnostic measure. Dr. Kolischer then, by means of sketches, showed what can be accomplished by ureteral sounding. It is possible to push a stone back into the pelvis of the kidney, rendering it accessible for surgical interference later. In his opinion, ureteral catheterization can be resorted to successfully only by experts. The same is true of the Harris segregator. The latter instrument should be used only by skillful and trained hands, because the urinary organs are not to be trifled with.

DR. EDWARD H. OCHSNER discussed the historic aspects of adenoma of the kidney and of the so-called adrenal tumors described by Grawitz. The aberrant tumors of Grawitz cause severe systemic disturbance; the principal symptom is greatly increased arterial pressure. The patient rapidly becomes very sick, and dies usually with a hemorrhage into the brain or other important structures.

DR. JAMES B. HERRICK said that adrenal tumors not infrequently produce bone metastases in the same manner as tumors of the mammary gland, of the prostate, and of the thyroid, so that in case of bone metastasis, where the primary trouble can not be exactly located, the region of the kidney should be interrogated for possible tumor of the adrenal.

As to the diagnosis of renal neoplasms, some authors place considerable stress on the occurrence of varicocele on the right side as indicative of pressure on the spermatic vein, while others give it a passing notice. In renal tumor there may be a distinct systolic bruit or murmur, sarcomata rich in blood-vessels may give rise to a very loud systolic bruit, and with a palpable tumor aneurysm might be suspected. A tumor of the kidney may simulate an aneurysm by pulsation. The distinct expansile pulsation may be pronounced in cases of renal tumor rich in blood-vessels.

A few weeks ago Dr. Murphy showed him a case in which there was a large mass in the left side of the abdomen, in front of which was the colon. The patient, a man, 40 years of age, had had some hematuria, and showed distinct secondary anemia and cachexia. This mass enlarged very rapidly as the symptoms only dated back two or three months. He has never seen, in any case of aneurysm, a more perfect expansile pulsation than there was in this large mass. From the hematuria, the location of the mass—which reached to the median line and bulged between the flanks—the cachexia, and the relation of the mass to the colon, it seemed to be a large tumor of the kidney.

DR. BERTRAM W. SIPPY has had occasion to look up the subject of tumors of the kidney very thoroughly from a clinical and pathologic standpoint, and, so far as he knows, there have been only four cases reported in which adrenal tumors of the kidney have been accompanied by greatly increased arterial tension. Two cases, which he observed post-mortem in Vienna, were accompanied by cerebral hemorrhage sufficient to cause death. The arterial tension was high in these. Neisser has reported a case in which the arterial tension was very high, likewise Dr. Holmes. He did not consider increased arterial tension a very valuable diagnostic sign, because in twenty-nine cases reported by one author no clinical manifestations were noted.

DR. FRANK BILLINGS does not think the treatment of renal calculus is purely surgical, for there are many cases which, by the grace of God, and the drinking of lots of water, get well, and nothing further is heard of them after the first attack. It is not unusual, in taking the histories of patients, to have them say that they have suffered from renal colic years before, and have escaped further attacks. He has had, in his own experience, patients who suffered from renal colic at intervals of every two or three years, but who, under a systematic course of hygiene, careful dieting, plenty of fluids, as diluents, would cease to have these attacks, and he believes that such hygienic treatment is scientific and not entirely symptomatic. While he has not looked up his case histories, nor would he care to state exactly how many cases have been operated on, yet he believes that he has had more recoveries from such treatment as he has mentioned than from operative intervention.

DR. HAROLD N. MOYER saw at least a dozen cases of renal calculus in the years of his earlier practice that were not operated on, and the patients are well to-day.

DR. L. L. MCARTHUR, in closing the discussion on his part, emphasized the use of the ureteral catheter in septic conditions of the bladder. The segregator should be used rather than take the chances of infecting one kidney which it is hoped is normal.

DR. DANIEL N. EISENDRATH narrated a case that was operated on by Dr. McArthur and himself with unfortunate results. A man came to him three years ago, presenting all the symptoms of renal calculus. Operation was advised, but refused. The man was placed on medical treatment for nearly two years, at the end of which time he returned with pyuria. He was very much emaciated, and had chills and elevation of temperature. The urine was carefully examined with the Harris segregator, with a view to determining the working capacity of both kidneys; apparently each kidney secreted the normal amount of urine, although the urine secreted from each side contained some pus corpuscles. The percentage of urea was normal, and there was no albumin other than that which could be ascribed to the presence of pus corpuscles. Operation revealed several large stones in the pelvis of the kidney with quite an amount of detritus. Later the man developed anuria, and died. Post-mortem examination showed a kidney considerably smaller than normal on the opposite side. In addition to one large calculus, the six or seven smaller ones removed varied in size from a pea to a marble, each one of which showed the characteristic phosphatic coating over the urates, showing that medical treatment had resulted in an extra deposit of the previously existing calculi.

DR. M. L. HARRIS, in rebuttal, stated that Dr. Kolischer had misunderstood him. He did not say in his paper that the only value of catheterizing the ureters was in collecting the urine from each kidney. He did not go into details regarding this point in his paper.