has occurred in the use of the vaccine. Of course, it is not suggested for a moment that the above is the condition prevailing normally in the Manila Laboratory. The latter has, in a few years, and most deservedly, taken a place amongst the first-class institutions of its kind in the world. It is only unavoidable now to make clear that their accident has not been conditioned by the peculiarities of the anticholera vaccination method, as they believe it has.

It is to be hoped that the above explanations will not be viewed as implying any want of consideration for the Manila scientists whose efforts, ever since the establishment of their laboratories, have enriched science with numerous contributions of a truly remarkable character.

SUBMUCOUS PERINEORRHAPHY

A REPORT BASED ON SIX YEARS' EXPERIENCE WITH AN ORIGINAL METHOD *

W. WAYNE BABCOCK, M.D.

Surgeon-in-Chief to the Samaritan Hospital
PHILADELPHIA

The methods of perineorrhaphy that have been chiefly employed for the past twenty-five years suggest more of a mathematical than an anatomic basis for their existence. For the most part, they have consisted of excisions of mucous membrane from the posterior vaginal wall having geometrical patterns that vary as do the fancies of different surgeons. In support of the various operations much has been written about the laceration in the muscular and facial planes and of the effectiveness of particular operations, yet one who studies the work of various gynecologists will be impressed by the thought that usually the precise anatomic restoration of the perineum occurs only in the theory of the operator, for the operation, as a rule, consists of little more than the removal of an area of mucous membrane and the union of the wound edges. If at times the operator's needle is made to sweep in various directions with the specification that certain muscles are caught in its grasp, the precise evidence that such muscles are included and especially any evidence that the important fascial planes of the perineum are restored is rarely observed.

In the Emmet operation, which has long been praised as the best method of restoring the pelvic floor, the crest of the rectocele is caused by the crown stitch to be brought close to the posterior edge of the vaginal opening. As normally the point indicated by the crest of the rectocele occupied a position some distance above the introitus, the operation shortens the posterior vaginal wall. The lateral triangular denudations obliterate the normal lateral sulci; so that after the Emmet operation the vagina in this region loses the normal H-shaped section and becomes less dilatable. But few operators carefully investigate the muscular structures in doing an Emmet operation and fewer make an exposure of the muscle or use an accurate method for the muscular restoration, while the fascial planes of the pelvic floor are almost invariably ignored in the operation, although they may at other times be the basis of an elaborate theoretical discussion. In this type of operation the through-and-through sutures serve as setons, carrying vaginal secretion to the depths of the wound, and by irritation tending to in-

crease the amount of fibrous tissue in the perineum. The sutures in the vagina entangle secretion, interfere with proper cleanliness and at times are difficult of removal, and if overlooked or partly removed may remain in the tissues and cause abscess or sinus formation, and protracted disability. Similar serious objections may be brought against most of the conventional operations for the restoration of the perineum. In an endeavor to obviate some of the disadvantages of the older technics I devised a method of perineorrhaphy that was first put in execution in 1902. The operation proved to be satisfactory and the method has been demonstrated to students and acquaintances for the past six years, and during this time I have employed it exclusively for all secondary repairs of the perineal floor. In 1904, my colleague, Dr. A. C. Applegate, delivered without perineal laceration a patient on whom the operation had been done the previous year, of a full-term child, and the experiences of others who have tried and adopted the method and the tendency of certain other operators¹ to devise independently, methods of perineal repair based on similar principles emphasize the conclusion that this is a type of operation distinctly superior to the older operations of Emmet or Hegar. To object to the present operation because it gives a greater muscular and fascial support to the vagina than is normal is not consistent with the acceptance of modern operations of herniotomy.

The operation is based on the following principles:

First, no tissue is removed or extensive denudation made.

Second, buried, absorbable, layer sutures are used exclusively, none of which penetrates the skin or mucous membrane.

Third, the operation is done from the outside of the vagina, rendering the introduction of sutures easier and the exposure of tissues better than with those operations done from within the vagina.

Fourth, each structure is sutured with precision under the guidance of the eye; there is no blind groping with the needle for tissues not seen and perhaps not felt.

Fifth, each of the layers of the perineal floor—vaginal wall, submucosa, muscular supports, fascial planes, and skin—are united seriatim in layers after the plan of the better types of herniotomy.

Sixth, the vagina is not separated from the rectum and therefore there is no danger of wounding the bowel.

The operation has the following advantages:

It restores and increases the length of the vagina instead of shortening the posterior wall of the vagina as occurs with many of the older operations. Although no tissue is removed it is questionable if any other type of perineorrhaphy is capable of producing greater narrowing of the external portion of the vagina or of affording greater support to the anterior vaginal wall. As it greatly supports the anterior vaginal wall, slight or moderate degrees of cystocele may not require additional operation. I have only done the additional operation when the cystocele was marked and yet have thus far had no reason to regret the practice. The operation restores the normal H-shaped section to the vagina instead of destroying or obliterating the lower lateral sulci. There is the added advantage that no sutures require removal; that no foreign bodies in the shape of shot, suture ends, or the like are left in the vagina to promote and to hold secretion and that the

* Read before Philadelphia Obstetrical Society March 4, 1909.

1. Hill: Restoration of the Pelvic Floor. THE JOURNAL, April 4, 1908.

seton action of through and through sutures is avoided. Moreover, the proneness of knots in catgut to untie when exposed on a mucous membrane is obviated by the burial of all knots in the tissue. Despite the number of buried sutures in none of my cases has the perineum been destroyed by suppuration, and the resultant perineum is more elastic and resilient as less fibrous tissue forms than after the older operations with the use of denudation and through-and-through sutures. In one of the early cases in which the mistake was made of freely separating the vagina from the rectum, the rectum was opened with the result that the posterior portion of the wound became infected. In this case the vaginal support, however, was not destroyed. In my experience in only one case has the perineum given way and in this

In doing the operation, one should realize that the vagina runs nearly parallel with the skin perineum, and that the finger introduced to the cervix may be but half an inch distant from the finger of the external hand placed on the skin adjacent to the coccyx. Lateral to the vagina one may nearly always distinguish the thick broad bands of the levator ani muscle. In the nulliparous woman the muscle is close to the introitus, in the multiparous it is more deeply situated. By placing a finger against the anterior edge of this muscle on either side and comparing the relative depths of the muscle from the surface, one may acquire an idea as to the relative injury to the two muscles. Rarely ever will one find the muscle completely divided lateral to the vagina; but on the side of greater laceration the

Fig. 1.—Step 1: The first incision made by scissors or knife at the base of the carunculae. The incision is U-shaped, enters the submucous areolar tissue, and is slightly external to the orifice of Bartholin's glands. According to the degree of contraction of the orifice required, the incision starts and ends anterior to, opposite to or posterior to the level of Bartholin's glands.

the healed wound was ruthlessly torn open by the husband a week or ten days after the operation.

Comparing my experience with the Emmet operation with the present one under equal aseptic precautions, the liability to suppuration seems less, the required period of disability shorter, and the inconvenience of suture removal, and the danger of non-absorbable sutures obviated in the present operation. Comparing the end-results of the two methods a more perfect perineum with a better muscular support and a closer approximation to a normal restoration is secured by the method of submucous perineorrhaphy.

Fig. 2.—Step 2: Isolation and exposure of the internal edges of the levator ani muscle. The vagina is not separated from the rectum in the median line, and therefore there is no risk of wounding the rectum. The posterior edge of the incision is retracted by a tenaculum to secure better exposure. A sharp pair of scissors is then thrust through the fascial plane, lateral to the vagina, opened and withdrawn, affording access to the levator ani. This incision is enlarged by stretching with the fingers (as indicated by the dotted lines), or, if the tissue be cicatricial, by further incision. The finger is then thrust through the incision close to the external wall of the vagina and the levator ani isolated from the vagina. The external surface of the levator ani is likewise freed and the muscular edge pulled into the wound by a hook or tenaculum forceps. The edge of the opposite half of the levator ani is similarly freed and brought into the wound.

edge of the muscle will usually be found to be from one to several centimeters deeper than that of the opposite side.

TECHNIC OF THE OPERATION

The parts are carefully aseptized in the usual manner and the anus covered by a sterile towel or gauze

pad held to the skin by appropriate clips. The labia having been separated and supported by the fingers of an assistant, the sharp point of a pair of scissors is introduced into the subcutaneous cellular tissue just external to the orifice of Bartholin's gland. The incision is carried around the posterior margin of the introitus just external to the carunculae until a point external to the orifice of the opposite Bartholin's gland is reached. A pair of tenaculum forceps is then fastened to the outer edge of the posterior part of the incision to serve as a retractor. No separation of the vagina from the rectum is necessary except when the tissues about the posterior commissure are so cicatricial that the usual gaping of two or three centimeters of the wound edges is not obtained. If the posterior wall of the vagina is

edge of muscle, then opened and withdrawn. This penetrates the remains of fascia of Colles, the two layers of the triangular ligament and perhaps the thin and at times almost inconspicuous layer of the transversus perinei profundus which lies between. The opening is enlarged by stretching with the fingers, or if necessary, by a few touches of the knife, and the finger is passed into the opening close to the lateral vaginal wall until the edge of the levator ani is felt. The muscle is freed on its inner and outer sides by the finger, grasped with a pair of tenaculum forceps, and pulled into the wound. On the side of greater laceration the muscle will usually be found to have the deepest situation. The isolation and exposure of the muscle is usually easy and requires but a few seconds of time. Very rarely will the muscle

Fig. 3.—Step 3: Submucous closure of vaginal wall. With a continuous suture of plain, number one catgut, beginning below, the edges of the vaginal wound are united. This suture does not penetrate the mucous surface and serves to lengthen the vagina and to narrow the introitus. If the posterior vaginal wall be very redundant, this first suture may be buried by a second row of interrupted or continuous sutures, catching the tissues lateral to the first and so rendering the vaginal wall thicker, more rigid and more convex. When a level above that of the emergence of the muscles has been reached this suture is temporarily laid aside.

bound down and deformed by cicatricial tissue this should be loosened sufficiently to overcome the deformity (Fig. 1.)

The second step consists in the exposure of the levator ani. The edge of the muscle is located between one finger placed against the lateral wall of the vagina, and a second finger or thumb placed in one side of the wound. Having located the muscle in the wound, a pair of sharp-pointed scissors is thrust through the depths of the wound to the situation of the anterior

Fig. 4.—Step 4: The edges of the levator ani muscles are united in the median line under the united vaginal wall by mattress sutures. This muscular sling accentuates the lateral vaginal sulci and restores the normal H-shaped section to the vagina. In certain cases a continuous or simple interrupted suture may be substituted. Some of the sutures should catch the under surface of the vagina to prevent the formation of a dead space. Below a modified figure-of-8 suture is shown, which serves to unite the levator ani with the sphincter ani and to support the sphincter. This figure-of-8 type of suture in this situation is not usually required and frequently no support or reinforcement of the sphincter is necessary. If the sphincter be divided, however, the ends are to be isolated in the wound and united by buried sutures of chromicized catgut, and in extreme cases the sphincter may also be reinforced by slips of the levator ani split off from each internal edge, crossed and wrapped about the sphincter and sutured into place. No additional incision is required in a complete tear, but merely sufficient dissection in the wound to expose the divided ends of the sphincter, which are then united.

be found to be completely divided. The opposite edge of the levator ani is then exposed and brought into the wound in the same manner (Fig. 2).

The third step of the operation consists in the repair of the vaginal mucous membrane. With a curved needle and No. 1 plain catgut the mucous membrane is united beginning near what was the posterior commissure. A continuous suture is used engaging the submucous cellular tissue a short distance from the wound edges and uniting the upper border of what was a transverse incision in a vertical manner. In doing this, the part of the vagina that was near the posterior commissure comes to occupy a point several centimeters above the introitus and the length of the vagina is restored. If there is much redundancy a second row of sutures may be applied catching the tissue outside suture number one and so rolling the submucous tissues together that the convexity on the mucous surface of the posterior

Fig. 5.—Step 5: Suture of the inferior fascial plane (urogenital trigone) over the united muscles. This supports and reinforces the united muscles, and, as the transversus perinei profundus muscle lies within this fascia, the action of this muscle is reinforced. A continuous, interrupted, mattress, or imbricating suture may be employed, depending on the strength of the suture line required in the particular case. The ends of suture one are shown emerging above and below.

wall of the vagina is markedly increased. Suture number one which has thus made a partial submucous closure of the vaginal wall is then temporarily laid to one side to be finished later (Fig. 3).

The fourth step of the operation consists in uniting in the median line the edges of the levator ani muscle, which are located by the attached tenaculum forceps. Interrupted, continuous or mattress sutures of chromic catgut may be employed, but the muscle must not be united so far anteriorly as unduly to constrict the orifice. Some of these sutures catch the underlying submucous tissue to prevent the formation of a dead space. Posteriorly, a simple or figure-of-eight suture is employed to bind together the various structures that meet here, and if necessary, to give support

to the sphincter ani. In certain cases, it may be desirable to reinforce the sphincter ani by splitting off strips from the anterior edges of the levator ani, wrapping these about the sphincter and fixing them in place by sutures. Rarely is a similar procedure desirable anteriorly to reinforce the vaginal support. Should the levator ani be found to be torn, which is not usual, the divided ends should be pulled well forward, carefully isolated and repaired by buried sutures. With the union of the levator ani muscles the rectum, anus and vagina will be found to ascend to a higher position in the pelvis, and to be carried forward toward the pubis (Fig. 4).

The fifth step of the operation consists in the suturing of the inferior fascial plane to reinforce the muscular support. The outer edges of the incision through which the levator ani muscles have been brought are united

Fig. 6.—Step 6: The unfinished suture of step three is now continued under the cutaneous edges of the incision until the fourchette is reached, where the two ends of this suture are tied and the knot permitted to sink into the wound out of sight, the ends being cut short. The tying of the ends of this first suture binds the various structures—vagina, muscle, fascia and skin—together and prevents the formation of dead spaces. This elevation and reinforcement of sphincter ani is indicated.

in the median line by continuous, interrupted, mattress or imbricating sutures of No. 1 chromic catgut, as may seem necessary in the particular case. This support, which includes the posterior part of the urogenital trigone, affects the layers of the triangular ligament and the deep transversus perinei muscle, and posteriorly serves to increase the tension on the transversus perinei muscle. To obliterate any dead space some of the sutures may catch the under-lying muscle (Fig. 5).

The sixth and final step of the operation consists in completing suture number one which after completing the submucous union of the vagina is continued posteriorly under the skin uniting the subcutaneous and Colles fascia until the posterior portion of the incision is reached. The two ends of suture number one are

then tied, binding together the various structures that have been united; the knot is permitted to sink under the skin, the ends being cut short (Fig. 6).

Thus there has been built up in layers a perineum consisting of united mucosa, muscle, fascia and skin. All the sutures being buried and the layers bound together compactly there are practically no dead spaces. A small strip of iodoform gauze introduced into the vagina is permitted to hang down over the skin sufficiently to cover the skin incision. This serves as a guide to the nurse in catheterizing and protects the incision. The catheter is used for two days after which the gauze is removed and if desirable a daily douche is given. The patient may sit up on the tenth day, and cautiously resume her duties after two weeks.

2033 Walnut Street.

THE TREATMENT OF SCOLIOSIS

ARTHUR STEINDLER, M.D.

Assistant Orthopedic Surgeon to the Home for Destitute and Crippled Children
CHICAGO

In treating scoliosis by means of orthopedic corsets I think that it is difficult to meet the two most important requirements, the one of giving a reliable and uni-

marily involved in the curvature, not enough hold may be gained for correction and the maintenance of the corrected position of the spine.

The sideward shifting of the body will always result in a lifting of the pelvic belt of the corset from the crista ilei of the opposite side, unless its position there is otherwise secured. In order to obtain this advantage, and, on the other hand, to exert some correction on the costal prominence, I attached to the pelvic belt on the healthy side a thigh-piece connected with a hinge and embracing the leg by means of a padded ring right above the knee.

On the side of the scoliosis a pad corresponding to the costal prominence articulates by means of a hinged steel piece with the pelvic belt, a similar pad being attached on the corresponding anterior costal prominence of the opposite side.

On both pads a considerable amount of elastic pressure is secured by a couple of springs. It is easily to be seen that, if the attachment to the leg is given an outward bent and then fastened closely above the knee, an opposite movement of the pad will result in bringing the sideward shifted body over the median line to the other side.

Two uprights in the back on both sides of the spinous processes give the body rest and allow the shoulders to be pulled back by means of shoulder-straps (Figs. 1, 2, 3 and 4).

Both patients with right dorsal lateral curvature, shown in the photographs, were given a previous mechanical treatment of several months. It consisted in gymnastic exercises, manual redressement, especially directed toward the lumbar curvature and the detorsion of the scoliosis, and some massage.

After I had convinced myself that the lumbar portion of the spine had gained a sufficient amount of flexibility I had the patient wear the contrivance, as described above, while the gymnastic exercises were being continued.

The considerations which led me to pursue this mode of treatment are, briefly, the following:

I have to a great extent given up mere extension as a method of correcting curvature, because of the fact that the detorsion of the body would not keep pace with the extension of the

Fig. 1.—Patient with scoliosis.

Fig. 2.—Same patient as in Fig. 1, with orthopedic corset.

form support to a deformed spine, the position of which has been improved by mechanic or gymnastic treatment, and the other of exerting a correcting influence on certain parts of the body by means of the orthopedic corset itself.

A number of good and practical corsets have been devised, giving satisfactory results to a considerable extent. While the Hessing corsets furnish only a thorough support to the spine, a good correcting corset has been described lately by Bradford, the correcting force being exerted by means of shoulder straps, circular transverse springs and a swinging upright for pressure on the neck. But it seems to me that in cases in which there is a considerable shifting of the body to one side, for instance, in which the lumbar part of the spine is pri-

marily involved in the curvature, the formation of flail intervertebral joints and increased flabbiness of the spine being the result. Most of my attention is directed toward the detorsion of the chest and the mobilization of the lumbar part of the spine, which, as a rule, is the last to maintain a certain degree of flexibility, when the dorsal part of the spine is already hopelessly fixed. Then, if the sideward displaced body is shifted to the other side by means of the apparatus and an over-correction exerted, the upper part of the spine will have to respond with a self-correcting secondary curvature to the side of the disease in order to regain the median line.

Thus a considerable improvement of the deformity is caused under the influence of weight-bearing and without causing flail joints, while other