21. Mitchell: Glasgow Medical Journal, vol. xlv.
22. Wiener med. Presse, 1898, p. 1477.
23. Says the Spectator of Feb. 9, 1901.
24. Article on Malaria, Twentieth Century Practice of Medicine.
25. The Quarterly Review, July, 1899.
26. Travels in the Equinoctial Regions of South America, vol. ii, p. 205.
27. In the New York Evening Post for Sept. 6, 1901.
28. There was a play published in 1768, called "The Devil on Two Sticks," by Samuel Foote. In it the President of the College of Physicians is represented as delivering a lecture. He says: "Brethren and Students—I am going to open to you some notable discoveries I have made respecting the source or primary cause of all distempers incidental to the human machine; and these, brethren, I attribute to certain animalcule, or piscatory entities, that insinuate themselves through the pores into the blood, and in that fluid sport, toss, and tumble about like mackerel or codfish in the great deep; and to convince you that this is not a mere *gratis dictum*, an hypothesis only, I will give you demonstrative proof. Bring hither the microscope! Take a peep. . . . Those two yellow drops were drawn from a subject afflicted with the jaundice—What d'ye see?"
29. *Lust*.—Some little creatures like yellow flies that are hopping and skipping about.
30. *Hcl*.—Right. Those yellow flies give the tinge to the skin, and undoubtedly cause the disease; and now for the cure! I administer to every patient the two-and-fiftieth part of a scruple of the ovaria or eggs of the spider; these are thrown by the digestive powers into the secretory, there separated from the alimentary, and then precipitated into the circulatory, where finding a proper nidus or nest, they quit their torpid state and vivify, and upon vivification, discerning the flies, their natural food, they immediately fall foul of them, extirpate the race out of the blood, and restore the patient to health.
29. Transac. Congress Amer. Phys. and Surg., 1891, vol. ii.
30. The Practice of Medicine Among the Burmese. Edinburgh, 1879.
31. Chapter on Inoculation, Nothnagel's Encyclopedia of Practice of Medicine, American Edition, 1902, vol. ii.
32. Nothnagel's Encyclopedia of Practical Medicine, American Edition, vol. ii, 1902.
33. Edinburgh Review, 1899, p. 345.
34. Says the British Medical Journal of Dec. 14, 1901.
35. Editorial in American Medicine, July 27, 1901.
36. Exposition Times, quoted in above editorial.
37. The following letter of a traveler in Bengal (*Revue Scientifique*, February, 1901, reprinted from *Gazette hebdomadaire de Med.*) is of interest. He says: "Three months ago a mad dog bit six or seven men, among them two of my bearers, wounding them badly. I at once had some iron heated white hot to cauterize the wounds. But the natives looked on laughingly. 'El Sahib' said they, 'it's nothing at all; we have an excellent remedy; you shall see.' The dog ran again. One of the men seized a stick and killed him; another ripped open the abdomen, took out the palpitating liver, cut some pieces off and gave them to each of the wounded men, who swallowed them, raw and bloody as they were. 'The danger is over now,' they said. As I was incredulous they brought me a young man on whose legs were large scars. Bitten by a mad dog some five years before, this man had eaten a piece of the animal's liver and had felt no evil effects from his wound. The case I witnessed happened in March. It is now the 3rd day of July. The wounds have healed and the men continue in good health. The natives even go so far as to maintain that if this remedy be given to a man already stricken with hydrophobia, it will infallibly cure!"
38. *Revue Scientifique*, Feb. 20, 1892.
39. *Scientific American*, Oct. 7, 1893.
40. *Jour. Amer. Folk Lore*, July, 1901.
41. *Jour. Amer. Folk Lore*, April, 1901.
42. G. A. Stockwell: *Ther. Gazette*, 1887.
43. *American Medicine*, Sept. 20, 1901.
44. Weiss: *Wiener Med. Presse*, 1898, p. 1477.
45. *Med. and Surg. Reporter*, March 26, 1881.
46. *Jour. Amer. Folk Lore*, April, 1901.
47. Flückiger and Hanbury's *Pharmacographia*.
48. F. P. Smith: *Mat. Med. and Nat. Hist. of China*, Shanghai, 1871.
49. Pliny: *Lib. xxvii*, C 76, also xxv, 25; quoted by Flückiger and Hanbury.
50. Flückiger and Hanbury's *Pharmacographia*.
51. R. H. True: *Jour. Am. Folk Lore*, April, 1901.
52. *History of Coca*. J. H. Vail & Co., N. Y., 1901.
53. *Johns Hopkins Bull.*, April, 1902.
54. *Pharmacology and Therapeutics*, 1880. *Gulstonian Lectures*.
55. *Glasgow Medical Journal*, vol. xlv.
56. R. H. True: *Jour. Amer. Folk Lore*, April, 1901.
57. L. E. Monnet: *Therapeutic Gazette*, 1885.
58. Hoffmann: *Medical and Surgical Reporter*, Feb. 22, 1879.
59. True: *Jour. Amer. Folk Lore*, April, 1901.
60. *The Medical Lore of Celtic Scotland*. The American Medical Journal, June, 1896.
61. W. Hamilton: *History of Medicine from the Creation to the Nineteenth Century*, London, 1831.
62. 1. ATROPIN. Discovered by Brandes, in the *Atropa belladonna*. 2. BRUCIN. Disc. in 1819, by MM. Pelletier and Caventou, in the false angustura, in the state of a gallate. 3. CATHARTIN. Disc. in 1820, by MM. Lassaigue and Feneulle, in the pods and leaves of senna. 4. CINCHONIN. Disc. by Duncan, in cinchona bark. 5. CYSTISIN. Disc. in 1820, by MM. Lassaigue and Chevallier, in the seeds of the *Cystitis laburnum*. 6. DELPHIN. Disc. in 1819, by MM. Lassaigue and Feneulle, in the seeds of stavesacre (*delphinium staphysagria*). 7. DIGITALIN. Thoughts of its existence entertained by MM. Lassaigue and Chevallier; actually discovered by M. Le Royer, in 1824, in the leaves of the *Digitalis purpurea*. 8. EMEZINE. Disc. in 1817, by MM. Pelletier and Magendie, in the roots of the *Viola emetica*, *Callicocca ipccacuanha*, and *psycotria emetica*, and by M. Boullay in the violet. 9. GENTIANIN. Found by MM. Henry and Caventou, in the root of gentian. 10.

106E. Disc. in 1813, by Courtois, in the mother waters of soda, as it is obtained from the sea-weed. 11. LUPULIN. By M. Planche, and nearly at the same time by MM. Ives, Payen and Chevallier, in the yellow matter of the flowers of hops. 12. MORPHIN. Disc. in 1817, by Serturner, in opium, in the state of a meconate. 13. NAUCOTIN. Disc. in 1802, constituting one-fiftieth of opium, by M. Derosne, who named it opiane. 14. PICROTOXIN. Disc. by Boullay, in the *Coccus orientalis* (the fruit of the *Menispermum coccinus*). 15. PIPERIN. Disc. by Erstaed, in black pepper. 16. QUININ. Disc. 1820, by MM. Pelletier and Caventou, in the cinchona. 17. RHUBARBARIN. Disc. by Pfaff, in the rhubarb cultivated in Europe. 18. SCILLITIN. Disc. by Vogel, in the bulbs of the *Scilla maritima*. 19. SOLANIN. Disc. in 1821, by Desfosse in the *Solanum dulcamara* and in the *Boletus esculentus*. 20. STRYCHNIN. Disc. in 1818, by MM. Pelletier and Caventou, in the *nux vomica* and the bean of St. Ignatius. 21. VERATRIN. Disc. in 1819, by MM. Pelletier and Caventou, in the cevadilla, white hellebore, and in the colchicum.—[From the *Lancet*, Nov. 15, 1902.]

63. Boston Med. and Surg. Jour., 1882.
64. Trans. New York State Medical Assoc., vol. viii.
65. J. M. Magill: Fort Wayne Medical Journal-Magazine, February, 1898.
66. Transac. New York State Med. Assoc., vol. viii.
67. J. Spear: The *Lancet*, 1878, vol. ii.
68. *Odyssey*: Book xxii, 481 and 482.
69. *Lancet*, Jan. 2, 1892.

THE SINGLE CUFF METHOD OF CIRCULAR ENTERORRHAPHY:

A NEW METHOD.*

O. BEVERLY CAMPBELL, A.M., M.D.
ST. JOSEPH, MO.

A NEW METHOD.

A consideration of the evolution in the methods of circular enterorrhaphy occurring in the past seven centuries, with the use of mechanical devices and sutures, and by the suture methods alone, is indeed interesting.

The goose's trachea of the four masters, wood canula, dried intestine, Scarpa's tallow candles, Watson's gelatin tubes, Chopart and Sabatier's prepared cardboard, later, Senn's bone plates, Paul's bone tube, Ashton's rubber rings, Murphy's button, Jessett's bone tubes, Brokaw's segmented rubber rings, the potato cylinder, the forceps of Laplace and others, the rubber bulbs of Treves, represent the progress in mechanical devices; while the Glover suture and the open treatment of the wound, Jobert's method, the Lembert suture, the Czerny-Lembert, the Maunsell method, Chaput's method, the Harris method, Kummer's method and the Connell suture, represent the progress in suture methods.

Considering the multiplicity of methods now in use, by mechanical devices and by suture, leads one to conclude that circular enterorrhaphy is still in an evolutionary state, that an ideal method had not as yet been devised.

Since Richat, Richerand and Treves, in the early part of the nineteenth century, taught that union is more readily obtained between serous surfaces than between mucous surfaces or between serous and mucous surfaces, nearly all of the methods that have since been originated have adhered to the principle of obtaining union of serosa to serosa. The Lembert suture, placed through serosa, muscularis and submucosa with the knot outside of the lumen of the bowel, is probably the most universally popular method of suturing. Yet it must be admitted that closely placed sutures of this character in pathologic states of the bowel are often defective, because of pressure necrosis. The method is slow of accomplishment, also requiring great skill in placing the sutures through the submucosa without penetrating the mucosa. Should the mucosa be penetrated, a new danger is added, that of capillary oozing. By this method a diaphragm is formed within the lumen of the bowel, lessening to some extent its caliber. All of the methods

*Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Surgery and Anatomy, and approved for publication by the Executive Committee: Drs. H. O. Walker, A. J. Ochsner and DeForest Willard.

in vogue which adhere to this principle produce a diaphragm within the lumen of the bowel, and some of the late methods have added to the danger of pressure necrosis, and have contributed a new danger in placing a foreign body within the lumen of the bowel, an occasional source of secondary leakage from necrosis, ileus, perforation and peritonitis.

Some of these later methods, however, possess the great merit of being time-saving measures, which must be recognized as extremely important in cases demanding resection of the intestine. Chaput, in 1889, developed an operation termed circular enterorrhaphy by abrasion and application, in which he coapts denuded mucous surfaces, demonstrating that permanent union may be thus obtained. Robinson, in 1891, in a paper entitled "Circular Enterorrhaphy by a New Method," demonstrated union between serosa and denuded mucous membrane. Robinson's method, as well as Chaput's, does not produce a diaphragm within the lumen of the bowel. Harris, in 1892, presented a method involving the same principles as Robinson's method, with, however, an improved technic, in that he does not use a

Fig. 1.—The mesenteric arteries are ligated, and a knife is passed through the mesentery and made to encircle the bowel, cutting through serosa and muscularis.

foreign body within the lumen of the bowel. Harris has also, besides extensive experiments on the dog, applied his method to the human intestine, demonstrating sero-fibrinous union. The requirements to be met in an ideal method are:

1. The union of the severed ends of the intestine by a method which provides against immediate and secondary leakage from the intestine.
2. The method must be applicable to pathologic conditions, as well as to the normal intestine.
3. Lessening of the caliber of the bowel at the site of union should be obviated.
4. Rapidity of execution.

I wish to present a method which, in my opinion, meets all of these requirements. The character of union obtained is serofibrinous.

DESCRIPTION OF METHOD.

The intestine is drawn out through the abdominal incision and its course determined. The contents of the intestine are stripped downwards and tapes or clamps applied. The mesenteric arteries are ligated. A knife

is passed through the mesentery and made to encircle the bowel, cutting through serosa and muscularis. A cuff of serosa and muscularis five centimeters in length with the mesentery, is rolled back by means of gauze dissection on the distal portion of the bowel, which is well protected with gauze. The bowel thus denuded is severed at its distal end within one centimeter of the cuff. At the proximal end it is severed flush with the primary incision. Eight through-and-through sutures, the knots placed within the lumen of the bowel, unite the proximal end of the bowel to the denuded stump at the distal end.

The cuff is now rolled over the site of union on to the proximal end of the bowel and made secure by a suture placed through the mesentery close to the bowel and three interrupted sutures placed in the cuff through serosa and muscularis of the underlying bowel.

PRINCIPLES INVOLVED IN THE METHOD.

The cuff completely covers the sutures placed through the proximal end of the intestine and the stump of the distal end, so that leakage can not occur, as plastic lymph and blood agglutinate the cuff to the serosa of the proximal end of the intestine.

Fig. 2.—A cuff of serosa and muscularis with the mesentery is rolled back on the distal end of the bowel, by means of gauze dissection. The denuded area is severed at the distal end within one centimeter of the cuff, at the proximal end flush with the primary incision.

There can be no increase of tension on the sutures placed through the proximal end of the intestine and the stump, on the resumption of peristalsis, because of the rapid union between the cuff and the serosa of the proximal end of the intestine. The caliber of the lumen of the bowel at the point of union is unchanged.

ARGUMENTS IN FAVOR OF THE METHOD.

1. It is practical, easy of accomplishment and quick of execution.
2. It is the most natural way of uniting the severed ends of the intestine.
3. The cuff so covers and protects the sutures placed through the proximal end of the intestine and the stump of the distal end that the direct line of leakage and infection is obviated.
4. Pressure necrosis and leakage is not probable, and union is quite certain, as the blood supply to the cuff and the proximal end of the bowel that is covered with the cuff is undisturbed.

5. It is the most rapid of all methods, requiring but from five to eight minutes for its complete accomplishment.

6. It is the least complicated of any of the different methods now in use, and it can be safely performed by the ordinary surgeon, who rarely is called on to resect an intestine, quite as well as one greatly skilled in intestinal surgery.

7. Convalescence is more rapid and fewer post-operative sequelæ are likely to occur.

8. It is applicable to any part of the intestinal tract.

9. The security against primary and secondary leakage, furnished by the cuff, makes it the safest of all methods.

RESULTS OF EXPERIMENTS ON DOGS.

I will now submit a report of the pathologic findings following operation in specimens of union secured from dogs. There were four specimens examined, viz., two days, four days, eight days and thirty days, following the primary operations. The following is a report from the Columbus Memorial Laboratory, Chicago, by Prof. W. A. Evans:

SPECIMEN 1 (The two days' specimen).—This is a piece of small intestine, with mesentery attached. It is $5\frac{1}{4}$ cm. long.

Fig. 3.—Eight through-and-through sutures are placed through the proximal end of the bowel and the denuded stump of the distal end.

Near the center of the specimen is a discolored area $2\frac{3}{4}$ cm. long. Four circular incisions were made, one just above the discolored area, two through it and one just below it. A circular piece showing the convexity of the bowel was removed from just about the middle of the discolored area. At this point there is about $\frac{1}{2}$ centimeter between the outer and the inner circular muscle coats. A second longitudinal section was taken through the line of juncture between the red and white at its proximal end.

SECTION A.—This is through the farther end of the cuff. Our sections are made lengthwise of the intestines. By microscopic examination we find the following: The mucous coat shows villi glands and other mucosa structures normal in every particular. The muscularis mucosa is normal. The submucosa is normal. It is rather thick just at the point of examination. This coat is quite thick, owing to it being the seat of a mucous fold. The circular muscle coat shows slight infiltration with leucocytes. The longitudinal muscle shows the same as the circular. Outside of the longitudinal muscle is a thin zone of mixed connective tissue fibers, fibrin, red and white blood cells. It is the union between the inner muscle tunic of the cuff and the serosa. It is as firm as it should be. This layer is about $1/10$ as thick as the longitudinal muscle. Cuff circular coat is somewhat more inflamed than is the inner. Cuff longitudinal

coat is somewhat more inflamed than the inner longitudinal. Cuff serosa does not show very well in sections.

CONCLUSIONS.—Union is good between the cuff and the underlying intestinal wall.

SECTION B.—This is a section made from near where the mucosa and the submucosa of one side was stitched to the wall of the other side. Mucosa: This coat shows villi glands and other mucosa structures. It is quite normal. At four points just on the mucosa side of the muscularis mucosa there is a moderate inflammatory reaction around the thread fibers. There are some leucocytes among those fibers. The muscularis mucosa is in two coats. In the submucosa there is moderate inflammatory reaction, especially along the vessel radicles. One thread is partly in this coat and partly in the mucosa.

INNER CIRCULAR MUSCLE.—This muscle shows one thread puncture just about its juncture with the longitudinal muscle. This layer shows a fair amount of inflammatory reaction along the vessel radicles.

LONGITUDINAL MUSCLE.—This shows about the same amount of inflammatory reaction as the circular coat. Outside the longitudinal muscle is the layer of fibrinous adhesions. This layer is four millimeters thick as it appears on the slide. It is composed of fibrin infiltrated with leucocytes and larger cells with considerable perinuclear protoplasm. In this district the thread shows at several places.

CUFF.—Circular muscle fibers show considerable inflammatory reaction and extravasation of blood. The cuff longi-

Fig. 4.—The cuff is now rolled over the site of union on to the proximal end of the bowel. A suture is placed through both folds of the mesentery of the cuff close to the bowel above the anastomosing branch of the mesenteric artery, and through both folds of the mesentery of the proximal end of the bowel. Three interrupted sutures are placed through the cuff into the serosa of the proximal bowel.

tudinal muscle shows much extravasation of blood and a fair amount of inflammatory reaction. The serosa, except for extravasated blood, is negative.

SPECIMEN 2.—This is a piece of small intestine with partial mesentery attached. It is 8 cm. long. Length of the discolored area in this specimen is 2 cm. The incisions were made just above the discolored area, $4/10$ cm. below its upper border, 1 cm. lower down, and three other incisions each $5/10$ centimeters apart.

SECTION A.—This extends longitudinally through the intestine at the tip of the cuff. The sections here show the same as in the section marked "Section A, Specimen 1." The photograph shows a very good union, still fibrinous, between the cuff and the underlying intestine. The other layers are about as reported in Specimen 1.

SECTION B.—This is a semicircular section of the convex half of the intestine near the point of union of the mucosa and submucosa, to the mucosa and submucosa of the unstripped end. It is mucosa. The epithelial elements of the mucosa are perfect. In the connective tissue structure of the mucosa is more inflammatory reaction than in

any section yet examined. The muscularis mucosa is in two coats, the longitudinal coat being the heavy one. The submucosa shows considerable inflammatory reaction. The circular muscle is filled with cells of inflammation, the fibers being considerably displaced. The longitudinal muscle shows the same as does the circular. The sections are evidently beyond the point of suture, i. e., in a region not split for the flap.

SECTION C.—These sections are just up the cuff from the line of suture of the mucosa. They show everything shown in "B" and in addition, external to the longitudinal muscle coat, is a coat of fibrous adhesion, and outside of this are the two muscle layers of the cuff.

SPECIMEN 3.—A piece of larger intestine than in the other specimens. The length of the piece is seventeen centimeters. The length of the discolored area is two centimeters. The bowel has been cut longitudinally just above the proximal end about 0.3 cm. below it, 0.5 cm. lower down, 1 cm. lower down, the last being outside the line of discoloration.

Sections were made as follows: A, longitudinal section through the cuff. B, the convexity of the bowel about the point of union of mucosa. B¹ near B.

Fig. 5.—a, Mucosa. b, Muscularis mucosa. c, Thread.

SECTION A.—This is a longitudinal section near the tip of the cuff. The mucosa, muscularis mucosa, submucosa and longitudinal muscle are quite normal, showing very little inflammatory reaction. The line of union between the longitudinal muscle and the circular muscle of the cuff shows the following: The space covered is about 50 mikromillimeters. It is filled in by a loose reticular fibrous tissue. The fibers have no uniformity of direction. There are only a moderate number of blood vessels. These are young. There are many cells around the embryonal vessels. The vessels that I identify as such run lengthwise of the intestine. The two muscle coats of the cuff show about as much inflammation as the muscle coats of the intestinal wall. The serosa is thickened by inflammatory reaction until it is about as thick as one of the muscle coats.

SECTION B.—Longitudinal through the point where submucosa and mucosa of one side are joined to the intestinal wall of the other. The mucosa is perfect except for leucocytic infiltration. There is a deep V-shaped dip at the point of union, the apex being toward the muscle. Around the apex of the V runs the muscularis mucosa. The submucosa is also continuous. It is thin at the apex of the V. The circular muscle of the side is continuous and into the other. This is true of the longitudinal

muscle and the serosa. The muscle coats of the other side come into the inverted V and turn up to form one of its sides.

They are joined to the circular muscle of the other side by a small amount of fibrous tissue. (The threads are knotted in the lumen and are firmly held in the wall.) All of the coats contain large numbers of leucocytes.

SECTION B.¹—Circular beyond the line of sutures. This shows as indicated for Specimen 2, except that there is greater leucocytic infiltration in each of the coats.

SPECIMEN 4.—This was a piece of intestine, split longitudinally. Microscopically we do not find any evidence of the operation procedure, though this is probably due to our not finding the proper area for section. There is a heavy wall of the usual coats. One half or more of the specimen has a serous coat that shows some peritoneal thickening. The other portion is not so covered.

EXPERIMENTS ON DOGS.

I have performed 75 primary operations on dogs, with the following results: 72 dogs recovered and 3 died, a mortality of 4 per cent. The convalescence in every case was short, the dogs being in every instance turned into the general ward at the end of forty-eight hours. The following is a report of the cases that died:

Fig. 6.—a, Cuff. b, Line of union.

CASE 1.—Dog No. 36 died five days following operation. Cause of death was general peritonitis. The great omentum was infiltrated with blood and pus, and was adherent to the cuff. Free pus was found in the peritoneal cavity. The intestines were not much distended, but were agglutinated together in several places. The site of operation showed the cuff rather dark, but union of the cuff to the underlying bowel wall macroscopically seemed good. A section of intestine involving the site of union was removed, and examined macroscopically, but no evidence of non-union was discovered. The intestine was gently distended with water, but no leakage was found. The cause of death was believed to be an infection from without, due to a faulty technic.

CASE 2.—Dog number 48 died four days following operation. The cuff was distended with blood and pus, a perforation was found in the cuff in the line of union of the proximal end of the bowel to the stump, with positive evidence of leakage.

CASE 3.—Dog No. 62, died thirty-six hours following operation. No evidence of peritonitis. The appearance of cuff was good, and was seemingly well adhered to the underlying bowel wall. On section the intestines were found quite well filled with blood, showing that death was due to hemorrhage.

Thirty dogs were reoperated on at periods varying from two to thirty days following the primary operation, with two deaths, a mortality of 7.5 per cent. The deaths occurring in dogs operated on at the end of forty-eight hours following the primary operation, and in each instance within twenty-four hours.

CONDITIONS OF THE CUFF FOUND.

In nearly every instance there were slight adhesions of the omentum to the cuff. The union of the cuff to the underlying bowel was good in every instance. In one instance the intestine above and below the cuff for a distance of five or six inches was adherent, making a bend in the intestine just below the cuff, but in no way obstructing the lumen of the bowel. In every respect the results were universally good.

WORK ON THE HUMAN.

I have only used this method in two cases, in the one a somewhat modified technic was followed, in the other the same technic, as above described, was adhered to.

CASE 1.—Mr. B., 50 years of age, was operated on Feb. 6, 1903, for the removal of a tumor in the region of the sigmoid flexure of the colon. The tumor, with several inches of the intestine, was removed. The proximal end of the intestine, which was to be covered with the cuff, was denuded of its serosa, so that the cuff came in direct contact with the muscularis. The patient made an uninterrupted recovery. The tumor removed proved to be an adenocarcinoma.

CASE 2.—Miss B., aged 20, had been operated on by the writer five months previously for the relief of chronic obstruction of the bowels, due to tubercular peritonitis. The patient was relieved of the obstruction, improved in general health, and it seemed for a time that her ultimate recovery was assured. Feb. 1, 1903, about five months following operation, she again developed pronounced symptoms of intestinal obstruction. Her condition grew worse from day to day, although slight bowel movements were secured by the use of the high rectal tube. February 12 she had stercoraceous vomiting, and a fecal discharge made its appearance in an abdominal fistula remaining from the previous operation.

Although the condition of the patient could hardly be more extreme, and but little hope could be given her of an operation benefiting her, yet she was desirous of the few chances that it might offer. The abdomen was opened in the median line. The intestines were greatly distended and very much adherent to the parietal peritoneum, and greatly matted together. The site of obstruction was searched for and found. The fecal fistula was located, the intestine just above the site of obstruction was perforated in two different places, so that it was necessary to resect about six inches of it, as it was impossible to attempt its repair. There was no difficulty in obtaining the cuff, the above described method being practiced in every detail. The time occupied for the complete resection was ten minutes. In the further liberating of the adherent intestines a large tear was discovered in a badly ulcerated area, and a second resection of probably four inches of intestine made.

The second resection was a little more difficult on account of adhesions and the extensive pathology in the mesentery; however, it was accomplished in a reasonably short period of time. The patient died at the end of thirty-six hours. A postmortem was denied, so that we can not say as to the behavior of the parts of the bowel united.

The method of operating had nothing to do with the results in this case, as it was evidently hopeless by any method of procedure. It has, however, demonstrated that a cuff can be readily formed of pathologic tissue and that the method is applicable in all cases where resection of the intestine is necessary.

REFERENCES.

Ovariectomy and Abdominal Surgery, by Harrison Cripps, F.R.C.S., Chicago Medical Recorder, September, 1892.
Transactions of the Western Surgical and Gynecological Association, 1900.

Abdominal Surgery, Greig Smith, vol. II.
Chicago Medical Recorder, vol. XII.
Senn's Experimental Surgery, 1887.
Annals of Surgery, 1891, xliii, 427.
Treves' Operative Surgery.

INTESTINAL RESECTION,

WITH A REPORT OF SIXTEEN CASES: NINE WITH THE MURPHY BUTTON AND SEVEN WITH THE SUTURE.*

JAMES H. DUNN, M.D.

MINNEAPOLIS.

With the exception of small lesions toward the free border of the tube, surgical repair of the intestines usually implies a resection more or less extensive, hence the title intestinal resection has seemed more nearly descriptive of the matter on which it is my purpose to elicit discussion than the more usual term "intestinal anastomosis."

Every surgeon who undertakes even cases of strangulated hernia, intestinal obstruction or penetrating wounds of the abdomen, as well as all who open the abdomen for whatever cause, must be prepared to face, often in the most unexpected manner, a resection or repair of the intestinal canal. In the vast majority of cases intestinal surgery is decidedly emergent. The urgent nature and comparative rarity of such operations in any one surgeon's practice renders a perfect technic exceedingly difficult. To skilfully execute such an operation on a moment's notice under the most varying conditions, and perhaps at rare intervals, is a severe task. These considerations lend the subject a somewhat general interest, and place it among those topics which may be frequently discussed with profit even though nothing new be brought out, every surgeon having reason to frequently review his knowledge and rehearse his technic.

The first difficulty one meets is the great multiplicity of methods and the almost endless novelties of technics and operations from which to select. With the mind so full of many theories, and the preparatory drill spread over endless ways, the operator is in the worst possible condition to meet an emergency. Have we not arrived at a point where we can stop investigating new frills and settle on the simplest procedures with advantage?

During the past six years I have had occasion to resect the intestine 16 times, as follows: For strangulated hernia, 3; for fecal fistula, 3; for sarcoma of the small intestine, 2; for carcinoma of the cecum, 2; for carcinoma of the sigmoid, 2; for intussusception, 1; for gunshot wound, 1; for injury during operation, 2. In these the Murphy button was used 9 times, with 3 deaths; the suture 7 times, with 1 death.

The somewhat prodigious number of methods for uniting the bowel may be first divided into three general classes: First, mechanical appliances, to remain in the intestine; second, pure suture methods; third, special appliances to facilitate suture, and then to be removed.

The latter class, to say the least, belongs to those things which are more ingenious than necessary. For my part I do not use them, for many reasons, one of which is perhaps quite sufficient, viz., as good work may be done more conveniently without as with them.

Mechanical means versus the suture in intestinal anastomosis is often debated pro and con. As a simple general proposition, this is manifestly unjust. Possibly all anastomoses may be made by the suture, but cer-

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Surgery and Anatomy, and approved for publication by the Executive Committee: Drs. H. O. Walker, A. J. Ochsner and DeForest Willard.