3. Patients lying in bed should have the diseased joints exposed to the direct rays of the sun, their heads and eyes being protected by green glasses or shades. The joints may be covered with blue, so as to secure easiest passage of the ultra-violet actinic rays, and local medications rich in iodine may be also employed as desired.

4. Tent life on the hospital grounds, or better, in the open pine forest, can be successfully employed through both summer and winter.

5. Sanatoria should be established for tuberculosis of the hard tissues, as well as of the soft.

6. The concentration of the sun's rays by lenses, as suggested by Butler, Finsen and others, is of positive benefit in bactericidal influence. As final curative agents, however, the direct sun's rays are most effective. Electrical rays can be used when sunlight is lacking.

7. The Roentgen rays in the laboratory have an inhibitory power on the tubercle bacilli, and may prove useful in restraining the growth of these micro-organisms in living tissues.

8. The actinic rays and the x-rays are both apparently helpful in the fight with tuberculosis, but several years will be required to determine accurately their effect. They should be employed not to supercede, but to antedate and to supplement operative procedures, to assist the mechanical protection of the joint, and to increase the general therapeutic measures employed.

BIBLIOGRAPHY.

1. Trans. Amer. Orth. Asso., xv, 1902, p. 10.
2. Annales de Chirurgie et d'Orthopédie, vol. xiii, 1900, p. 19.
3. Lancet, May, 1901, 1396.
4. THE JOURNAL. Amer. Med. Asso., April 11, 1903, p. 966.
5. Allgem. Wiener Med. Zeitung, 1902, p. 26 (Kattenbracker).
6. Recherches sur l'inflammation provoquée par l'action de la lumière solaire. N. B. Finsen.
7. Pleasanton. The Influence of the Blue Ray of Sunlight, and the Blue Color of the Sky in Arresting Disease. 1867.
8. Jour. Advanced Ther., May, 1902.
9. Auerbach: Centralblatt für die Med. Wissenschaft, xix, p. 1.
10. H. Reider: Münchener Med. Wochenschrift, March 11, 1902.
11. Milton Franklin: N. Y. Med. Record, Oct. 25, 1902, p. 652.
12. Finsen: Phototherapy.
13. Finsen: "Ueber die Anwendung von Concentrirten Chemischen Lichtstrahlen in der Medizin." Leipzig (Vogel), October, 1899, p. 52.
14. Williams; R. R.: In Med. and Surg. McMillan & Co., 1902.
15. Hopkins, G. G.: Phil. Med. Jour., Oct. 7, 1899; Dec. 9, 1899; Oct. 27, 1900; Sept. 7, 1901.
16. Downs and Blout: Effect of Light on Bacteria. Proc. Royal Soc. Lon., vol. 26, 1877, S. 488.
17. Duclaux: L'influence de la lumière du soleil sur la vitalité des micrococci. August, 1885.

A STUDY OF THE HISTOLOGIC CHANGES IN EPITHELIOMA UNDER THE X-RAY.

A PRELIMINARY REPORT.*

J. CLARK STEWART, M.D.

Professor of Principles of Surgery, Medical Department University of Minnesota.

MINNEAPOLIS.

The striking and unforeseen curative effects of the Roentgen ray on epitheliomata make it probable that a careful study of the histologic changes occurring in these tumors while under treatment must prove both interesting and instructive.

Medical literature is rather barren in studies of this sort, most opinions put forth being founded on insufficiency or absence of observed facts. Note the opinion solemnly set forth in a journal devoted to the exploitation of x-ray work: "The beneficial influence which the agent (x-ray) undeniably exercises on superficial cancer is due to x-ray vibrations acting on the cell elements of the growth, which elements are too weak and lack-

ing in vitality to reach maturity and become the specialized tissue for which they were destined of themselves in such a way as to give them new vitality, and hence ability to reach maturity and their normal ultimate structure."

That is, the x-ray is a sort of reformatory treatment by which the erring cancer cells are spanked and brought to a realizing sense of their vagrant ways, become reformed and proceed to grow into normal adult tissue, a very beautiful effort of the imagination, but needing a substratum of facts.

Many equally remarkable explanations of the effect could be quoted, but this one will suffice.

Dr. A. G. Ellis has made an important contribution to this subject from a strictly scientific standpoint in the *American Journal of Medical Sciences*, January, 1903. He studied four cases, two of which were epitheliomata which had been subjected to varying amounts of exposure to x-rays, and has determined some interesting facts. If he gives the literature at length, it will not be necessary to quote authorities.

His conclusions are that the study of these cases shows: 1. Necrosis of cells and trabeculae of varying

Fig. 1.—Photograph of Case 1 before treatment.

degree. In Case 1 (a cancer of the breast) there was marked fatty degeneration. 2. Increase of elastic tissue. 3. Fewer areas of lymphocytic infiltration in one case after exposure, about equal numbers in others. 4. A tendency to occlusion of vessels by deposits on their inner surfaces. 5. Practically entire absence of infiltration by polymorphonuclear leucocytes. He says: "The unsatisfactory clinical results, as well as the microscopic changes in Case 3 (epithelioma) can probably be safely attributed to the presence of the exceedingly numerous keratinized areas or 'pearls.'"

The results of the present study are quite at variance with the above in many respects, as will be seen later. The larger part of the material studied was obtained from one case, which came under observation December, 1902, and the research work was begun December of some time before the publication of Dr. Ellis' paper.

CASE 1.—A. B., Norwegian, aged 52, single, gave the following history: Eighteen months ago he cut the back of his right hand with a beer bottle, and the cut became infected and healed slowly with a dense raised scar. Soon after an elevated wart-like growth appeared in the scar, which was repeatedly cauterized with pure carbolic acid without much effect. His general health has been good. There is no cancer in his ancestry.

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Surgery and Anatomy, and approved for publication by the Executive Committee: Drs. A. J. Ochsner, J. E. Moore and De Forest Willard.

or environment. An examination shows a fungus ulcer about three inches in diameter on the back of the right wrist. It is raised above the skin more than one-half an inch, the elevation being greatest at radial edge. The extensor tendons of the fingers move freely, but just at their radial side the growth

Fig. 2.—Fatty degeneration in epithelial pearl. Hermann fixative 500 diameter.

has deep attachments. There is no enlargement of the axillary lymph nodes. In view of the impossibility of excision it was decided to subject the growth to x-ray treatment, as it seemed the only chance of saving the hand.

On December 10, 11 and 12 he was given twelve minutes' exposure to a one-inch tube at five inches. On

Fig. 3.—New blood vessel containing red blood cells in center of epithelial pearl.

December 16 and 17 he was given fifteen minutes at five inches. On December 18 the first block was excised, being cut from the non-sensitive surface of the ulcer, half an inch from the skin edge. December 22 and 23 he was given twenty-five minutes' exposure to a low tube

at five inches. December 23 block excised. December 29 block excised. From December 29 to January 2 he was given daily twenty minutes' exposure to a low tube at five inches, and blocks were removed every other day until the whole fungus part of tumor had been softened and came away. In all, seven blocks were secured, covering the time from the fifth exposure up to the eighteenth, in all, thirty-six days.

HISTOLOGIC CHANGES NOTED.

Each block was cut in three portions and fixed in different fixatives. Hermann, Mann, Zenker and 25 per cent. formalin being used. The blocks were carried up, embedded in paraffin and cut in sections of from three to ten microns in thickness. The sections were examined unstained from the Hermann fixative for fat, and others were stained in various ways. The Hermann sections were washed in ether to differentiate fat from other substances taking osmic acid. Hematoxylin and eosin, hematoxylin and Van Gieson, Weigert's resorcin stain for elastic fibers, and Pianese IIIb for degenerations, Plimmer's stain for parasites were used. By these methods quite a number of changes could be made out, which can be briefly tabulated as follows:

Fig. 4.—Circle of new vessels in epithelial pearl.

1. *Fatty Degeneration.*—This is marked in all sections from all blocks. In the early sections it is most marked in the epithelial cells at the periphery of the pearls, the fat occurring as granules grouped about the nucleus, and in many cases in the nucleus itself. Fat exists, however, in a large percentage of the epithelial cells, and this infiltration seems to increase with the number of exposures to the x-ray, until the cells disintegrate, when in the final stage of detritus it is seen mainly in the leucocytes, either as a degeneration, or possibly as a phagocytic inclusion.

In view of Unna's statement that fatty degeneration does not occur in epitheliomata, this is certainly a noteworthy effect of the treatment.

2. *Vascular Changes.*—In the earliest slides there are marked evidences of the formation of new blood vessels. They are seen between the epithelial cells and noticeably in the centers of many pearls. They are at first filled with red blood cells with an occasional white cell, but later the red cells disappear and the vessels are packed tightly with the polymorphonuclear leucocytes. These leucocytes are also seen between the epithelial cells, and

finally separate them into bands and detached cells surrounded by a reticulum filled by white cells.

3. *Elastic Tissue*.—Weigert's resorcin stain shows considerable elastic tissue in fibers and granules, usually in the immediate neighborhood of a new vessel, but not in its walls.

4. *Hyaline Degeneration*.—As this degeneration is a regular part of the life history of an epithelioma, it is not surprising that it should form a large part of the picture in many of the sections. It is especially noticeable in both pearls and single cells about the middle of the series, being replaced by coagulation necrosis in the latest slides. In many of the later slides the hyaline bodies of Unna are plentiful, as well as many cell inclusions, which, stained by the Pianese IIIb, or by Plimmer's method, are indistinguishable from the so-called Plimmer and Russell bodies as figured by Gaylord. As these do not appear in the early sections, but only when karyokinesis has ceased and degeneration of various kinds are very marked, it is difficult to under-

mented nuclei and other detritus, while interspersed through the mass are small clumps of epithelium, single cells, which have no nucleus and have lost their outline, and are just ready to pass into coagulation necrosis. In areas where the final fluid detritus was defined the microscopic appearance was like a so-called

Fig. 5.—Elastic tissue. Weigert resorcin-safranin.

stand how these fuchsin bodies can have any etiologic relation to cancer. They are indistinguishable as to staining reactions from the other hyaline bodies noted above.

5. *Necrotic Changes in Cells*.—In some of the early slides karyokinesis is well marked, but later it entirely disappears, except an occasional cell in blocks taken from near the skin border of the ulcer. Chromatolysis begins quite early and increases until in the last sections there is hardly an intact nucleus. At the same time there is fusion of the cell bodies and loss of all cell outlines. Hyaline degeneration of single cells is very common, as before stated, as are the fuchsin bodies.

6. *Final Result*.—The latest sections obtained before destruction and liquefaction show an interesting and remarkable picture. Nothing is left of the epithelial structure but some dimly staining masses of cells in coagulation necrosis, with the rest of the section composed of a fine reticulum taking the acid stain, having its spaces filled with polymorphonuclear leucocytes, frag-

Fig. 6.—Shows hyaline cell inclusion similar to Plimmer bodies.

mentable pus, and the microscope showed only polymorphonuclear leucocytes, broken-down epithelial cells, and free fat.

The study of sections from other epitheliomata, at varying exposure, gave results corresponding to the above, although in small growths it was impossible to get more than one or two blocks. One case especially, an epithelioma of the ear, after four ten-minute exposures, gave a picture like the final one from Case 1, except

Fig. 7.—Shows general breaking down with infiltration of leucocytes into and between the epithelial cells, hyaline degeneration of the cells and at outer edge coagulation necrosis of whole mass.

the leucocytic infiltration was almost entirely composed of plasma cells, instead of the polymorphonuclear cells.

It will be noticed that there are several very striking differences between the results of the study noted above and those obtained by Ellis in the two cases of epithelioma.

ma studied by him. These are the vascularization and the leucocytic infiltration which are probably only phases of the same process, and the difference was probably due to the destructive effect of the treatment in my cases, as compared with the negative clinical effects noted by him.

Certainly his explanation of the resistant character of the pearls, preventing curative effects, seems doubtful, at least, in view of the very early and decided changes in them noted above.

The explanations offered by some writers that the degenerations observed and final necrosis are due to decreased blood supply, which is caused by the thickening of the inner coats of the blood vessels, find no support in this study, as the abundant formation of new vessels and their distension makes improbable any obliteration of the different vessels. Unfortunately, none of the sections cut were from locations which could give any information as to whether the endarteritis observed by so many was present.

It seems probable that different epitheliomata may offer different degrees of resistance to the disintegrating effects of the x-ray, just as their bearers may show different degrees of susceptibility to the burning effects of the ray, but when necrosis is finally attained there will be characteristic changes in the cells.

It is to be regretted that sections were not taken from Case 1 before treatment, and more frequently during the active treatment, but the idea of making a study of the case occurred only when rapid results began to be noted after four treatments, and the subsequent progress was so rapid that all available tissue was destroyed when the seven blocks were obtained.

It may be of interest to hear the subsequent history of this case which began so auspiciously. Early in January, after some six weeks of treatment, all the elevated portion of the growth was destroyed and there was left a deep portion which extended into the bones of the carpus, and under the small tendons of the thumb toward the wrist. After wasting a great deal of time in a vain attempt to cause necrosis of the deeper parts, a small, apparently new growth was noticed at the radial edge of the ulcer. This determined an erasion under an anesthetic, which showed that the growth had progressed under the skin at the radial edge of the ulcer, in spite of vigorous x-ray treatment. All the epithelioma was carefully scraped away, and all overlying skin was excised, as well as the small extensor tendons of the thumb, which were hopelessly involved, and there is now a clean defect, with no signs of epithelioma, which is healing rapidly under mild x-ray treatment.

The tissues excised, which included nerves, tendons, vessels and new-formed scar, all exposed to many x-ray treatments, will furnish material for a later communication.

CONCLUSIONS.

1. It is probable that when epitheliomata react favorably to x-ray treatment that characteristic histologic changes will be found.
2. The important early changes are fatty degeneration and vascularization of the epithelial pearls.
3. Leucocytic infiltration and various degeneration processes complete the destruction.
4. Bodies indistinguishable from "Plimmer's bodies" multiply as epithelia degenerate.

NOTE.—I wish to acknowledge with thanks the aid in this study of Prof. Margaret S. Nickerson, Drs. Louis B. Wilson and Arthur T. Mann, without whose assistance the work would have been impossible.

DISCUSSION

ON PAPERS OF DRs. WILLARD AND STEWART.

DR. HENRY A. CHRISTIAN, Boston—I wish to add one case to the report of Dr. Stewart, a case of epithelioma of the back of the hand of the distinct cauliflower type. Before any treatment was undertaken a portion of the tumor was excised and sent to the pathologic laboratory of the Boston City Hospital for examination. It was a typical specimen of epithelioma, with the usual cell nests and with no particularly marked amount of inflammatory reaction. After treatment for a considerable length of time a second portion was excised. Pathologic examination of this showed that the epithelial cells in the cell nests were dead. They no longer stained and, what is particularly interesting, they were in many places surrounded by foreign body giant cells. In other words, the dead epithelial cells had changed from a living acting part of the growth and now simply acted as a foreign body and were being destroyed by the giant cells. There were no polymorphonuclear leucocytes present. There was no change in the connective tissue cells, while the epithelial cells were entirely dead.

Another point of interest was that a portion of the hand protected by the lead shield and supposed to be non-carcinomatous, i. e., to lie beyond the border of the growth, was in reality carcinomatous. This portion protected from the x-rays by the lead shield showed no such change. In other words, the epithelioma survived where the lead shield protected the growth; and there was death of the cells where the lead shield did not protect it.

Just what the process in these cases is I am not prepared to say. I do not think that too much importance should be placed on the fatty degeneration described by Dr. Stewart. Fatty degeneration has, so to speak, lost in recent years some of its pathologic standing. We no longer look on it as such an important degenerative process. Then, fat is frequently present in cells of the body under normal circumstances. Furthermore, the reagents used by Dr. Stewart for obtaining the reaction demonstrating fat are open to some objection. The main reagent used was osmic acid. We know that osmic acid stains other things than fat, for instance, certain keratin granules in the epithelial cells of the skin, also, the fatty substance of the sebaceous glands. Now, as the epithelioma originates from the skin these substances normally present must be borne in mind. If a substance stains black with osmic acid, this is not positive proof that it is fat, nor is the presence of fat always evidence of a degeneration.

Another interesting point is the presence of fat in the polymorphonuclear leucocytes. This occurs very frequently; in fact, it is generally present where there is an inflammatory process. It is also to be found in the polymorphonuclear leucocytes circulating in the blood. This is true particularly where there is a marked leucocytosis. I have had an opportunity to study these cells, particularly in association with pneumonia, and have found fat very constantly present in them.

Another point on which emphasis was laid is the great number of polymorphonuclear leucocytes present in this case. Is that due to the effect of the x-ray or to something else? In this case I understood that the surface skin was ulcerated. In such a condition you would expect to find polymorphonuclear leucocytes present at the outset. Again, when we have polymorphonuclear leucocytes and an inflammatory process we generally have the formation of granulation tissue; in other words, the formation of new blood vessels, so that the statement that the formation of new blood vessels is caused by the x-ray is possibly also open to criticism. I think the fact that the epithelial cells are dead, as shown by the case reported by Dr. Stewart and the case I have reported, are rather important factors. It is possible that simple inflammation may have had something to do with the formation of the polymorphonuclear leucocytes and the new formed blood vessels. The question remains open as to how far fatty degeneration, inflammation and cell death are due directly to x-ray action. Very general conclusions are not, of course, to be drawn from single cases.

DR. EDWARD H. NICHOLS, Boston—I agree with the beginning of Dr. Stewart's paper, but there are one or two things later on which I am compelled to differ with him. In regard to those lesions which were selected for this *x-ray* treatment as epidermoid cancer or epithelioma, which, after a greater or lesser number of treatments, have disappeared, the ulcers healed over, the capillary growth melted down and entirely disappeared and new skin formed, such cases are constantly being reported as cured. I think we should consider all those statements of cures with the greatest suspicion. It is true that for some reason or other the surface condition has apparently improved. The fact that the new skin has formed over the area of the wound is far from proving a cure of that case of cancer. I have had considerable opportunity during the last three years of observing the histologic changes following *x-ray* work. The cases referred to by Dr. Christian I have had the opportunity of seeing. The changes as we have seen them following the use of the *x-ray* have shown no peculiar characteristics. The changes in cases of epidermoid cancer treated by the *x-ray* show, as a rule, an increase, and in many cases stimulation of the cells. And I would defy anybody to recognize this change as a characteristic one. You see exactly the same changes as you would see in cases of degenerative carcinoma, and this has been deemed by some men to be due to the action of the *x-ray* on certain portions of the epidermoid growth, but they are really indications of the growth of the cancer. The *x-ray* acts as an irritant, and acts on those portions of the cancerous growth which are less resistant than the sound tissue. I think it can be perfectly explained by saying that the *x-ray* acts as a mild irritant, not producing burns, yet sufficient to cause increase in the epithelial cells at first and later disintegration. One other thing that needs to be borne in mind is that the *x-ray* can also produce epidermoid cancer. At a meeting of the American Dermatological Association a year ago last August, a case was reported of extensive epidermoid cancer of the hands of a man who had been using the *x-ray*, followed rapidly by metastasis. There is at the present time an operator in Boston who has lost three fingers on one hand and two on the other due to extensive epidermoid following rapidly after very extensive *x-ray* irritation, he having had a distinct *x-ray* burn. In regard to the bodies that Dr. Stewart mentioned, I have seen their work in their own laboratories, so perhaps I can express an opinion, and I have seen the definite Plimmer bodies of Greenough, and I do not think these objects seen by Dr. Stewart were the Plimmer bodies. I think every surgeon should bear in mind very carefully the fact that the *x-ray* is liable to produce cancer. In certain cases it may do good, and for that reason I think it is better to first remove the epidermoid cancer and then use the *x-ray*, and in certain cases it may be of advantage, but I think in all cases operative treatment should be the first procedure.

DR. JOHN RIDLON, Chicago—In regard to the treatment of tuberculosis of the joints by sunlight, I do not think that we should assume too large a hope in this direction. Sunlight or any other light alone does not cure tubercular joints; they must be treated by mechanical means, immobilized, or else no good can result from the exposure to the sunlight or any other light. As to the effect of sunlight on tuberculosis of the joints, I am somewhat in doubt as to whether it is likely of itself to produce beneficial results. All of us who practice in large cities and large towns can see great gain if a child is exposed to the sunlight. On the other hand, in the case of tubercular joints in the children of farmers and people who live in the country, who are more exposed in the sunlight and fresh air, there is usually not so much improvement from further exposure to sunlight and open air. It would seem then that from clinical observation it is the change of environment that has the effect rather than the sunlight and air without the local environment.

I have had no personal experience with the *x-ray* treatment of tuberculosis. My office associates, Dr. A. D. Harmer and Dr. John L. Porter, have done considerable experimental work with the *x-ray* on tuberculosis of the joints, and such

cases as we three together had that seemed to be favorable experimental work we have subjected to that treatment. Cases were those in which the progress had been unfavorable under ordinary mechanical fixation. It was thought only that such cases should be selected and not those which doing well under ordinary treatment. A considerable number have been subjected, during the last year and a half, to *x-ray* treatment. Many have improved more rapidly than have been expected without this treatment. The number, however, has not been sufficiently great for us to form any worthy conclusions. It has reminded me very much of iodoform injection treatment, which was very much in vogue about ten years ago. Almost all of us used these injections of iodoform in the cases of tuberculosis, and we all thought we were getting better results. We all know now that injection of iodoform into tuberculosis of the joints is practically useless. So I fear that this *x-ray* work will turn out in the same way, and that, although a certain number of cases do appear to be cured, it will not amount to much. What we have learned is that when the superficial joints are benefited by the *x-ray* there must be very much irritation of the skin. Tuberculosis of the hip joint and the spine have not been benefited. Dr. Harmer has deliberately burned the skin in many cases, and he has found he does not get as good results when the skin is not subjected to strong irritation.

DR. FRED. B. LUND, Boston—I have seen, during the last five years, cases of cancer treated by the *x-ray*. Surgeons recognize that epithelioma of the face, particularly of the rodent ulcer type, is a non-malignant form, that recurrence is slow, and that in the cases that have recurred there has usually been no glandular involvement. I have seen a great many cases of cancer of the face, many of the rodent ulcer type, that were really benefited by the *x-ray* treatment, and leave a most excellent scar, and some of our surgeons in Boston have stated (very prematurely, it seems to me) the day of the knife in the treatment of epithelioma is past, and that we had no business to use the knife in these cases, but that they should be turned over to the *x-ray* treatment. On the other hand, I have seen some very sad results following that advice. A patient came in suffering with epithelioma of the lip in a very early stage. Glandular involvement appeared soon after it was exposed to the *x-ray*, and he was treated by the *x-ray* without the slightest result. I have seen a number of cases of recurrent carcinoma in breast glands treated without the slightest result. There is a certain number of cases of epithelioma of the face, of a more malignant type, in which it is of no value. I operated three years ago on an old lady 68 years of age. The cancer was excised and the wound covered with skin grafts; excellent results followed for a time. Then recurrence took place and I excised the growth under local anesthesia. After operating on her 12 or 15 times, I tried the *x-ray* treatment six months. At the end of this time the growth recurred with extraordinary rapidity, and the soft parts of the face had to be excised down to the nasal bones. In other words it seems to me that the *x-ray* is in the nature of a mild irritant or caustic, that it has an excellent influence on the malignant growths, but that the malignant growths had to be excised. The ideal way perhaps is to first excise the growth and then in case of recurrence use the *x-ray*, but temporary relief with the *x-ray* in the first instance is not only useless but dangerous.

DR. JOHN W. BRANDAU, Clarksville, Tenn.—Is there any effect produced on the kidneys by the use of the *x-ray* in cancer of the cervix? In the past few months I have treated two cases of carcinoma of the cervix. One was of the flower type, the other hard cancer. I had excellent results in both so far as the cancer was concerned. The cauliflower cancer underwent rapid degeneration, and after ten treatments of fifteen minutes each every second day the cervix was practically normal in size. In the other case the cervix was very different. There was sloughing of the cervix and seventeen treatments covering a period of three months

both cases the pain was extreme previous to treatments; opium in some form being administered. After the second exposure there was no more need for opium. One of the women went to her home apparently cured of the cancer to await the results of the sloughing, with instructions to return in the course of three or four weeks for examination. In about four weeks there was suddenly almost complete suppression of urine, which was highly laden with albumin. In a few days she died from uremia poisoning. The other case died about eight weeks after treatment was discontinued from exactly the same cause, and what little urine I could get by catheterization was almost solidified by albumin on boiling. I would like to know whether any of the other gentlemen present have observed any bad effects on the kidneys from this treatment.

Dr. J. F. THOMPSON, Galveston, Tex.—A few months ago a young man 29 years of age, with a large tumor over the stomach, was operated on, and it was found that the entire anterior part of the stomach and the lesser curvature was involved with a growth that afterward proved to be an epithelioma. There was almost complete obstruction of the pylorus for several weeks before the operation. I did a posterior gastroenterostomy and got immediate relief of the symptoms of obstruction. The growth appeared to be an extensive one, and about three days after the operation it occurred to us that it might be well to try the effect of the *x*-ray on this tumor. About forty-eight hours after the operation, I believe, we reopened the wound and packed the opening between the peritoneal parieties and the tumor, leaving this mass, which was about the size of a large orange, in the wound. We got profuse exudate. After 24 hours it ceased. We took out our own packing and found that the mass had been transformed into an external one. We then commenced the use of the *x*-ray, and after having used it for about a week, we noticed signs of increase in size, and a few days later the mass broke down and discharged profusely a cheesy matter mixed with pus. It then became apparent to us that we were going to have a fistula, because sloughing was extensive, and a few hours later, the nurse came to me and informed me that there was something in Mr. A.'s bed that looked like his breakfast. We found this to be true. A large opening was found in the stomach. We succeeded in packing this off, so that the patient could retain food sufficiently long to pass through our very large anastomosis into the bowel. He went to bed, was comfortable, took his nourishment well and enjoyed it. He seemed to have an especial liking for such indigestible foods as ham and bacon. He improved slightly in weight for two or three weeks, when he began to go down, and after about a month died from extreme emaciation. This, to my mind, seems to teach us some very interesting points. One is that we can not break down hard epithelioma with the *x*-ray. The second is that it is rather dangerous to try to break down a cancerous growth in the abdominal cavity. Third, the *x*-ray did not appear to have any prominent effect in checking the general growth of the epithelioma, which had become metastatic at the time of the operation.

Dr. DE FOREST WILLARD, Philadelphia.—In answer to the question as to whether a blue medium is of value either with the *x*-ray or with the Finsen light, or with sunlight, it seems to me to depend entirely on the power of the medium to increase the force of the ultra-violet rays. Since the ultra-violet rays are therapeutically the potent ones, of course, the larger the proportion the better will be the result. The rays have proven themselves of value already in superficial tuberculosis and cancer, but we have by no means achieved the limit of possibility in the use of the sun's rays. I stated in my paper that we should use these agents as supplementary to our general treatment and not depend on them entirely. In regard to the use of the *x*-ray in tuberculosis, the benefit has certainly been sufficient to encourage me to use it further. Its influence in cancer and epithelioma has in my hand been helpful, but not curative. The Finsen light and the *x*-ray are still only in the experimental stage, and I have no anticipation that they will do more than act as accessories to general treatment. I

expect to test them for five years. I do not expect to obtain any results that will be of value in a shorter time.

In reply to Dr. Ridlon, I would say that in my conclusions, I stated that sunlight and fresh air should be employed, "together with fixation of the joints." In another conclusion I said that this method should be employed, "not to supersede, but to supplement the mechanical and operative appliances." I took it for granted that no element would be neglected, and particularly the mechanical fixation, which is so important, in fact the most important element in inhibiting a destructive tubercular process.

In regard to the criticism that many farmer's children who live in the open air have tuberculosis of the joints, I would say that I have not infrequently found farmers houses in far more unsanitary conditions than prevail in the city.

Our patients are benefited in the city hospitals largely because we bring them out of the darkness, and give them good food, in addition to mechanical appliances.

The point that I wished to emphasize was that, in addition to all our other methods, we should employ sunlight, the *x*-ray and the Finsen ray, in fact everything that will help us in the desperate fight with joint tuberculosis.

Dr. J. CLARK STEWART, Minneapolis.—In answer to Dr. Christian I would like the permission of the chair to ask him a question, and that is, whether the washing of sections stained by osmic acid in ether and thus discharging the black color does not prove the blackened substance to be fat. The technical work was done for me by Prof. Margaret Nickerson of the University of Minnesota, and I am confident that Dr. Christian may accept all statements of fact as correct. In regard to the fatty degeneration, as I stated in my paper, the slides were washed with ether, demonstrating the presence of large amounts of fatty degeneration, and as Unna states that fatty degeneration never occurs in epithelioma, this pronounced fatty degeneration was considered worthy of mention. In regard to the vascularization being simply that of granulation tissue, I would say that the presence of the vessels long antedated the presence of anything which could be termed granulation tissue, and even of the leucocytes themselves. Later, where at first red blood cells had appeared in the new vessels, these were crowded with leucocytes, and still later, the leucocytes infiltrated the extravascular tissue. All the changes described in my paper were distinctly progressive during the treatment of the case. I don't attach much importance to the polymorphonuclear form of leucocytes in these sections as in two other cases where the resulting changes were about the same, the leucocytes were entirely of the plasma cell type. I was glad to hear Dr. Nicholls' remarks about the nonoccurrence of Plimmer's bodies in epithelioma. The bodies figured by me closely resemble those pictured by Gaylord in his article in the *American Journal of Medical Sciences*. The study of these cases shows certain positive results which must be of more or less value in explaining the results attained by successful treatment.

In regard to Dr. Willard's paper I can add just one observation of interest. I have treated one case of caries sicca of the shoulder joint with rather striking results. These cases, as you know, usually go on to ankylosis after about two years' disability, while the case under consideration attained about the same result after two months' treatment with the *x*-ray. Apparently the cicatrization and contraction of the tubercular tissue about this joint was very greatly hastened by the effect of the *x*-ray.

English the Universal Language.—A Spanish physician taking a post-graduate course at Vienna writes to the Madrid *Siglo Medico* that he had been informed that some of the lectures were delivered there in French and other languages, but found that this was a mistake. He says: "The lectures are exclusively in German or in English, and the latter is the truly universal language, and he who says the contrary, lies. (*y en ingles, que es hoy el idioma verdaderamente universal: y el que dijera lo contrario miente.*") He adds that the North American colony there is "tremendous," and "Yanquis" are to be met everywhere, very frequently as the assistants at operations.