Effect of serum on fever: 60 subsided, 56 reduced, 26 stationary, 84 not recorded.

Effect of serum on night sweats: 69 subsided, 17 unchanged, 140 not recorded.

Result of serum on weight: 125 increased, 15 unchanged, 27 decreased, 59 not recorded.

Result of serum on strength: 154 increased, 9 unchanged, 24 decreased, 39 not recorded.

Result of serum on appetite: 114 increased, 15 unchanged, 31 decreased, 66 not recorded.

Result of serum on local signs: 40 disappeared, 58 mitigated, 28 unchanged, 100 not recorded.

Result of serum on tub. bacilli: 40 disappeared, 103 reduced, 7 altered, 76 not recorded.

Result of serum on general well being, exclusive of the 40 recoveries: 145 improved; 9 unchanged, 32 not recorded.

Number of recoveries that seemed complete and permanent . . .	40
Number of apparent recoveries with existing lesions (cavities) in statu quo . . .	3
Number of improved capable of performing usual duties . . .	41
Number of improved to a lesser degree . . .	69
Number of deaths reported . . .	32
Number of cases disappeared from observation or under various treatments . . .	41
	226

As to the pulmonary cases, the extent, stage and importance of the conditions at the beginning of were as follows:

In Class 1 there were	20 cases in advanced stage; 3 in early stage
" 2 " " "	33 " " " "
" 3 " " "	12 " " " "
" 4 " " "	9 " " " "
" 5 " " "	12 " " " "
" 6 " " "	7 " " " "
	93
	15

Not classified accurately enough for satisfactory description of the stage . . . 120

I beg to say today that those cases reported as having recovered, dating back from one to three years, are still living and apparently well. None of these cases recorded as having improved greatly have succumbed yet, but some of those reported as having up to that time remained unchanged or mitigated or slightly improved, have since died. I have been unable to get an absolute record as to the number of these, although we are using a method of tabulation and clinical report chart which we send out to physicians using the serum. It seems that not one of the cases reported by doctors as having recovered, or in which the disease seemed to have been arrested permanently, have had a relapse, so far, although most of them remain in an unfavorable climate where the disease originated, as in St. Louis, for instance.

I beg now to present a report of 67 cases, which were not included in the foregoing, and which have been treated in the last year, and purposely recorded since the last meeting of the AMERICAN MEDICAL ASSOCIATION, concerning which I have direct information.

Total number of cases 67.

Effect on fever: 17 subsided, 22 reduced, 5 increased, 23 not recorded.

Effect on night sweats: 26 subsided, 4 unchanged, 27 diminished, 10 not recorded.

Effect on weight: 40 increased, 11 unchanged, 4 decreased, 12 not recorded.

Effect on strength: 34 increased, 10 unchanged, 4 decreased, 19 not recorded.

Effect on appetite: 34 increased, 10 unchanged, 4 decreased, 19 not recorded.

Effect on local signs: 17 disappeared, 22 mitigated, 27 not recorded.

Effect on tubercle bacilli, 17 disappeared, 35 reduced, 15 not recorded.

Number of recoveries that seem complete at present . . .	17
Number of improvements to a considerable degree . . .	35
Improvements to a lesser degree . . .	11
Deaths . . .	4

Thus the grand total of cases recorded above properly enough for reliable report is, not including the various doctors' reports herein mentioned . . . 293

Of these, have recovered . . .	57
Apparent recoveries with lesions in statu quo . . .	3
Improved capable to perform again usual duties . . .	76
Improved to a lesser degree . . .	80
Deaths reported (probably some not known) . . .	36
Cases disappeared from observation . . .	11
	293

This report does not include the numerous cases in which serum was tried in moribunds or in more favorable cases for too short a time. Nor does it include a number of cases improved in the last few months by using serum by rectal injection.

In concluding this report, I wish to say that too much is expected of sero-therapy in tuberculosis, or of any treatment for that matter. It can never be possible to arrest consumption when the tissues are so destroyed and their generative energies so enfeebled that they are beyond the power of stimulation. It is only in the early stages that one may expect the best results (and in the early stages it is a most efficacious remedy). One great trouble we have to deal with is mixed infection and this can be reached only by the use of antitoxins prepared specially for the germs producing the complications, assisted occasionally by other measures. Tubercle antitoxin can not act directly against microbic complications. Furthermore it should not be forgotten that the destructive process of tuberculosis is so great, so comprehensive, that no means of wise special or general treatment should be spared to assist sero-therapy or any other special treatment in the work of repair. Most of them were cases of an experimental character. The cases which I have submitted here have been treated almost exclusively by the use of serum.

I am fully convinced that no one is warranted today in proclaiming the absolute and exclusive curative properties of an exclusive specific alone in tuberculosis of all kinds. Every case must be treated on its merits, considering in each the symptoms, lesions, predispositions, inherited conditions and the various susceptibilities present. Sero-therapy is nature's remedy and offers the greatest assurance of benefit in the subjugation of the specific cause or causes, but in most cases, as just stated, it should be supported in its splendid work by such hygienic, dietetic and medicinal measures as are considered wise for each individual case.

THE CAUSE AND MECHANICAL TREATMENT OF SUBLUXATION OF THE KNEE-JOINT.

Read at the Third Annual Meeting of the American Academy of Railway Surgeons held at Chicago, Sept. 23, 24 and 25, 1896.

BY S. L. McCURDY, M.D.

SURGEON TO THE PENNSYLVANIA CENTRAL RAILWAY.
PITTSBURG, PA.

In order that the treatment of subluxation of the knee-joint may be properly carried out its cause must be thoroughly understood.

Subluxation of this joint is an under-dislocation of the head of the tibia back of, or posterior to, the condyles of the femur. Such a deformity does not exist as a result of chronic disease when the femur is kept parallel with the tibia, during tubercular, synovial, traumatic, gonorrheal and other inflammatory diseases in and about the knee-joint. It can only result from direct violence and from faulty position during the course of treatment.

The series of experiments has been carried out to determine the relative power of the flexor and extensor muscles of the leg, as well as to determine at what

angle, in flexion, the flexor muscles equal the extensors in power. We must remember that extension of the leg is limited to a parallelism with the thigh, and that flexion is only limited by the resistance of the calf with the thigh muscles, or through an arc of 90 degrees. In extension the flexors and extensors are antagonists, and oppose each other with equal advantage, so far as the position of the bones themselves are concerned, because they both have direct action. The advantage, however, is in favor of the extensors, because they are always more powerful. When the leg is flexed the extensors lose their power of direct action and act over a pulley. This pulley is formed by the action of the quadriceps extensor femoris, patella and ligamentum patellæ over the condyles of the femur. As the leg is flexed the extensors lose power and the flexors gain. In order to determine the power of these sets of muscles and to more particularly determine at what angle of flexion the flexors equal the extensors, experiments have been carried out.

Without giving in detail the various experiments which were made upon different individuals, it was found that the extensor muscles upon an average had a resistance of thirty-two pounds, and the flexors twenty-four pounds and that at an angle of about 35 degrees the resistance of the extensors was reduced to twenty-four pounds. This shows that the extensors and flexors have equal power when the leg is flexed to an angle of 35 degrees. A slight variation from the above is found in different individuals.

The experiments were carried out by the use of a goniometer, a pair of spring scales, and four hooks on the jam of a door, one on either side opposite the knee and the other two below these near the floor. To test the extensors, a tape is secured to the hook in front of the knee and thrown around the knee above the joint. The scale is hooked to the hook back of the foot, and another tape is thrown over the ankle and the free ends passed through the ring of the scale. With the leg in extreme extension the tapes are drawn through the ring of the scales. The scales will first register the extreme power of the extensors, but as it is drawn back to about 10 degrees it requires less force to make flexion, hence, a reduction on the scale dial.

To test the flexors, the scales are placed in front of the foot and the knee secured from the hook immediately back of it.

It is quite customary to treat the diseased conditions of the knee-joint in flexion, the degree of flexion being determined by the attending surgeon. Some reasons given for this are that it places the leg in a better position for use during after years, or it is more comfortable to the patient, etc. All of the reasons given are fallacious and founded upon anything but sound reasoning.

In the case of traumatic and other forms of synovitis it is probably true that a greater area of synovial surface is included between the heads of the bone and semilunar cartilage. Such a reduction in actual surface pressed upon could not diminish the pressure, but would place the leg in a less secure position and thus increase muscular spasm and as a result increase intra-articular pressure.

It must be remembered that all muscles which pass over, or the tendons of which pass over or control the action of a joint in inflammation, are in a state of spasmodic contraction. This increased action on the

part of the muscles is an effort on the part of nature to protect the sensitive joint. The deformity which follows is in the direction of the most powerful muscles, or sets of muscles.

In hip-joint disease it is flexion, adduction, or abduction. In tarsal disease it is valgus, varus, flexion, extension, etc. In knee-joint disease it may remain normal, as it generally does in diseases of the shoulder or wrist, so long as the leg is kept in a position favoring this. Just as soon, however, as the flexors have a greater power than the extensors we have an increase of flexion.

As a result of my experiments an approximate estimate of the muscular power showed that the flexion at an angle of 45 to 50 degrees, the flexors had absolute control of the leg. This will admit of two things, an increase of flexion and a tendency of the head of the tibia to glide further back under the condyles. This is not only correct theoretically, but it is borne out clinically and has no doubt been observed by many of my hearers.

I have thus dwelt at length on the cause of the condition under consideration in order that the treatment might be preventive rather than that we be called upon to correct the deformity after its existence.

The treatment of subluxations may be summed up briefly as follows:

It must first be determined whether true or false ankylosis exists, whether the dislocation is partial or complete. In true ankylosis a complete section of the bone must be made either by the use of chisels or saw, and if the relationship of the bones can not be restored without it, a wedge-shaped section should be made, the bones brought together and or otherwise secured.

In cases of ligamentous or false ankylosis with a partial subluxation, *brisement forcé* with tenotomies of the hamstring tendons is generally sufficient. The most difficult cases to manage are those where the adhesions are more or less firm and a complete luxation of the bones exists.

Tenotomies of the tendons will free the lower portion of the head of the tibia, but great force does not as a rule restore the joint surfaces to a normal relationship. It is quite possible, after tenotomy, to bring the leg parallel with the thigh, but it will be found that the leg is upon a lower plane than the thigh. This is due to the fact that the head of the tibia does not slip down and forward upon the condyles, but instead the relationship of these bones are not disturbed.

Immediate reduction of subluxation of the knee-joint is prevented: 1. By the adherent patella and synovial membrane to the condyles. 2. By the impinging of the anterior portion of the head of the tibia against the condyles. In cases where the hamstrings are not cut this burrowing in of the bones upon each other is much greater, and an immediate reduction is less likely to be effected. Gradual and continuous leverage offer the best means for correcting the deformity mechanically. In this degree of deformity a rapid cure can not be promised, except by a radical operation.

A complete correction mechanically must not be attempted at the time of the practice of *brisement forcé*, but instead the leg should be carried forward to an angle that would permit a gradual correction of the subluxation. If plaster of Paris is used as a support, it should be reapplied every ten to fourteen days.

DISCUSSION.

Dr. MAYO—Dr. Gillette of St. Paul, in a number of cases, has broken up the adhesions by means of great force, using some form of mechanical apparatus in one sitting and without tenotomy. I have myself been afraid to apply this extreme degree of force, and, if the deformity is sufficiently great, much prefer to make an osteotomy of the femur after the method of Macewen: this, it appears to me, gives the best results with the least loss of time and danger. The mechanical apparatus of the doctor I believe would succeed better in the hands of the inventor than with the rest of us. In my experience I have seldom been able to succeed with another man's splint as well as does the inventor himself, and as a rule do better with the plaster of Paris cast.

Dr. HAWES—I wish to take exception to one statement in the paper; no mention is made of complete reduction of the subluxation at one sitting. I think the best result I ever had was where it was necessary for me to use force; I actually tore the skin in the popliteal parts. The splint I have been accustomed to use has two bars on the outside and one inside, extending from the inside of the thigh to the heel and outside to the heel and carrying the bandage around it all, behind the tibia. It seems to me I ought to get better results than I would with this splint.

THE DIAGNOSIS OF MALARIA.

BY ARTHUR R. EDWARDS, A.M., M.D.

Professor of Therapeutics, Northwestern University Medical School;
Professor of Practice and Clinical Medicine, Northwestern University Woman's Medical School; Attending Physician to
Cook County Hospital and St. Luke's Hospital;
Pathologist to Wesley Hospital, Chicago.

The diagnosis of malaria, like its pathogenesis, has a scientific life of scarcely two decades. The subject has been roughly handled since an acquaintance with its microscopic diagnostic methods has reached the general profession from the laboratories of scientific biologists and clinicians. As in all newly developed methods, medical and otherwise, there is a tendency to swing from one extreme to the other. Great skepticism greeted Laveran's initial reports that certain parasitic elements existing in the blood of malarial patients and hitherto confused with pigment-bearing leucocytes, were the cause of malaria. The general skepticism of Laveran's colleagues is now superseded by equally universal credulity. While previously no parasites were admitted, now they are even confused with the constituents of the normal blood. Five years ago the existence of malaria in Chicago was denied, while today many vague diseases, especially obscure or obstinate fevers, are diagnosed malaria.

I. DIAGNOSIS OF THE EXISTENCE OF MALARIA.

(a.) *Positive blood findings*, i. e., the detection of the plasmodium of malaria, establishes the fact of malaria, since malaria is always caused by the parasite, and again the organism is almost always found in malaria and in malaria only. A few microscopic examinations will convince the greatest skeptic. It must not be forgotten that, in certain instances, two diseases may occur simultaneously. We have seen malaria in conjunction with various ancient heart lesions, ulcerative endocarditis, pulmonary tuberculosis, chronic nephritis, although never with typhoid fever. These coincidences are, however, infrequent, and the presence of malaria plasmodium makes possible positive differentiation from other diseases, e. g., the frequent error of overlooking or misinterpreting an incipient pulmonary tuberculosis attended with

chills. Atypical fevers, associated with chills and sweating, which have few or dubious local findings and are therefore difficult of diagnosis, can at least be differentiated from malaria; for example, analagous typhoid, septico-pyemia occulta, generalized miliary tuberculosis, febrile gastritis, cholecystitis suppurativa, central pneumonia, deep-seated sarcomatosis, ulcerative endocarditis, etc.

(b.) *Negative blood findings*, in suspected malaria, are not definitive from one examination. Not infrequently is more than one microscopic search necessary for the positive exclusion of malaria. While suggestive, then, a single negative finding is far from conclusive. The parasites may be indistinguishable in the first few days of the disease. In certain forms they swarm in internal organs, avoiding the peripheral circulation; and lastly, in chronic and recurrent types they are found with great difficulty. Sometimes visceral (splenic) puncture is needed for their detection.

(c.) *Fallacies*.—Certain deformities in the red blood corpuscles are often mistaken for plasmodia, e. g., crenations, poikilocytosis and vacuole formation. Not only can the more intimate structure of the red blood discs retract, simulating plasmodia, but the exterior of the hemacyte is far more plastic than is commonly acknowledged, even to the extent of protruding pronounced pseudopodia-like processes. These are but too frequently mistaken for parasites, being found in very many instances of apparently otherwise normal blood. Vacuole formations are characterized by their sharp contour and high luster.

Melaniferous leucocytes are readily distinguishable from the plasmodia by their large nuclei and by their ameboid movement, always absent in adult parasites of equal size. Unstained spores may be confused with the blood plaques, which are, however, structureless and contain no pigment. An Austrian pediatricist lost a docentship for reporting as malaria cases whose blood preparations afterward proved to contain only blood plates and no plasmodia. Coagulation products have been confused with flagella. Many of the small dots seen in malaria which resemble micrococci and were mistaken for such by the earlier Italian observers are similar to those found in most anemias and described by Ehrlich as degenerative changes.

II. DIAGNOSTIC TECHNIQUE.

Complex methods of staining and counterstaining the parasite have been in vogue, but the simplest and most accurate is the direct examination of the freshly-drawn unstained blood, a method we have used with entire satisfaction for several years. In this procedure injury to the corpuscles and staining of the blood plaques are obviated.

The lobe of the ear is cleansed, pricked and a quite small drop is gently expressed. A clean cover glass is held in a pair of forceps to avoid the heat and moisture of the hand, and is carefully brought in contact with the top of the drop. The heat and moisture of the hand or rudely placing the cover against the drop favor imperfect spreading from precipitate drying of parts of the blood. Rubbing the slide well facilitates equable spreading of the blood. Examination is best made with an oil one-twelfth inch immersion lens, although Laveran used lenses of lower magnification. Permanent preparations are procured by allowing the covers to dry, to remain half hour in equal parts of absolute alcohol and ether and by painting with filtered eosin and methylene blue. The use of stains is not usually