rapid pulse. When this patient was admitted to the hospital the heart sounds were strong and distinct, but through an error the operation was delayed three hours after admission. The child perished during this time.

In the third case the uterus grasped the child so firmly at operation that the lower end of the uterine incision was slightly torn in delivering the fetus. This patient had one rise of temperature during her convalescence, which soon subsided.

Kerr believes that craniotomy should be positively declined when the child is living and in good condition. He believes that sterilization should not be practised, because the mortality is now not sufficiently great to prevent a repetition of the operation if necessary. He believes that it is well to select the time of operation to have the circumstances favorable. It is important not to wait until the lower uterine segment has developed, because this is readily torn and easily infected. The placenta in each case was on the posterior wall of the uterus and the Fallopian tubes were close together upon the anterior wall. In the Fallopian tube tissue excised decidual cells were not found.

Puerperal Tetanus.—In the *Indian Medical Gazette*, vol. xxxvi., No. 5, 1901, KEDARNATH DAS reports the case of a Hindoo woman delivered at full term of a living child. On the tenth day the infant was seized with trismus neonatorum, and died within twenty-four hours. The mother developed tetanus, and was admitted to a hospital. She was treated by washing out the uterus with an antiseptic solution, with the free use of chloral by rectal injections and by hypodermoclysis. The spasms continued six weeks before the patient began to grow essentially better. An abundance of liquid food was given and the bowels were kept open regularly. During her illness the patient received 8 ounces, 6 drachms, and 10 grains of chloral hydrate. Das has collected 68 cases of puerperal tetanus, showing a remarkably small percentage of recovery. He could find but five recoveries, giving a mortality of 92 per cent. There were 21 cases treated by antitoxin, given either by subcutaneous or intracerebral injections. Of these but one recovered, a mortality of 95 per cent. Among these cases of tetanus but 4 occurred in India, where the disease does not seem to be common. One of these recovered, giving a mortality of 75 per cent.

Prolonged Retention of the Fetus During Labor by the Enormously Dilated Cervix.—RAPIN (*La Semaine Médicale*, November 13, 1901) reports the case of a primipara, aged twenty-five years, who was suffering from a tedious labor. The head came down into the pelvic cavity and upon the pelvic floor, when the attention of the midwife was attracted by a tumor situated between the symphysis and the umbilicus. The descent of the head ceased and assistance was summoned. On examination the pelvis was found somewhat contracted in its antero-posterior diameter. The head was in the pelvic cavity, the smaller fontanelle obliquely and toward the left. The fetal heart sounds were not heard, and upon the presenting portion of the head there was a tumor which bled somewhat. A tumor of elastic consistence was found between the pubes and the umbilicus. Thinking that this was a distended bladder, the patient was catheterized, but the tumor did not dis-

appear. The forceps were then applied, but the head could not be delivered with moderate traction. Digital examination could discover no apparent reason for the retention of the fetus. The patient was accordingly transferred to the hospital. Upon examination there it was discovered that the two arms of the fetus seemed engaged in the pelvis beside the head, and that if these could not be dislodged the child could not be delivered. Accordingly the fetus was cautiously delivered by version and found to be dead and somewhat macerated. It was perfectly formed and in no way monstrous. Immediately after delivery there was free hemorrhage, which ceased after the expulsion of the placenta, a hot intra-uterine douche of lysol and compression of the uterus. The abdominal tumor disappeared. Upon further examination it was found that the abdominal tumor had been composed of the upper uterine segment containing the placenta. The child had been contained in the enormously distended lower uterine segment and cervix.

He also reports the case of a multipara who had had six normal confinements previously. She had a flat rachitic pelvis. The head was presenting and fixed in the pelvis, the heart sounds feeble and irregular. Meconium was escaping from the vulva, and there seemed to be fetid gas in the uterus. It was evident that the cervix was enormously distended. The forceps were applied at the superior strait, and a male child asphyxiated was delivered. The child was resuscitated. The placenta was removed and the uterus thoroughly doused with a hot antiseptic solution, when upon examination it was found that the upper uterine segment had been invaginated into the lower. It was enormously distended.

Both of these patients recovered, which must be considered an excellent result in view of the very serious distention which was present.

Labor Brought on by Alcoholic Neuritis.—LE PAGE reports in the *Comptes Rendus de la Société d'Obstétrique de Paris*, 1901, vol. iii., the case of a patient in her second pregnancy whose first pregnancy had terminated normally. In the present gestation the patient suffered severely from nausea and vomiting, and had great difficulty in walking, with much pain in the lower limbs. A physician whom she consulted ordered baths and a milk diet. She finally became much worse, with frequent pulse, elevated temperature, dry tongue, delirium, and marked and excited movements. The lower extremities were paralyzed, the feet extended upon the legs, and the hands were immobilized and flexed upon the forearms. There was no inequality of the pupils and no strabismus. The abdomen was slightly distended, and there were sordes on the borders of the gums and teeth. There was nothing abnormal about the heart and lungs and no albumin in the urine. The appearance of the patient was that of a severe typhoid infection.

The serum diagnosis was negative, and labor was induced by the introduction of an elastic bag. During the treatment of the case it was learned that the patient had used large quantities of alcohol, drinking rum very freely. The delivery was spontaneous. The patient afterward made a very tedious recovery, with atrophy of the muscles, which yielded slowly to electrical treatment. The case was a typical one of polyneuritis, undoubtedly of alcoholic origin.