SECTION OF ANATOMY AND PHYSIOLOGY.

CONGENITAL MALFORMATION OF THE HANDS.

By S. W. ALLWORTHY, M.A., M.D., F.C.S. ;

Physician, Belfast Hospital for Skin Diseases.

[Read in the Section of Anatomy and Physiology, January 31, 1902.]

THE photographs and radiographs of this case taken by me are interesting as illustrating some peculiarities of development. They are of a man's hands, aged thirty, who was a seven months' twin. In the right hand all the bones of the middle finger are absent, and there is not a trace of them to be seen in the skiagraph. The left hand has all the bones, but the first phalanx of the index finger evidently developed across the palm of the hand (which is well shown in the skiagraph) and its pressure wedged the bones of the second and third fingers into each other, so that they became united. The piece of bone between the second metacarpal and the second phalanx is probably an overgrowth of the base of the first phalanx, due to the displacement during ossification. The hands are wonderfully useful, and there is no other deformity.

There is no history of any similar occurrence in the family, and the only fact of interest related by the mother is, that during the early months of pregnancy she fell across a wall whilst stretching over to feed pigs.

The other twin was smaller, but otherwise perfectly developed.

A second case came under my notice a short time ago in

which both hands had all the bones of the middle finger missing, and presented an appearance identical with the right hand in the above. There was no history of importance, and my patient, who was not a twin, in this instance said his mother always attributed the deformity to the loss of a wedding ring.

Congenital Malformation of the Hands.—No. I.

Congenital Malformation of the Hands.--No. II.

DR. T. G. MOORHEAD ON THE ABDOMINAL AND PELVIC VISCERA
IN ADVANCED PREGNANCY.