abating a little; bowels again relieved at midnight. Pulse: morning, 96; night, 100. Hand treated as before; reduced solution of morphia to three minims. A little more support taken during the day.

Nineteenth day.—Slept well; hand treated as before. Pulse: morning, 100; night, 104. Not so much pain; bowels not relieved; took more support; muscles still very rigid along the whole of spine.

Twentieth day.—Slept well; hand treated as before. Pulse: morning, 108; evening, 100. Half a pint of beef-tea and about a quart of milk taken during the day; bowels not relieved; more pain towards night; spasm very strong again; hand looking very healthy; pupils fairly dilated.

Twenty-first day.—Slept well; hand treated as before; pain increasing towards night. Pulse: morning, 100; evening, 104. Not quite so much support taken; bowels not relieved; six drachms of castor oil given.

Twenty-second day.—Slept well; spasms not so violent, about an hour between each; took support better; hand treated as before. Pulse: morning, 104; evening, 96. Bowels relieved early in morning; very cheerful when not in pain.

Twenty-third day.—Did not sleep so well; support taken freely; a great deal of pain through night and day. Solution of morphia increased to five minims; hand treated as before. Pulse: morning, 108; night, 100. Bowels not relieved; muscles still very rigid.

Twenty-fourth day.—Slept better through night; hand treated as before. Pulse 100, night and morning. Spasm not so violent; bowels not relieved; took fair amount of support.

Twenty-fifth day.—Slept well; spasm abating; hand treated as before; ate part of pigeon, and able to open mouth a little. Pulse: morning, 100; evening, 96. Bowels relieved; rigidity of muscles abating a little.

Twenty-sixth day.—Slept well; spasm not nearly so violent. Pulse 104. Ate a little fowl; hand washed with carbolic lotion (1 to 40), and dressed with lint; poulticed over dressing at night.

Twenty-seventh day.—Slept well; hand treated with carbolic lotion as before. Pulse: morning, 100; night, 108. Bowels relieved three times during day; spasm more frequent towards night, but of no length.

Twenty-eighth day.—Did not sleep so well; hand treated as before, and poulticed over dressings at night; very obstinate, and shouted a great deal, causing spasm. Pulse: morning, 100; night, 108. Bowels relieved twice during the day; tongue bitten very much during the night.

Patient slept with his mother from this date, and progressed favourably. On the thirty-second day a spasm was brought on by touching loose tooth. On the thirty-sixth day he was able to move his head backwards and forwards, lift himself up in bed by the aid of a swing, but could not bend to sit. Put his tongue more than half out on the thirty-eighth day. The morphia was discontinued altogether on the thirty-ninth day. He was able to walk with help on the forty-fourth day.

The patient went on well from this date. The granulations on the thumb did not heal very well up to the sixtieth day, when it was dressed with styptic colloid, after which it rapidly healed, and the lad is now in good health, and goes to school.

Ancaster.

NOTES

ON THE USE OF THE HÆMACYTOMETER IN ANÆMIA.

BY JOSEPH W. HUNT, M.D. LOND.,

PHYSICIAN TO THE WOLVERHAMPTON AND STAFFORDSHIRE GENERAL HOSPITAL.

DR. GOWERS, by the introduction of his modification of the hæmacytometer, has opened up to the medical profession in this country a vast field for research into the pathology and therapeutics of diseases of the blood and the blood-making organs. He himself has made and published some observations in this direction.¹ Having for some time used his instrument in a large number of cases, I have been induced to state the result of my observations, with especial reference to the condition of blood in anæmia.

An extended use of the hæmacytometer only confirms the statement of those who estimate the number of red corpuscles in health as being about 5,000,000 in the cubic millimeter, or 100 in the two squares (Dr. Gowers' "hæmic unit") of the hæmacytometer; though an excess of 1,000,000, or a diminution to the same amount, is quite compatible with the most perfect health. The number of white corpuscles is very much more variable; but by taking the average of a much larger number of squares than suffices for the enumeration of the red corpuscles generally—that is to say, 150 to 200 squares—I have found the average not so inconstant as I at first supposed from a more limited examination. The maximum number of white corpuscles has been stated as .3 per "hæmic unit," or 1500 in the cubic millimeter of blood; but they may frequently fall below this.

In all investigations with the hæmacytometer it must be remembered that we cannot attempt to compute by it the number of corpuscles in the body, but rather the corpuscular richness of the blood—that is to say, the proportion which the corpuscles bear to the serum of the blood; and again, that the number of the corpuscles alone does not represent their value, for as much perhaps depends upon their quality—i.e. the amount of hæmoglobin—as upon their quantity.

In ten cases of anæmia which I have carefully followed throughout the greater part of the course of the disease, the average of the red corpuscles at the onset of treatment has been 62 per cent., and in the three worst cases the average was less than 46 per cent., the corpuscles in the most severe case of all averaging as low as 42 per cent. None of these patients were laid up by their disease, all being able to walk to the out-patient room, and some of them a pretty considerable distance. All were young girls, of ages varying from fourteen to twenty-two years or thereabouts, and engaged as pupil-teachers, seamstresses, barmaids, shopwomen, or some such occupation.

One of the most striking facts brought out by the hæmacytometer was, as Dr. Gowers was the first to point out, that the corpuscular richness of the blood is seldom accurately represented by the appearance either of the skin or the mucous membranes. Thus in some patients, when the skin has been markedly pale and the mucous membranes equally anæmic, the corpuscles have averaged 80 to 90 per cent., or as much as they have averaged in my own blood when in a fair state of health. In apparently the worst case of anæmia I have met with, a young girl seen by me in consultation with my friend, Mr. Bunch, of this town, the patient was confined to her bed, from which indeed she could scarcely stir; the pulse was very frequent; there were numerous syncopal attacks, causing her medical attendant to be summoned more than once at midnight, as the patient was believed by her friends to be dying; there was frequent vomiting; the face was nearly as pale as the sheets which covered her, and the conjunctivæ were almost blanched except where one or two bloodvessels pursued their solitary course. Together with these symptoms there was some suspicion of hysteria. I examined the blood with much interest, and found the number of corpuscles not far from normal, so that the case was evidently not one of true anæmia. Under appropriate treatment, the patient was, to all intents and purposes, well in a fortnight.

Again, the corpuscles reach the normal amount some time before the patient's symptoms have disappeared or the normal colour returned, and, in some cases at least, long before the catamenia have recurred in their accustomed quantity and at the regular intervals; that is, as Dr. Gowers has shown, iron may increase the number of corpuscles out of all proportion to the increase in their hæmoglobin. In the greater number of my cases it has taken very little more than three weeks for the corpuscles to reach their normal amount, and sometimes much less than that time. Thus in one case of advanced anæmia, the corpuscles, which before treatment were as low as 60 per cent., reached 110 per cent. in two weeks, and after another week had advanced to 120. In a second patient the corpuscles rose in thirteen days from 42 to 98 per cent., but afterwards sank again on the iron being discontinued. On the other hand, some cases are much less amenable to treatment; and in one patient it took seventy-two days to raise the corpuscles from 48 to 96 per cent., but during this period the catamenia recurred three times, and the patient was one week without medicine.

The condition of the heart in most of these cases proved

¹ Practitioner, July, 1878.

the severity of the disease. In the majority there was well-marked dilatation, as expressed by the character of the impulse and increased lateral dulness. Mitral systolic murmurs were not infrequent, and, with I think one exception, there were distinct basic murmurs, and in most an intensified second sound was heard over the pulmonary cartilage. Carotid murmurs were heard in each case. These physical signs all improved under treatment, and speedily so, the improvement taking place apparently *pari passu* with the increase in the number of the corpuscles, and previous to any marked change in the outward appearance of the patient.

Absence of the catamenia, sometimes extending over a long period, was a marked feature in many of these cases, or if not entirely absent, they were very deficient in quantity and quality (as shown by their colour). The return of the catamenia to the normal condition occurred generally at a later stage in the period of recovery than the improvement in the other symptoms, but even this frequently took place before four weeks had elapsed. In one patient, in whom the catamenia had never appeared, the anæmic condition was cured in eighteen days, but neither then nor during the rest of the time that she remained under treatment did the catamenia appear. The return of the catamenia was almost invariably accompanied by a diminution in the number of corpuscles, or at least with a temporary discontinuance of any increase in their number. Thus, in one patient the corpuscles fell from 64 per cent. before the onset of menstruation to 58 after it, and at the second period from 84 to 74 per cent., at which time, however, the patient was also a week without medicine. In another patient the corpuscles were examined the day before menstruation recommenced, and though for the last four weeks they had been steadily increasing from 73 to 107 per cent., they had now sunk to 104. The catamenia lasted two days, and when the patient was again seen, four days after their cessation, the corpuscles had continued their rise to 118 per cent. In a third patient they rose in three weeks to 120 per cent., but when re-examined at the end of the fourth week, the catamenia had recurred slightly the week before, and the corpuscles had diminished in number to 106 per cent. Similar instances could be added. Possibly owing to the examination of the blood not being made every day during the catamenial period, the results may be less conclusive than they might otherwise have been; but, imperfect as they are, they appear in strange contrast to the observations of Dr. Duperié, who states that menstruation provokes the formation of a very considerable number of corpuscles. It is true that in one or two of my cases the gradual increase in the number of the corpuscles, noticed before the onset of the catamenial period, was continued when the blood was examined the first time afterwards; but, on the other hand, in no case where the corpuscles averaged the normal amount, or above this, did menstruation occur without a diminution in the number of corpuscles, either immediately preceding or found immediately after such period.

In all the cases mentioned above the treatment was the same, and consisted of good plain nourishing food without stimulants, open-air exercise, a cold bath in the morning when practicable, together with half an ounce to an ounce of compound iron mixture thrice a day, and aperient every night, mostly five grains of aloes-with-iron pill or aloes-with-myrrh pill. With regard to the number of the white corpuscles throughout the disease, I noticed that they were not reduced in anything like the same proportion as the red corpuscles when the patients came under treatment, but were generally about the normal number, and that their number was far less affected by the treatment pursued.

The marked effect of iron in increasing the number of the red corpuscles was well seen in a case of surgical anæmia. The patient, a young man of about eighteen years old, was admitted into the hospital under the care of my friend and colleague Mr. Vincent Jackson, with a fractured skull. A considerable portion of the skull was removed by operation. Secondary hæmorrhage from some meningeal vessel which could not be reached came on, at the commencement of which the corpuscles averaged 76 per cent. The hæmorrhage continuing, they decreased in a week to 50 per cent., when the common carotid was ligatured. The operation was a bloodless one, and the hæmorrhage ceased from that time forward, but notwithstanding this, the corpuscles still diminished in amount, till three days after the operation they were as low as 38 per cent. The patient was then ordered iron (ferrum redactum, five grains in the twenty-

four hours) and good nourishing diet. In two days the corpuscles had risen again to 50 per cent., and in another fortnight, notwithstanding the profuse continuous suppuration from extensive wounds and the sloughing off of a small hernia cerebri, the corpuscles reached 100 per cent., and when again examined, five days after this, they were the same in number. The patient at this period was practically well, only remaining in bed for some superficial wounds to heal and for the recovery from hemiplegia due to the injury.

It is a point of considerable interest to observe the lowest number of corpuscles compatible with life. Dr. Gowers says that any patient with less than 20 per cent. is in imminent danger, and that, with one exception, he has not observed a lower percentage than 18 per cent., but this latter low average was maintained for some weeks in a case of pernicious anæmia. The exception referred to was that of a case of very profuse hæmatemesis, where the corpuscles numbered about 9 per cent. immediately before death. In one case of leucocythæmia, recorded by Sørensen, the red corpuscles were reduced to about 9 per cent.; but as the white corpuscles averaged as high as 10 per hæmic unit, the corpuscular poverty of the blood was not so profound as the number of red corpuscles suggests. In a case of purpura hæmorrhagica admitted into the hospital under my care, with epistaxis, bleeding from the gums, purpuric spots, and retinal hæmorrhage, the patient had only 18 per cent. of red corpuscles. Large doses of iron, turpentine, ergot, and aromatic sulphuric acid failed to produce any effect. In two days they had still further diminished to 14 per cent., at which period the prick with the lancet only yielded a little colourless serum, and it required considerable squeezing of the finger before any coloured blood could be obtained. On the third day after admission the corpuscles had sunk to 12 per cent., or rather less, at which number they remained till death took place on the fifth day. The white corpuscles varied from '2 (the normal amount) in the two squares, on admission, to '03, '05, and '08 on subsequent occasions. In a second case of purpura hæmorrhagica, when admitted, the red corpuscles were 75 per "hæmic unit"—i.e., per cent.,—and the white as high as 4'5. Owing to successive hæmorrhages, the number rapidly diminished; and when death took place on the fifth day, they numbered 23 per cent., and the white '15. In a third case, the day before death, the corpuscles averaged 30 per cent., and the white '6. Death took place rather suddenly, so that the blood was not examined again during life; but immediately after death the red had sunk to 24 per cent., and the white had risen to 2 per cent.

Wolverhampton.

A NEW FORM OF FEMALE SYRINGE.

By HENRY GREENWAY, M.R.C.S.

THE want of an efficient instrument for bringing various forms of injections in contact with the entire surface of the vagina and the os uteri has long been felt, and many of the leading writers on the diseases of women refer to the inadequate means generally employed for this purpose. In order to cleanse the vagina, or locally treat its diseases by injections, it must be dilated in such manner that its walls may form a tube, the dilator itself offering as little surface as possible; and the fluid to be injected should be delivered freely and from many points. These requirements can be fulfilled by the invention now brought before the notice of the medical profession. The accompanying engraving gives a very good idea of the new appliance. It consists of four small-size rods, about six inches long, curved at their upper ends so as to meet at a central point, and connected at their lower ends with the outer margin of a cylindrical stock, the spaces between the rods being equal. This part may be termed the dilator. In the centre of the stock is a hole in which is fixed a vagina-tube, which extends upwards nearly to the point of the dilator, and downwards about an inch below the stock, and it has four rows of perforations which face the spaces midway between the rods of the dilator, each perforation being directed diagonally upwards. On the lower end of this tube is fixed three or four inches of india-rubber tubing to receive a nozzle. The vagina-tube should be made of vulcanite, or some incorrodible material, but the