by precipitation by galvanic agency, as given by thin plates of iron and zinc, and iron and silver. The quantity of the poisonous salts taken by deceased was eight or ten drachms, consisting of two or three drachms of the commercial sulphate of iron and seven of sulphate of copper, and which had been purchased for agricultural purposes.

No medical man was called in during the exhibition of the poison; the symptoms were detailed by the witnesses at the inquest held upon the body.

August, 1856.

ON A

CASE OF DIFFUSE ABSCESS IN THE NEIGHBOURHOOD OF THE KNEE-JOINT.

By C. H. MARSTON, Esq., M.R.C.S.E., Henlow.

On the 26th of May last, I was called upon to see J. M——, a young man, twenty years of age, by trade a builder. Six days previously he had knelt for a length of time upon some damp timber, and in the evening felt a soreness and smarting in the right knee, which shortly afterwards became red and inflamed. Three days afterwards he had severe rigors, followed by headache, fever, and increased pain and swelling.

I found him lying on his back, the countenance expressive of much suffering; the right thigh was laid perfectly straight, the knee being much swollen, red, and painful; the swelling surrounded the limb, and extended from the spine of the tibia to about an inch above the patella. To the touch the tumour was hard and tense; no tendency to point was obvious at any place. The pain was described as throbbing and starting, the patient being continually awakened by "something shooting through the knee" when falling off to sleep. No appetite; tongue covered with a brownish fur; pulse 98; skin hot and damp; bowels have been opened by senna and salts. To apply at once twelve leeches, encouraging the bleeding by poppy and camomile fomentations; ten grains of the compound ipecacuanha powder to be taken at bed-time, followed by an aperient draught in the morning.

On examining the knee next morning, I found the swelling somewhat softer, and yielding a more decided sense of fluctuation to the touch. The patella could scarcely be distinguished by the finger, nor could the line of the ligamentum patellæ be discerned. It was evident that the suppuration was external to the joint, and surrounding the whole of the limb, the extent of the tumour excluding the idea of its being an inflammation of the bursa patellæ. I had the limb placed on a hard mattress, the parts surrounding the knee being well padded, so as to prevent "bagging." A valvular incision was then made just over the outer tuberosity of the tibia, and about twelve ounces of sanious pus escaped. A probe on being introduced into the wound was found to pass freely in all directions on the circumference of the limb; and on passing one probe externally, and another internally, they were found to meet beneath the integuments behind the knee.

It was evident that we had a diffuse abscess in the cellular tissue overlaying the deep fascia of the leg, this structure being itself uninjured. The extent of integument separated from the deeper parts was very considerable—the integuments being so extensively undermined that sloughing was feared. For twenty-four hours after opening the abscess, poultices were applied, the posterior and lateral parts of the limb being well supported by pads.

On the second day, the whole limb (leaving a small space for the discharge of pus) was bandaged on the principle recommended by Mr. Chapman; a portion of skin, about the size of a crown piece, was all that perished; and the walls of the abscess were, in the course of three days, found firmly adherent. The resulting ulcer was for some time disposed to be rather indolent in its action; but, with some humouring, completely closed in the course of six weeks.

I may just mention the great benefit derived from Curtis's "pasma," with which the sore was on several occasions dusted. I was anxious to have adopted during the whole treatment Mr. Chapman's method of bandaging, but so great was the irritation during the extremely hot weather which once or twice occurred, that I was compelled again and again to abandon it. This irritability of skin may be accounted for: during the progress of recovery, several crops of bullæ, some of which threatened trouble, showed themselves; they were, however, controlled by emollient lotions, and the internal use of Dono-

van's solution. The patient's health, during the whole period that followed the evacuation of the abscess, was vigorous. He ate well, drank well, slept well, and, to use his own expression, "felt well." Never have I seen so troublesome an eruption of bullæ, combined with so good a state of health.

There are two or three points worthy of note, as—

1. The intense inflammation set up in so close a proximity to the knee-joint.
2. The extensive undermining of the integuments.
3. The efficacy of the wetted strips of bandage in promoting the adhesion of the separated integument, and preventing further sloughing.
4. The troublesome cutaneous complication.

Bedfordshire, August, 1856.

FOREIGN BODY IN THE EYE.

By JAMES MARTIN, Esq., M.R.C.S.E., &c.,
Quebec, U.S.

PERHAPS the following case may be worthy of a corner in your journal. It was brought to my recollection on reading Mr. Critchett's paper in THE LANCET of Nov. 17th, 1855.

John R——, aged twenty-four, was brought to my surgery on March 10th, 1854. He stated to me that, about an hour previously, while in the act of driving a copper bolt, a splinter of the bolt flew off, and struck him on the right eye. On separating the lids, I found the cornea divided horizontally in its whole extent. I found also a slight wound on the edge of the lower lid. He did not complain of inconvenience beyond the peculiar sensation of the sudden deprivation of the sight of one eye. I could detect no foreign body in the eye, and did not feel justified in exploring for one. I imagined that the wound had been produced by the fragment of copper glancing swiftly past. I ordered him to live upon arrowroot and gruel, to keep ice-water constantly applied, to take every morning a wineglass of compound infusion of senna, and to remain in a darkened room.

For ten days he went on without complaint, sleeping well, and feeling comfortable. On the morning of the eleventh day, I was sent for in haste. I found my patient in great agony; he had been suffering since early in the morning. It struck me at once that the fragment had lodged in the eye, and I proceeded to remove it. With a pair of dressing forceps closed, I explored the eye, and having touched a hard substance, I seized it. With a slight amount of force, I drew from its bed, in the posterior part of the globe, a splinter of copper, three quarters of an inch long, one quarter of an inch in breadth, and weighing twelve grains. In a short time the pain was relieved, and the patient from that time suffered nothing. The globe, collapsed, would now form an excellent bed for an artificial eye, did his circumstances allow him to obtain one.

1856.

CASE OF

RUPTURE OF THE MENINGEAL ARTERY; OPERATION.

By HENRY WATSON, Esq., M.R.C.S.E.,

LATE SURGEON TO H. M. STEAM TRANSPORT "GOLDEN FLEECE."

JOHN HAWKINS, aged twenty-seven, a spare, delicate-looking man—the "Boots" in the *Golden Fleece*,—was seized, on the 18th of April, at the commencement of a voyage to the Crimea, with a fit of epilepsy, for which I was called to see him. I found him lying on the deck, with all the usual symptoms of that affection. In a few minutes he recovered his consciousness, and was able to answer my questions; and walked to his berth, apparently well. On my seeing him half an hour afterwards, to my surprise, I found him drowsy, stupid, and nearly insensible; in an hour's time he became completely so. His pulse became full, slow, and labouring; the pupils fixed and dilated; the skin hot; and his motions were passed involuntarily. I ordered his head to be shaved, and carefully examined it to see if there was any external fracture. None, however, was perceptible, a slight bruise only being visible on the right side. I therefore ordered a large blister to be placed on the scalp, fifteen grains of calomel to be given through the