the patient to any great extent, at least not for any length of time, for the urine flows into the bladder and over the excoriated and irritable surface of the tumor, and thus necessarily leads to permanent pain. The other method, which furnishes more satisfactory results, consists in establishing permanent kidney fistulæ and thus diverting the urine from the bladder; the patient is at once relieved from all pain and is made quite comfortable. A belt carrying two small pouches keeps the patient comparatively dry.

The choice of anesthesia is of the utmost importance in all cases of bladder surgery. Endovesical operations can, as a rule, be carried on under the influence of an antipyrin anesthesia supported by the rectal application of morphin. For the cutting operations, we always employ, as a matter of routine, the anesthesia produced by laughing gas. This anesthesia, besides being free from danger, so far as the heart is concerned, possesses a great advantage in that the patient returns to full consciousness immediately after the completion of the operation; this enhances the possibilities of natural micturition, of very early breathing exercises, and of the early administrations of food and drink. Up to the present time we have never observed any untoward influence of gas-anesthesia on the kidneys.

Fig. 10.—Inner aspect of the bladder showing the implanted ureter. Flap operation.

The next question that naturally arises is whether or not it is possible to determine, previous to the operation, the character of a bladder tumor; we do not hesitate a moment to answer in the affirmative and to say that such diagnostic work can be done with accuracy sufficient for all practical purposes. Any man with experience in cystoscopic diagnosis can recognize a benign tumor from its appearance (Nitze, Weinrich); tumors of doubtful nature are to be considered and treated as malign. In this connection it should be borne in mind that the benign papillomata comprise the majority of bladder tumors (Civiale, Thompson, Guyon, Soemmering, Kuester).

It was formerly held by a great many surgeons that, because of the poor permanent results and the quickness of the relapses, patients with malign bladder tumors should never be operated on. This view is shared at the present time by some writers. In our opinion, however, the soundness of such reasoning is, to say the least, open to discussion because, in the first place, a great many patients were and are still being operated on in an irrational manner, and, in the second place, because the diagnosis is quite often made so late that the chances for a successful operation are greatly reduced; the more

general employment of the cystoscope, however, will greatly improve matters as to the latter point. Furthermore, it should be taken into consideration that the use of rational operative methods and the uniform employment of gas anesthesia will certainly reduce the primary mortality of these procedures.

As it is, there are two cases of vesical cancer on record in which total extirpation of the bladder accomplished a cure without relapse for five years (Hogge), and for fifteen years (Pawlik), respectively. Rafin reports five cases of partial resection of the bladder for carcinoma that remained cured for more than three years. In Czerny's sarcoma case the patient lived for twelve years after the operation without relapse. Quite a few surgeons report cases of malign bladder tumor which, after partial vesical resection, have remained cured for from one to three years (L. Davis, Pousson, Pitha, Leschnew, Albarran, Alessandri, Schmidt, Kolischer).

Mikulicz, in his last publication before his death, declared that he felt convinced that he had relieved the sufferings and prolonged the life of his patient every time he had operated for cancer of the bladder. Surely so impressive a message gives warrant for our belief that, even though nothing else were accomplished than the adding of a few years of comfort to the sufferer's life, the operation would be fully justified.

CONCLUSIONS.

In conclusion we desire to submit, as the result of our work along these lines, the following fundamental propositions:

1. All benign tumors of the bladder should be approached from the inside, all malign tumors from the outside of the viscus.
2. In all malign cases in which the loss of substance is not too great the bladder should be closed completely by sutures after the removal of the tumor.
3. In cases of malign tumor the incision into the bladder should be made in accordance with the location of the growth, as defined by the cystoscope.
4. The permanent catheter should be absolutely abolished.
5. Gas anesthesia should be employed exclusively.
6. A constant cystoscopic surveillance should be maintained over any bladder that has ever been operated on for tumor.

ONYCHOMYCOSIS TRICHOPHYTINA,

WITH REPORT OF TWO CASES.*

A. RAVOGLI, M.D.

CINCINNATI.

The nails, like any other epidermic tissue, are exposed to the invasion of the trichophyton, yet cases of the kind occur rather rarely. Saubouraud, in the department of the ringworm affections in the Hospital St. Louis, considers onychomycosis trichophytina a rare disease. Neuberger¹ referred to the fact that Anderson, in Glasgow, among 11,000 cases of cutaneous diseases, of which 178 were of trichophytic nature, did not have a single case of trichophyties of the nails. White, of Boston, in 5,000 cases of diseases of the skin, with 180 cases of tinea trichophytina, did not have one case of trichophytic onychomycosis. On the other hand,

* Read in the Section on Cutaneous Medicine and Surgery of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

1. Ein Fall von Onychomycosis Trichophytina, Derm. Ztschr., xi, 551.

Pellizzari, Arnozan, Dubruille have often seen cases of the kind, and Pick considers this affection of the nails not so rare as stated by the others.

In some cases of onychomycosis trichophytina, the origin of the fungus is clearly found as patches of herpes tonsurans in other parts of the body, showing the infection of the nail to be due to the act of scratching. Mahon noted the coincidence of herpes tonsurans of the scalp, with that of the finger-nails, which are in this way more frequently affected than the toe nails.

In rarer cases the nails only are infected with trichophyton, and yet the patient has never had any trichophytic affection in other parts of the body. A case of this kind was reported by Gerson,² where a man had the nail of the second finger of the left hand for two years affected with trichophyties, and gradually all the other finger-nails were equally infected, without showing any patch of trichophyties in any part of the body. The case of Neuberger¹ is of the same kind and shows that onychomycosis trichophytina can be found without there being other places of infection on the body, and that this condition exists a long time.

The assertion of Zinssers,³ that tinea tonsurans of the nails never occurs unless preceded by attacks of the same in other parts of the body, falls to the ground.

ETIOLOGY.

It is easy to understand that any one who is affected with trichophyties of the body, and more especially of the beard and of the scalp, by scratching with his finger-nails, gets the spores of the fungus underneath the nails, and so the nails are infected.

To this class of cases is to be referred the case of Bosc and Galavielle⁴ in a man who was suffering with an extended attack of trichophytosis of the body, and consequently showed alterations of the finger-nails, which microscopically and through culture revealed the presence of the fungus.

The presence of the trichophyton in the nails without any previous attack of trichophyties in other parts of the body gave to Hirtz and Jacquet⁵ the idea that in these cases the fungus does not produce the disease primarily, but only secondarily, as a consequence of trophic disturbances of the nail. They reported two cases of trophic alterations of the nails, which showed great similarity, and in one they found the presence of trichophyton, while in the other it was absent.

I thoroughly agree with Krzystalowicz,⁶ who maintains that, in order to establish the diagnosis of onychomycosis trichophytina, there must be found the fungus under the microscope.

Recently in my practice I have had two cases of onychomycosis, which are interesting to consider, one after the other, on account of their different clinical and pathologic features.

CASE 1.—History.—A man of 54, of splendid physique, married, had always enjoyed good health. In himself and his family no history nor trace of syphilis, psoriasis nor eczema could be found. He had been suffering for over twenty-years with his finger-nails and toe-nails, which did not cause him pain, but only discomfort. He was by occupation a flour merchant,

and when he began to notice his finger-nails rough and diseased, he was under the impression that it was produced by an insect called "weaver," which is found in the flour. The first finger-nail to be affected was that of the second finger of the left hand. At the time I saw him it had completely disappeared and was replaced by a thin, atrophic and somewhat irregular scar. Some time later the nail of the third finger of the same hand was affected and there remains (Fig. 1) only a thin atrophic peeling. Gradually every nail showed signs of disease by losing its brilliancy, getting chalky, scaly and irregular. The nails were easily broken, causing inconvenience in the use of the hands. Some time afterward several toe-nails were equally affected. In all these years the patient never had any reproduction of trichophyties in any other part of the body. He had in the meantime consulted several physicians and many remedies had been given. The only benefit the patient had ever obtained was from bathing the fingers in a solution of bichlorid of mercury.

Examination.—A few months ago the man called on me to see if something could be done. On the first examination, the long standing of the affection, the absence of any ringworm

Fig. 1.—Hands of patient in Case 1, showing destruction of nails.

attack in any part of the body, were against the possibility of trichophyton infection. Every nail of the fingers and toes was affected in a different degree according to the time of its infection, so much so that while in the second and third fingers of the left hand the nails had been entirely lost, in the other fingers they were affected in different degrees. The nails were somewhat thick, opaque; their surface rough, uneven and showing longitudinal folds. The nails kept on breaking and splitting in longitudinal direction, and pieces of nail were hanging on the nail-bed. Where the nail had fallen or had been removed, as on the thumb of the left hand, an ugly irregular surface of the nail-bed remained, looking as if riddled by small holes. In order to establish the diagnosis a piece of nail was taken from the thumb of the left hand, it was softened in a solution of potash and then examined under the microscope. The fibers of the nail-plate contained spores and mycelia of trichophyton, which were arranged in rows between the edges of the epidermic cells forming the structure of the plate of the nail. Another piece of nail was washed in alcohol, crushed

2. Berlin. Dermat. Gessellsch., January, 1899; ref. Arch. f. Derm. und Syph., xlviii, 136.

3. Encyklopedie der Haut und Geschlechtskrankheiten, 1900, p. 225.

4. Onychomycose trichophytique, Arch. f. Derm. und Syph., xli, 419.

5. Hysterotoxia, Onychomycose trichophytique, Arch. f. Derm. und Syph., li, 447.

6. Die Pathogenen Haarpilze, Poln. Zeitsch. f. Derm. und Ven. ref. Monh. f. Pract. Derm., xlii, 428.

and was dropped into a culture medium of maltose-agar. After eleven days, around the piece of nail there appeared a light whitish area, which gradually extended by peripheral progress until it had invaded the whole surface of the culture. Some of the vegetation of the fungus was examined under the microscope, it proved to be a clear specimen of the ordinary trichophyton. The diagnosis was confirmed as that of onychomycosis trichophytina.

It was difficult, however, to conceive how it was possible that nails could get infected without any other point of the body ever having shown trichophyties, and that the toe-nails could also be infected without producing ringworm infection in other parts of the body. The explanation has to be found in the anatomic structure of the nail-plate and in the way that the fungus vegetates therein.

The photomicrograph (Fig. 2) shows the fungus penetrating between the interstices of the nail-plate, and the nail fibers separated just as we see it occur in the hair. In order that the vegetating fungus may be

Fig. 2.—Photomicrograph of nail-plate of patient in Case 1, showing fungus penetrating between the interstices of nail-plate. Bausch and Lomb 1/6 in. objective.

introduced in the interstices of the nail fibers, there is necessary a trauma of the nail. This patient used to trim with thin scissors the affected nails and then with the same scissors the other nails. It is clear that with the scissors he carried spores of trichophyton from one nail and inoculated them deeply into the structure of the other nails.

The fungus vegetating inside of the fibers of the nail is protected and can not easily be transported to other parts. This satisfactorily explains how the trichophyton could remain for so many years in the nails without being conveyed to other parts of the skin.

It will now be of some interest to make a brief comparative study of the other case of onychomycosis, where the scalp had been affected with tinea trichophytina.

CASE 2.—History.—In this case a few weeks after the patient had the affection of the scalp three finger-nails of the right hand became badly infected. The nails were of those

three fingers which he used in scratching his head to relieve the itching. The occipital region of the scalp was covered with large irregular trichophytic patches (Fig. 3), evidently the result of several round spots coalescing. The spots were partially bald, studded with stumps of hair, covered with thin, dirty, adherent scales, showing the clearest features of the trichophyties of the scalp. The nails were inoculated a short time afterwards and Figure 4 shows that the appearance of the affection consisted in a more acute process, and the injury produced in the nails was much more perceptible. Indeed the nails were entirely of a dirty yellow hue, chalky, thin and atrophic towards the matrix, and thickened and irregular towards the free edge of the nail-plate. The surface showed holes and crevices in the nail substance and large pieces of the nail were breaking and detaching themselves in a transverse direction. This is easily explained when we remember that in this case the fungus had not been brought to the internal structure of the nail-plate, but it had been placed on the surface of the nail in its eponychium, hyponychium and in the subungual epidermic layer. Consequently the fungus had to work its way through, so as to reach the laminar structure of the nail.

This is well proved by the microscopic examination and by the photomicrograph (Fig. 5) which was made by my assistant, Dr. George H. Werk.

The action of the fungus on the epidermic structure, according to Leslie Roberts, is that of digestive power on the horny tissues. It penetrates into them, making them its habitat. It detaches the epidermal cells one from another, and by digesting the hard membrane appropriates their plasma as nutrition and consequently the epidermal cells die.

Fig. 3.—Scalp of patient in Case 2, showing large trichophytic patches.

In the first case where the nail-plate had been primarily affected, the resulting alterations were somewhat different. The fungus with its spores and mycelia was disposed in long rows between the epidermic fibers composing the nail-plate, just in the same way as it is disposed in the cortical layer of the hair. Consequently the fibers composing the body of the nail were torn one from the other and the nail was breaking in a longitudinal way, according to the direction of the fibers. In the second case where the fungus from the eponychium had invaded the nail superficially, the nail had shown a tendency to break irregularly in large scales and more often transversely.

In order to ascertain whether the difference of the alterations could be produced by different kinds of trichophyton I resorted to the culture. Maltose-agar was used as a culture medium in the Erlenmeyer bottles. One culture medium was inoculated with a piece of nail of the patient in Case 1, and in the two others one was

inoculated with scales of the scalp and the other with a piece of nail of the patient in Case 2. In all positive results were obtained. The culture medium inoculated with the epidermic scale of the scalp showed, four days after the inoculation, a whitish, dusty halo around it, which gradually began to spread peripherally. The nail culture of the second patient showed the same condition after six days, while the nail culture of the first patient did not show signs of any result until after eleven days. The result of the culture was the same in every one, consisting of an abundant whitish, purulent, thin growth of the fungus, trichophyton megalosporon endo-ectothrix, with large spores of the common kind often taken from domestic animals.

The cultures proved that the fungi were the same, but this did not furnish any explanation in reference to the difference of the pathologic alterations. The only explanation I find is in the way of the introduction of the trichophyton into the nail structure. In the first case it was inoculated by the scissors in cutting and trimming the nails, and the spores were introduced

Fig. 4.—Showing infection of nails in same patient as Fig. 3, due to scratching of scalp.

into the interior structure of the nail-plate. In the second case the spores had been deposited in the eponychium and in the subungual layers, and consequently they had attacked the nail from the outside toward the inside of the nail-plate.

The distribution of the fungus in the superficial layers of the nails, or in the intimate structure of the nail-plate, is a sufficient reason for the difference of the course in the two cases. In the first case the disease lasted for over twenty years without giving a chance to the infectious spores to be transported into any other part of the body. The spores growing between the interstices of the fibers forming the nail-plate are protected, and it is not easy for them to be transferred to other parts except by cutting the nail. While in the second case where the parasite was vegetating on the eponychium and in the subungual folds of the finger, the nail was attacked from the external layers towards the inter-

nal. The distribution of the parasite explains how easily the spores of the trichophyton can be transferred from the nails to other parts of the body.

In my cases the ultimate result of the fungus on the nails has been that of causing complete atrophy. My experience is more in accordance with that of Kaposi,⁷ who maintained that in the trichophytic affections the nail disappears by atrophic process. Censi,⁸ however, referred to the possibility of the hypertrophy of the nail in form of onychogryphosis and made a variety of onychogryphosis trichophytina. I believe, however, that in those cases the alterations of the nails were not due entirely to the parasite, but the dystrophic conditions were due to other causes and the trichophyton was of later occurrence.

PROGNOSIS.

In cases in which the parasite is hidden in the interstices of the cells of the nail-plate, it can be reached only with great difficulty, and the obstinacy of the case, lasting for over twenty years, shows that the prognosis is bad. In the cases in which the fungus is on the surface of the nail-plate, however, it can be reached and the fungus can be destroyed, and with constant treatment we are rewarded by seeing the nail grow free from the disease.

TREATMENT.

In the case of the first patient it appears, from the condition of the nails, that anything which was used brought only a temporary relief, while in the case of the

Fig. 5.—Photomicrograph showing spores and mycelia of trichophyton invading substance of nail.

second patient with constant applications of Wilkinson's ointment the patient obtained perfect recovery within four months. This shows that in one case the spores were protected in the interstices of the cells forming the plate of the nail, and can be reached only with great difficulty. It is difficult, indeed, to bring a case of onychomycosis of the first type to recovery, so much so that in many cases it has been necessary to remove the nails. The affected finger is wrapped up in lint, well saturated with salve containing a large percentage of pyrogallie acid so as to produce an inflammatory condition of the nail, an artificial paronychia. In this way the nail is detached from the surrounding epidermis and it can be removed. This method is rather severe and not all patients are ready to submit to it.

7. Die Patologie und Therapie der Hautkrankheiten, Vien.

8. L'onicogriphosi Tricofitica; Giorn. Ital. delle malattie veneree e della pelle, xxxiii, 5.

My patient, when the pyrogallie acid began to inflame the skin, stopped its application and discontinued treatment.

Pellizzari⁹ reported one case of onychomycosis as cured by means of the Roentgen rays. He claimed that nine exposures of ten minutes each were sufficient to effect the recovery. It was my intention to apply the rays, but the patient did not come back.

The other patient was kept in the hospital and was put to work as an orderly in the ward. Good results were obtained from the application of unguentum Wilkinson to his scalp and fingers. I ordered his scalp and his fingers to be scrubbed with pure green soap once a day, and then rubbed and covered with the ointment. When he left the hospital the trichophytic patch of the scalp was perfectly healed, and the nails had all grown out new, showing no signs of alterations. The patient, so far, has not returned, which I consider a proof of permanent recovery.

DISCUSSION.

DR. J. B. KESSLER, Iowa City, said that six or seven years ago a farmer reported having barn itch. He had a fungoid eruption on the back of the hand, and its inflamed, elevated border contained pus. Accompanied by a veterinarian, Dr. Kessler visited this farmer's place and found about the noses and necks of the calves circles that looked like asbestos. These lesions were scraped close to the roots of the hair, where numbers of trichophyton fungi were found. Several calves and one horse were affected. Farmers thus infected are very apt to carry the infection to the barber shop. Dr. Kessler considers this the most common method of spreading the disease. It is a curious fact that, as soon as spring time comes and the farmer turns his calves out to pasture, they recover very rapidly from this affection, and when they are again confined to the pens during the cold weather they again contract the disease. Those who do not live in a farming district will probably find, on investigation, that trichophytosis is spread in the barber shops from human to human, and that this type of disease is comparatively easy to cure, but when a patient is thoroughly inoculated from a calf, the affection is much more virulent and more difficult to eradicate.

DR. M. L. HEIDINGSFELD, Cincinnati, said that he is firmly of the opinion that the infection of the trichophyton does not take place directly in the nail substance, but in the tissues from which the nail is originally formed. It is difficult to conceive how a hard, non-succulent substance like nail, after it has once been perfectly formed, can be a culture medium for any form of infection. Products like the nail and hair once removed from the body show no effect after inoculation with fresh, active cultures. The infection of the trichophyton occurs primarily in the nail-fold and extends thence to the matrix, which becomes inflamed and so pathologically changed that any nail substance which is subsequently formed is rough and irregular in outline, opaque in appearance and contains areas of cells corresponding to layers of imperfect keratinized cells, derived from the matrix. These round, imperfect keratinized cells are the round bodies which Dr. Ravogli has mistaken for spores. These conclusions were drawn by Dr. Heidingsfeld in a paper presented to this Section some seven or eight years ago. That nail substance when placed on culture media should grow fungus, possesses no significance inasmuch as contamination of this character occurs from almost any substance under similar conditions.

DR. GRANVILLE MACGOWAN, Los Angeles, Cal., agreed with Dr. Heidingsfeld that the infection in these cases takes place in the nail-bed. It is an exceedingly disagreeable affection, particularly in women, whether they be servants or in the higher walks of society. Dr. MacGowan has seen a number of these cases during the past five or six years, and has been rather at a loss what to do for them in order to get a speedy recovery, until he happened to think of the old remedy com-

mon among farmers and known as "sheep dip," or Vlemineckx's solution, which is practically a solution of calcium sulphid, in which they soak their hands at night. He has found this very efficacious as a temporary expedient, while for permanent relief he prescribes an ointment of the oleate of tin, which, together with the x-ray, proves most satisfactory.

DR. M. L. HEIDINGSFELD, Cincinnati, expressed the opinion that the best results of treatment are obtained if attention is directed to the nail-fold and not to the nail itself. The remedy he has found most efficacious is a 1 per cent. solution of corrosive sublimate in ether and alcohol.

DR. A. RAVOGLI, Cincinnati, said in regard to the non-inoculability of the nail to trichophytosis, that the presence of the trichophyton in the nail tissue has been successfully proved by inoculation on culture medium. The spores can not be so well shown in the microphotograph on account of the difficulty of staining them.

TWO CASES OF TUMOR AT THE PONTO-CEREBELLAR ANGLE WITH AUTOPSIES.

THEODORE DILLER, M.D.

Neurologist to the Allegheny General Hospital, Visiting Physician to St. Francis Hospital (Psychopathic Department).

PITTSBURG, PA.

In presenting these two cases no attempt will be made to review the literature of the subject of tumors at the ponto-cerebellar angle, which has now become quite extensive. They are recorded as a contribution to the subject for what they may be worth as clinical and post-mortem records.

CASE 1.—Patient.—A single man, clerk, aged 29, was admitted to the Allegheny General Hospital on April 18, 1905, in the service of Dr. T. M. McKennan, to whom I am indebted for the early notes of the case and who was first to make the correct diagnosis.

History.—The family history was unimportant. The patient had enjoyed good health up to the time of his present illness. His personal habits had been good. He denied syphilis. The present illness began about a year previously with headache, which soon became of daily occurrence, and grew more and more severe. It was noted about this time that his gait was uncertain, like that of a drunken man. About a month later the patient complained of some failure of vision and also of occasional dizziness. By autumn the uncertainty of gait, augmented by dizziness, had grown to such an extent that the patient was compelled to use a cane in getting about. In November he had some difficulty in speaking and swallowing. Speech was not quite so clear, and an occasional difficulty in swallowing occurred. He masticated with difficulty on the left side; he used the right side exclusively. The head pains became especially severe by December and were localized chiefly in the occiput. About three weeks before admission these pains left him. They were nearly always associated with vomiting, which was sudden and forceful. He had sometimes vomited when he did not have headache. On admission he complained of numb sensation over the right side of the body. He also complained of some dizziness.

Examination.—(Dr. McKennan.) The patient was a strong, well-nourished man. The pupils reacted well to light and accommodation. No satisfactory examination of the eye-grounds. There was slight nystagmus. A slight internal squint in the left eye was present. The tongue was protruded toward the left side. There was no wasting of the tongue or lips. The patient's enunciation was not distinct. His saliva dribbled when he was speaking or eating. The sense of taste was absent over the left half of the tongue, and the sense of touch almost absent. On the right side taste and touch were present, but decidedly impaired. The patient masticated on the right side exclusively. Left masseter weak. With the left ear the patient could not hear the ticking of a watch. With the right ear hearing was better, but impaired. The sense of smell seemed good. There was some enlarge-

9. Lo Sperimentale, 1906, ix, 6.