VII.—*On the Physiological Action of Light.* By JAMES DEWAR and JOHN GRAY
M'KENDRICK, M.D. (Part I. Plates X., XI.)

(Read 21st April, 5th May, and 2d June 1873.—Received, 6th February 1874.)

I. INTRODUCTION.

Sensory nerves are divided into two classes, those of general sensibility and those of special sense. The nerves of general sensibility are distributed to the skin, muscles, or viscera, and convey influences to the brain which give rise to sensations of touch, heat, &c., or to those vague sensations, not definitely localised, which we include under the name of the muscular sense. The nerves of special sense are endowed with special and individual physiological properties. When a nerve of this order is irritated in any way, either by mechanical, chemical, or electrical stimuli, an influence is conveyed to the brain which gives rise to the same kind of sensation as that produced by the normal stimulus on the terminal organ. For example, pressure on the eyeball, as shown by NEWTON and YOUNG, electrical stimulation by a continuous current, as demonstrated by PFAFF,* HELMHOLTZ,† RITTER,‡ PURKINJE,§ DU BOIS-REYMOND,|| and SCHELSKE,¶ produce many of the phenomena of vision, including not only the perception of light, but the perception of various colours and tints. But while this is the case, it is equally certain that each terminal organ responds to its normal stimulus. Thus the retina, though capable of stimulation by pressure or electricity, is specially fitted by its histological structure for the reception of those minute vibrations of the ether which constitute light. But while the terminal organ is capable of receiving a most delicate action of the normal stimulus, the nerve in connection with it is not so affected. For example, although the retina is affected by light, the optic nerve is not so, as may be proved by MARRIOTTE's** well-known experiment, by which it may be demonstrated that when the image of an external object falls on the entrance of the optic nerve, there is no corresponding sensation. The nerve is thus insensible to the normal stimulus of the sense organ, the retina. The current in the nerve cannot be excited by the direct

* Thierische Elektrizität, 1795, p. 142.

† Physiologische optik, p. 203.

‡ Beiträge zur näheren kenntniss der Galvanismus und der Resultate seiner Untersuchung. Jena, 1805, p. 159, *et seq.*

§ Beobachtungen über Versuche zur Physiologie der Sinne. Band i.

|| Thiersche Elektrizität. Band i. p. 353.

¶ Zur Farben-empfindung. Graefes Archives ix. p. 49.

** Phil. Trans. 1668, p. 668; 1670, p. 1023.

application of the normal stimulus, but only through the intervention of a terminal apparatus. For each sense a special terminal organ is required, the function of which is to receive the normal stimulus. This stimulus affects a change in the terminal organ, and the result is conveyed by the nerve to the nerve centre.

Now arises the question of what is the specific effect of the external stimulus on the terminal organ. With regard to the ear, for example, there is the hypothesis that the hair-like processes of the cells situated on the delicate rods of Corti found in the *scala media* of the cochlea vibrate synchronously with waves of sound, and communicate these vibrations to the minute filaments of the auditory nerve which are in connection with them. With reference to vision, numerous theories have been put forward. NEWTON, MELLONI, and SEEBECK stated that the action of light on the retina consisted of a communication of mere vibrations; YOUNG conjectured that it was a minute intermittent motion of some portion of the optic nerve; DU BOIS-REYMOND attributed it to an electrical effect; DRAPER advanced the view that it depended on a heating effect of the choroid; and MOSIER compared it to the action of light on a sensitive photographic plate. Up to this date, however, our knowledge of the specific effect produced by light on the retina is without any experimental foundation.

Now, it is evident that, in accordance with the principle of the transference of energy, the action of light on the retina must produce an equivalent result, and this result may be expressed and measured as heat, chemical change, or electro-motive force. The change in the retina will likewise excite corresponding changes in the nerve, which will be conducted to the brain. What are the properties of a living nerve which can be affected by changes in the condition of the retina?

A nerve has three distinct properties: 1st, sensibility, or the property by which it is capable of receiving an impression and producing an influence; 2d, conductivity, or the property of conducting the influence to or from a nerve centre; and, 3d, electro-motive force, or the difference between the electrical potentials of two surfaces of the nerve, such as the longitudinal surface and transverse section. These three properties exist only in living nerve. When a nerve is removed from the body, and deprived of an adequate supply of healthy blood, it dies, and the rapidity of its death may be measured by the gradual loss of these properties. Now, we know that these properties are affected by a stimulus applied to the nerve. When a portion of a nerve is subjected to the influence of a continuous current of electricity, that portion passes into a condition termed the electro-tonic state, in which these three properties are altered, and altered differently in the neighbourhood of each pole. It is difficult, however, to ascertain with accuracy, by physical methods, the alteration in the properties of sensibility and conductivity; but it is comparatively a simple matter

to ascertain changes in the electro-motive force of the nerve. It has been shown by numerous physiologists that, in the case of a motor nerve, the electro-motive force suffers a negative variation ; that is, its amount is diminished during an active condition of the nerve. This negative variation of the natural electro-motive force may be looked upon as the electrical expression of the physiological process taking place in the nerve during the passage of the normal nerve current. Analogy would lead one to expect a similar change in the electro-motive force of a sensory nerve during the transmission along its fibres of the influence produced by the action of the normal stimulus on the terminal organ with which the sensory nerve is connected. Thus, variations in the electro-motive force of the retina and optic nerve may be regarded as functions of the external exciting energy, which in this case is light.

Such considerations led to this investigation. It resolved itself into a matter of careful experiment in two directions : 1st, to determine the electro-motive force of the visual apparatus and optic nerve; and, 2d, to ascertain whether or not, and to what extent, this electro-motive force was affected by light.

The only experiment bearing on the first of these questions, namely, the electro-motive force of the optic nerve, was one made by DU BOIS-REYMOND, and thus described :* “ Having prepared the optic nerve of a large tench in such a manner that one extremity was the artificial transverse section, and the other the globe of the eye freed from all adherent particles of muscle, found that every point of the external surface of the ocular globe was positive to the artificial transverse section of the optic nerve, but negative to the longitudinal section.” From this experiment it therefore seemed to be quite possible, by using a delicate instrument, to obtain a measurement of the electro-motive force of the retina and optic nerve.

With regard to the second question, namely, the action of light on this electro-motive force, the problem seemed difficult to solve, but an experiment made by Mr, now Justice, GROVE, held out the hope of success. It is detailed as follows :†—“ A prepared daguerreotype plate is enclosed in a box filled with water, having a glass front, with a shutter over it. Between this glass and the plate is a gridiron of silver wire ; the plate is connected with one extremity of a galvanometer coil, and the gridiron of wire with one extremity of a Breguet's helix—an elegant instrument, formed by a coil of two metals, the unequal expansion of which indicates slight changes in temperature ; the other extremities of the galvanometer and helix are connected by a wire, and the needles brought to zero. As soon as a beam of either daylight or the oxy-hydrogen light is, by raising the shutter, permitted to impinge upon the plate, the needles are deflected. Thus, light being the indicating force, we get *chemical action* on the plate,

* MORGAN'S Electro-Physiology, p. 458.

† GROVE. The Correlation of the Physical Forces, 5th edition, p. 153.

electricity circulating through the wires, *magnetism* in the coil, *heat* in the helix, and *motion* in the needles." A consideration of this remarkable experiment led to the hope that we might be able, by a somewhat similar method, to observe the action of light on the electro-motive force of the retina and optic nerve.

The apparatus adopted for obtaining the normal electro-motive force was that of DU BOIS-REYMOND, usually employed in determining the electro-motive force of muscle or nerve. This consists of DU BOIS-REYMOND's well-known non-polarisable electrodes, which are shallow troughs of zinc, carefully amalgamated, containing a solution of neutral sulphate of zinc, and having inserted into them cushions of Swedish filter-paper, on which to rest the preparation, which, in our experiments, was the eyeball or optic nerve. To protect the latter from the irritant action of the sulphate of zinc, a film or pad of sculptor's clay, moistened with 75 per cent. solution of common salt, and moulded, if necessary, by the fingers, to a point, is placed on each cushion. These electrodes are connected with a galvanometer, a key being interposed by which the current from the electrodes may be shut off at pleasure. The eyeball, carefully freed from adherent particles of muscle, is now placed on a glass support between the two clay points, and these are now adjusted to it so that one touches the cornea, and the other the transverse section of the optic nerve. On now opening the key, the galvanometer at once indicates a strong current passing from the longitudinal to the transverse section of the nerve. (The arrangement of the apparatus will be understood by referring to the plate. Plate, No. X.)

The first experiment we performed was on the eye of a rabbit. The galvanometer employed was a large multiplying instrument made by SAUERWALD of Berlin. The animal was killed by chloroform, and the eyeball removed as quickly as possible from the orbit. It was then placed between the troughs in the manner already described. We obtained a deflection of from 10 to 15 degrees. On allowing the beam of light from a magnesium lamp to impinge on the cornea there was no effect observable. This experiment was repeated many times with the same negative results. It then became evident that a much more sensitive instrument was required, because the amount of deflection was so small, and the variation, if any, of this deflection produced by the action of light would probably be so minute as to escape notice.

We then used a sensitive THOMSON's galvanometer, which was kindly placed at our service by Professor TAIT. With this instrument we at once obtained a deflection of several hundred degrees on a millimetre scale placed at a distance of 26 inches, so that we had now an opportunity of observing whether light produced any variation in the electro-motive force.

After placing the galvanometer in position, secondary difficulties now presented themselves. It will be necessary to allude to these difficulties, and the methods by which we avoided or overcame them.

1. *The gradual death of the Retina and Nerve.*—When the eyeball is placed on the cushions in the manner above described, and the key interposed between the electrodes and the galvanometer is opened, there is at once a large deflection. But this deflection diminishes as the nervous apparatus dies. At first the nerve dies by fits and starts, afterwards more slowly, and with considerable uniformity. After a lapse of from 20 to 30 minutes, the rate of dying, as measured by diminution of electro-motive force, is almost arrested, and the vitality of the nervous apparatus remains nearly constant. These facts will be evident on considering the data of the following experiment :—

RABBIT'S EYE.—*Rate of Death as measured by diminution of Electro-motive Force.*
 Zero, 164. Polarity, 156. Total of first deflection, 118.

Time P.M.	Readings on scale.	Difference.	Time P.M.	Readings on scale.	Difference.
H. M. S.			H. M. S.		
1 43 0	274	0	1 55 0	250	4
1 44 0	282	8	1 56 0	246	4
1 45 0	288	6	1 57 0	242	4
1 46 0	285	3	1 58 0	238	4
1 47 0	282	3	1 59 0	234	4
1 47 30	Zero 164.		2 0 0	231	3
1 48 0	278	4	2 1 0	230	1
1 49 0	274	4	2 2 0	239	1
1 50 0	270	4	2 3 0	228	1
1 51 0	266	4	2 4 0	228	0
1 52 0	262	4	2 5 0	228	0
1 52 30	Zero 164.		2 6 0	228	0
1 53 0	258	4	2 7 0	228	0
1 54 0	254	4			

Time occupied in above experiment, 24 minutes.

Total fall in electro-motive force during that period 46°.

The polarity remained constant throughout the experiment at 156°.

2. *Change of the Zero point of the Galvanometer from Magnetic Alterations.*—This was greater at certain times than at others. On several occasions the variations were so sudden as to prevent any accurate experiments being made.

3. *Changes in the Polarity of the Electrodes.*—This was one of the greatest difficulties in the inquiry, but we have employed the most approved methods of overcoming it—such as by the use of neutral sulphate of zinc, by moistening the clay with a 75 per cent. solution of pure chloride of sodium, and by connecting the electrodes together by a thick copper wire for several hours before using them. With a sensitive galvanometer it is almost impossible to maintain either a constant zero or a constant amount of polarity. This fact would be a very serious one in an inquiry such as this, if the variations to be observed in the deflections occurred slowly and through minute distances, because, in these circumstances, it would be manifestly impossible to discriminate between changes due to variations in polarity and those due to the action of light. But

we found that the changes produced by the action of light occurred suddenly, and lasted only for a brief period of time, so that there could be no possibility of mistaking them for changes in polarity. We have also eliminated as far as possible any error from variations in polarity by multiplying experiments and by performing them under as favourable conditions as could be devised. Frequently, during the course of a series of observations on an individual eye, the polarity was observed and any changes noted.

4. *Effects of Heat.*—The effects of radiant heat were eliminated by causing the light to pass through a spherical double shell of glass, having at least an inch of water between the walls. In our earlier experiments we found that heat produced immediately an increase in the electro-motive force. It was therefore absolutely necessary to eliminate radiant heat.

II.—EFFECTS OF LIGHT ON THE EYE OF THE RABBIT REMOVED FROM THE HEAD IMMEDIATELY AFTER DEATH.

Four series of experiments were made on one of the eyes of four individual rabbits, as follows:—

SERIES I.—RABBIT'S EYE.—*Source of light, a magnesium lamp. Zero, + 185; polarity, + 125; total first deflection, 130°.*

Observations.	Degrees of Galvanometer scale.					Remarks.
	Immediately before the impact of light.	Immediately after the impact of light.	Immediately after removal of light.	Effect of the impact of light.	Effect of the removal of light.	
1	255	259	250	Rise of 4	Fall of 9	*Fresh transverse section of nerves now made.
2	250	256	250	Rise of 6	Fall of 6	
3	250	250	245	No rise	Fall of 5	
4	254	258	254	Rise of 4	Fall of 4	
5	255	258	254	Rise of 3	Fall of 4*	
6	185	186	179	Rise of 1	Fall of 7	
7	175	175	165	No rise	Fall of 10	
8	165	166	155	Rise of 1	Fall of 11	
9	155	157	147	Rise of 2	Fall of 10	
10	150	152	142	Rise of 2	Fall of 10	

The result of this experiment was, that *on the impact of light there was an average rise of nearly two degrees; the electro-motive force slowly diminished; and on the removal of light there was a rapid fall of from five to seven degrees.* The rise in this instance on the impact of light, we now know, was owing to the action of heat. At this early stage of the inquiry it was deemed necessary to focus by a lens the light of the magnesium lamp on the eyeball.

SERIES II.—RABBIT'S EYE.—*Source of light, a magnesium lamp. Neutral point, + 195; polarity, + 175; total first deflection, 150.*

Obs. 11. Before the impact of light the reading on the galvanometer scale was at 225, giving a total deflection of 150. On the impact of light it fell at once to 215—a fall of 10. The electro-motive force now diminished in the dark as follows, at intervals of 15 seconds :—*Obs. 12 to 27*—190, 185, 180, 175, 170, 165, 162, 160, 161, 159, 157, 156 155, 153, 152, 150, 149.

Obs. 29. A heated poker brought near it caused an immediate rise to 190. On the removal of the poker, it fell slowly to 153. The following four experiments were then made, the source of light being the flame of a magnesium lamp held at a distance of three feet, and allowed to fall directly on the eye :—

Observations.	Readings of Galvanometer scale before impact, at intervals of 15 seconds.	Effects of the impact of light.	Effects of the removal of light.	Remarks.
30	153, 153, 152, 152, 152, 151, 151, 150	143, or fall of 7	146, or rise of 3	The result of these four experiments is a fall of about 5 degrees on the impact of light, and a rise of 2 degrees on its removal.
31	145, 146, 146, 146, 145, 144, 144, 143	138, or fall of 5	140, or rise of 2	
32	140, 140, 140, 140, 140, 140	135, or fall of 5	137, or rise of 2	
33	135, 135, 135, 133, 133, 133	128, or fall of 5	130, or rise of 2	

After readjusting the eye on the cushions, the following experiments were made, in which the light was passed through a flat-sided glass vessel filled with water, one inch in thickness. The source of light, as before, was a magnesium lamp :—

Observations.	Before impact.	After impact.	After removal.	Effect of impact.	Effect of removal.	Remarks.
34	110	105	106	Fall of 5	Rise of 1	Result is a fall of about 3 degrees on impact of light, and a rise of 2 on removal of light.
35	104	103	105	Fall of 1	Rise of 2	
36	100	97	98	Fall of 3	Rise of 1	
37	94	91	92	Fall of 3	Rise of 1	
38	90	87	89	Fall of 3	Rise of 2	
39	86	83	85	Fall of 3	Rise of 2	
40	81	78	80	Fall of 3	Rise of 2	

SERIES III.—RABBIT'S EYE.—*Source of light, magnesium lamp. Zero, — 60 ; polarity, — 80 ; total first deflection, 238.*

Observations.	Before impact at intervals of 15 secs.	After impact.	After removal.	Effect of impact.	Effect of removal.	Remarks.
41	158, 158, 159, 159	155	156	Fall of 4	Rise of 1	Light fell in these experiments on the cornea.
42	155, 152, 152, 152	150	154	Fall of 2	Rise of 4	
43	155, 155, 155, 155	150	152	Fall of 5	Rise of 2	
44	152, 152, 152, 152	148	149	Fall of 4	Rise of 1	
45	150, 150, 150, 150	155	157	Rise of 5	Rise of 2	Light fell in these experiments on the back of the eye.
46	155, 154, 154, 155	160	161	Rise of 5	Rise of 1	
47	150, 150, 149, 148, 148, 148	152	153	Rise of 4	Rise of 1	
48	168, 168, 168, 168	166	167	Fall of 2	Rise of 1	The eye was re-adjusted on the electrodes, and light fell on cornea. Readjusted.
49	168	166	167	Fall of 2	Rise of 1	
50	168	165	168	Fall of 3	Rise of 3	
51	170	166	167	Fall of 4	Rise of 1	
52	180	176	178	Fall of 4	Rise of 2	
53	180	176	180	Fall of 4	Rise of 4	
54	180	178	180	Fall of 2	Rise of 2	

The general result of these observations is, that when light fell on the cornea, there was an average fall of over three degrees on the impact of light, and a rise of nearly two degrees on the removal of light ; while on the other hand, when the beam fell on the back of the eye, there was always a rise, which we believe to be attributable to heat.

Three observations were made on this eye with a lighted lucifer match as the source of illumination. It was found that if the match was held sufficiently near the eye so as to heat it, there was a sudden increase in the electro-motive force, to the extent of from ten to fifteen degrees; but the light from the match was insufficient to affect the retina. The effect of heat in increasing the electro-motive force was observed in many other experiments.

SERIES IV.—RABBIT'S EYE.—*Zero, — 117 ; polarity, — 65 to — 80 ; total first deflection, 280.*

Observations.	Before impact.	After impact.	After removal.	Effect of impact.	Effect of removal.	Remarks.
55	200	197	200	Fall of 3	Rise of 3	Source of light from a gas-flame. Passed through $\frac{1}{2}$ in. of water.
56	196	195	No effect	Fall of 1	No rise	
57	195	194	No effect	Fall of 1	No rise	
58	196	194	195	Fall of 2	Rise of 1	
59	196	195	No effect	Fall of 1	No rise	
60	195	194	No effect	Fall of 1	No rise	

RABBIT'S EYE—(Continued).

Observations.	Before impact.	After impact.	After removal.	Effect of impact.	Effect of removal.	Remarks.
61	200	196	No effect	Fall of 4	No rise	} Eye readjusted; light from magnesium lamp.
62	195	193	No effect	Fall of 2	No rise	
63	240	236	237	Fall of 4	Rise of 1	} Eye readjusted; source of light was the gas-flame.
64	235	233	234	Fall of 2	Rise of 1	
65	244	242	243	Fall of 2	Rise of 1	
66	245	243	245	Fall of 2	Rise of 2	} Eye readjusted.
67	245	243	244	Fall of 2	Rise of 1	
68	248	247	248	Fall of 1	Rise of 1	} Source of light from magnesium flame, passed through $1\frac{1}{2}$ inch of water.
69	244	240	244	Fall of 4	Rise of 4	
70	242	239	240	Fall of 3	Rise of 1	
71	240	238	238.5	Fall of 2	Rise of 5	
72	240	237	No effect.	Fall of 3	No rise	
73	236	234	No effect	Fall of 2	No rise	
74	234	232	No effect	Fall of 2	No rise	

The result of the above twenty observations is that the impact of light produced a fall in the electro-motive force of about two degrees on the impact of light, while on the removal of light there was a rise of about one degree in most cases; but in a few there was no rise at all.

III.—EFFECTS OF LIGHT ON THE EYE OF THE FROG REMOVED FROM THE HEAD
IMMEDIATELY AFTER DEATH.

At this stage of the inquiry it became evident that, owing to the rapid death of the nervous textures of warm blooded animals, it was difficult to obtain sufficiently great galvanometrical readings, the variations of which could be readily observed. We, therefore, began to employ the eye of the frog, the tissues of which, as is well known, retain vitality long after the systemic death of the animal. The eye was prepared by removing as far as possible all adherent tissues from the sclerotic, and was so placed between the electrodes that one electrode touched the cornea, while the other was in contact with the transverse section of the optic nerve. When the eye was placed in this position a large deflection was obtained, and frequently it was so great as to pass beyond the scale. As was the case with the eye of the rabbit, the electro-motive force slowly fell to a point at which it was on the whole tolerably permanent. Readings were not made until it reached this stable condition. To secure not only a large deflection, but also sensitiveness to light, accurate adjustment of the eyeball on the cushions was necessary.

Ten series of experiments were made on ten frogs.

At this stage of the inquiry, it was found necessary to adopt arrangements for securing a sudden impact and a sudden removal of light without being obliged on each occasion to extinguish the flame which was the source of the light. For

that purpose we placed over the apparatus a rectangular box, blackened without and within, having a draw-shutter as seen in the diagram, Plate X. After placing the eye on the cushions the key was opened, and the amount of deflection observed. If not sufficiently great for experimental purposes, the eye was readjusted so as to secure accurate contact of the nerve with its corresponding cushion; if sufficiently great, the box was placed over it, and the eye was allowed for several minutes to remain in the dark. The impact of light took place on withdrawing the shutter. The change in the electro-motive force was noted at the instant of the impact. The subsequent change in the electro-motive force produced by continuance of light was not noted till the spot on the galvanometer scale became tolerably steady. When this occurred the shutter was closed, and we noted the effect of the removal of light. In these experiments, therefore, time was not especially taken note of.

SERIES I.—EYE OF FROG.*

The light from a magnesium lamp was passed through a glass vessel, containing a thickness of $1\frac{1}{4}$ inch of water, with the following result:—

Observations.	Before impact.	On impact.	During light.	After removal of light.	Effect of impact.	Effect of removal.
75	347	354	350	358,346	Rise of 7.	Rise of 8.
76	346	354	350	357,345	Rise of 8.	Rise of 7.
77	345	{ Went off } the scale. }	—	—	—	—

Two hours afterwards the following experiments were made with the same eye:—

Obs. 78. Galvanometer at 215. The eye nearly in the dark. On removing the paper from the galvanometer lamp, which was used as a shade, the electro-motive force at once rose to 238. During the continuance of light the electro-motive force fell to 225. On lighting the gas in the room it rose to 238, and then slowly fell during the continuance of light. On shutting off all the light it rose immediately to 230, and then fell to 205.

Obs. 79. Stationary at 205. Gas lit, rose to 224. In two minutes fell to 217, and in one minute more to 213. On turning off the gas, fell from 212 to 215, and then quickly to 197.

Two hours still later the following experiments were made with the same eye:—

Obs. 80. Stationary at 98. Gas lit in room, rose to 105, that is, a rise of seven degrees. During continuance of light, fell to 103. On removal of light, rose one degree.

* In the following series of experiments the zero, polarity, and total first deflection, are not given. These points, however, were always attended to, and observations were always made on a deflection amounting to, at least, 300 or 400 degrees.

Obs. 81. Stationary at 100. On impact of diffuse light, rose to 104, that is, a rise of four degrees. During continuance of light, fell to 102. On turning the light off, a slight and barely perceptible rise was observed.

SERIES II.—EYE OF FROG.

The eye having been adjusted on the clay cushions, the key was opened, and the spot of light reflected from the mirror of the galvanometer was at once deflected three inches off the galvanometer scale. Counting from the point of the polarity, this indicated a deflection of nearly 500 degrees. A small scale, divided into tenths of a millimetre, was attached to the galvanometer scale. The following observations were then made :—

1. A diffuse beam of light from a gas flame was passed through a globular glass jar, twelve inches in diameter, filled with a solution of ammoniacal sulphate of copper (blue light).

Observations.	Before impact.	On impact.	During light.	After removal of light.	Effect of impact.	Effect of removal.
82	57·2	59	56	58	Rise of 1·8	Rise of 2
83	56	58	55	59	Rise of 2	Rise of 4
84	55	56·5	54	55·5	Rise of 1·5	Rise of 1·5
85	55	56	55	58	Rise of 1	Rise of 3
86	58	59	56	57	Rise of 1	Rise of 1
87	52	53	52	54	Rise of 1	Rise of 2
88	51	52	50	53	Rise of 1	Rise of 3
89	51	51·5	49·5	50·2	Rise of ·5	Rise of ·7
90	45	46	47	48	Rise of 1	Rise of 1
91	46·5	47	46·2	47·5	Rise of ·5	Rise of 1·3

In all of the above experiments what we term the inductive effect, that is, the rise in the electro-motive force after removal of the light, was well marked. The impact produced a rise of about two degrees; there was a falling off during the continuance of light, and another rise on the removal of light.

2. In six experiments (*Obs.* 92 to 97) a lighted match, held up at a distance of two feet, produced a rise on impact of about one degree, a fall during continuance of light, and another rise, or inductive effect, of one degree on the withdrawal of light.

3. In all the previous experiments the glass of the galvanometer lamp was surrounded by a black shade, so as to prevent diffuse light from acting on the eye. It was now found (*Obs.* 98) that simply removing this protective shade caused at once a rise of 1·5 degrees; during the time the diffuse light acted on the eye there was a fall of about two degrees, and on again replacing the shade there was a rise of nearly two degrees.

4. The following experiments were then made by acting on the eye with

light after it had passed through a solution of bichromate of potash (red light) :—

Observations.	Before impact.	On impact.	During light.	After removal of light.	Effect of impact.	Effect of removal.
99	37	39	38	40	Rise of 2	Rise of 2
100	38	40	39	40	Rise of 2	Rise of 1
101	35·5	37·2	35·5	36·5	Rise of 1·7	Rise of 1
102	32	33·5	31	31·4	Rise of 1·5	Rise of ·4
103	36	38	34·5	35	Rise of 2	Rise of ·5
At this stage the electro-motive force fell so as to permit readings to be made on the true galvanometer scale.						
104	340	360	350	355 and sudden fall	Rise of 20	Rise of 5
105	270	283	260	268 do.	Rise of 13	Rise of 8
106	235	240	235	238 do.	Rise of 5	Rise of 3
107	242	248	245	248 do.	Rise of 6	Rise of 3
108	235	240	238	239 do.	Rise of 5	Rise of 1
109	224	231	224	225 do.	Rise of 7	Rise of 1
110	217	224	217	219 do.	Rise of 7	Rise of 2
111	209	215	207	208·5 do.	Rise of 6	Rise of 1·5

These experiments show a rise on the impact of light, a fall during the continuance of light, and another rise, or inductive effect, on the withdrawal of light.

SERIES III.—EYE OF FROG.

On placing the eye on the cushions, and opening the key, a deflection of nearly 500 degrees of the galvanometer scale was obtained. The readings were made on the attached scale just alluded to. The following observations were then made :—

Obs. 112. A platinum spatula was heated to a red surface in the flame of a Bunsen burner at a distance of $2\frac{1}{2}$ feet. Before it reached a red heat the galvanometer stood at 47. During the emission of red rays it slowly rose to 49. When spatula cooled, it fell to 48.

Obs. 113 and 114. A little carbonate of soda was placed on the spatula, and introduced into an almost colourless Bunsen flame. Rose from 48 to 51, and on withdrawal sank to 48. This experiment was repeated.

Obs. 115. A little nitrate of strontia was placed on spatula, and pushed into flame. Rose from 48 to 49·5. No inductive effect was observed.

Obs. 116. Experiment with nitrate of strontia was repeated. Rose from 50 to 53·5, and on removal sank to 50.

Obs. 117. Again repeated. Rose from 50 to 53. On removal rose from 52 to 54.

Experiments on the Eye of a Frog with a sudden Flash of Light.

A non-luminous Bunsen flame was held at a distance of $2\frac{1}{2}$ feet, and it was rendered instantaneously luminous by quickly placing the finger and thumb over the two air-holes at the base. The following are the results.—

Observations.	Before the flash.	After the flash.	Rise.
118	51	53.5	2.5
119	53	55	2
120	53.2	56	2.8
121	53	56	3
122	53	55.5	2.5
123	53	55	2

The Bunsen flame was removed to a distance of eight feet, and the experiment was repeated.

Observations.	Before the flash.	After the flash.	Rise.
124	53.2	55	1.8
125	53.2	54.8	1.6
126	53.2	54.6	1.4
127	53.2	54.4	1.2
128	53.8	54.4	.6
129	53.9	55	1.1

A series of observations of a similar nature to those just detailed were made on one of the eyes of other five frogs with exactly similar results; and it appears to us it would serve no useful purpose to give these in detail.

Experiments on the Action of Diffuse Daylight.

A number of observations were now made on the action of daylight on the eye of a frog. In these experiments diffuse daylight was admitted into the totally dark room by removing a piece of cardboard from one of the panes of the window. The following were the results :—

Observations.	Before impact.	On impact.	During light.	Removal of light.	Effect of impact.	Effect of removal.	Remarks.
130	85	100	100	105	Rise of 15	Rise of 5	* Eye readjusted on pads. Gradual rise of the electro - motive force during this experiment.
131	135	140	140	142	Rise of 5	Rise of 2*	
132	214	220	225	228	Rise of 6	Rise of 3	
133	230	240	240	250	Rise of 10	Rise of 10	
134	235	250	250	255	Rise of 15	Rise of 5	
135	274	285	285	290	Rise of 11	Rise of 5	
136	292	300	305	307	Rise of 8	Rise of 2	
137	310	326	326	328	Rise of 16	Rise of 2	
138	314	330	332	340	Rise of 16	Rise of 8	
139	325	338	338	342	Rise of 13	Rise of 4	

These experiments show that diffuse daylight produces results similar to those obtained by the action of rays from artificial luminous sources.

The action of light on the eye of the frog may be briefly stated as follows :
When diffuse light is allowed to impinge on the eye of the frog, after it has

arrived at a tolerably stable condition, the natural electro-motive power is in the first place increased, then diminished; during the continuance of light it is still slowly diminished to a point where it remains tolerably constant; and, on the removal of light, there is a sudden increase of the electro-motive power nearly up to its original position.

IV.—EXPERIMENTS AS TO THE ACTION OF THE VARIOUS RAYS OF THE SPECTRUM.

With the view of observing the action of the various rays of the spectrum, we attempted, in the first instance, to place the eye in different parts of the spectrum produced by interposing a prism in the path of a beam from a lime light. We made numerous observations in this way. Another, and more satisfactory method was to pass light through a rectangular glass vessel, about one inch in thickness, containing a coloured liquid, the absorptive powers of which had been previously determined by the spectroscope. Both of these methods produced substantially the same results, which will be seen in the following record of observations made on the eye of a frog, which was in the condition giving a rise on the impact of light and a fall on its removal:—

Observations.	Beam passed through.	Absorptive effect of fluid on the Spectrum.	Effect of light on the Eye.				
			Before impact.	On impact.	Removal of light.	Effect of impact.	Effect of removal.
140 } 141 } 142 }	Water.	None.	200	220	200	Rise of 20	Fall of 20
			200	220	200	Rise of 20	Fall of 20
			210	225	220	Rise of 15	Fall of 5
143 } 144 } 145 }	Bichromate of potash.	{ Absorbs all beyond the low green— <i>red light.</i>	220	230	222	Rise of 10	Fall of 8
			220	230	220	Rise of 10	Fall of 10
			220	230	220	Rise of 10	Fall of 10
146 } 147 } 148 }	Chrome alum.	{ Absorbs all the yellow, orange, and a portion of the low green, leaving low red and violet— <i>rose tint.</i>	226	230	227	Rise of 4	Fall of 3
			228	232	228	Rise of 4	Fall of 4
			226	230	227	Rise of 4	Fall of 3
149 } 150 } 151 } 152 }	Neutral chromate of potash.	{ Absorbs all above the high green— <i>yellow light.</i>	230	238	225	Rise of 8	Fall of 13
			220	231	221	Rise of 11	Fall of 10
			221	232	221	Rise of 11	Fall of 11
			221	232	220	Rise of 11	Fall of 12
153 } 154 } 155 } 156 }	Ammoniacal sulphate of copper.	{ Absorbs all below the high green— <i>blue light.</i>	235	239	234	Rise of 4	Fall of 5
			237	241	236	Rise of 4	Fall of 5
			240	244	239	Rise of 4	Fall of 5
			241	246	241	Rise of 5	Fall of 5
157 } 158 } 159 } 160 }	Sulphate of nickel.	{ Low and high end of spectrum absorbed — <i>green light.</i>	246	250	246	Rise of 4	Fall of 4
			244	250	246	Rise of 6	Fall of 4
			249	260	250	Rise of 11	Fall of 10
			250	261	251	Rise of 11	Fall of 10
161 } 162 } 163 }	Water, second time.	None.	256	270	257	Rise of 14	Fall of 13
			255	270	255	Rise of 15	Fall of 15
			255	269	255	Rise of 14	Fall of 14

These experiments show—(1), that the part of the spectrum which to our consciousness is the most luminous, namely, the yellow rays, produce the greatest effect in altering the electro-motive force of the retina ; (2), that if we divide the spectrum into two parts, as, for example, into one portion above the high green, and another portion below the high green, the summation of the effects produced is about equal to that of white light ; and (3), that certain of the colours of the spectrum may be arranged in the following order, in their greater relative power in altering the electro-motive force—yellow, green, red, blue.

V.—CRUCIAL EXPERIMENT AS TO THE EFFECTS OF LIGHT ON TWO EYES PLACED ON THE GALVANOMETER CUSHIONS IN REVERSED POSITIONS.

Two eyes of the same frog were dissected out in the ordinary way. The first, which we will call No. 1, was so placed on the cushions as to give a deflection of nearly 400° to the right (the observer being supposed to be seated with his back to the galvanometer and his face to the scale). The other eye, No. 2, was placed in the reversed position. A current was thus obtained from No. 2, which passed through the galvanometer in the opposite direction to that of No. 1 ; but as the current of No. 1 was stronger than that of No. 2, the deflection of No. 1 was only considerably reduced. It is evident that we could now study the effect of light on No. 1 by placing a small black paper cap on No. 2, and *vice versâ*. The effects were what might have been anticipated. When one eye was thrown out of gear by not being subjected to the action of light, the deflection produced by the electro-motive force of the other eye was altered in the usual way. Suppose, for instance, that the impact of light in both cases produced (as in this instance it did) an increase in the electro-motive force, we would expect that when No. 2 was covered, the spot of light on the galvanometer scale would go to the right (as indicated by the rise of numbers in the following figures) ; whereas, if No. 1 was covered, the spot of light would go to the left (as indicated by the fall of numbers in the following figures). That is to say, when No. 2 was covered, and No. 1 acted, light caused an increase in the electro-motive force of No. 1 (to the right) ; whereas when No. 1 was covered, light caused an increase in the electro-motive force of No. 2 (to the left). Such was the case.

Paper cap on No. 1, the deflection of which is to the right. No. 2 was in action.

Obs. 164	Before light, 165	Impact, 163	Continued, 153	Removal of light, 150
" 165	" 175	" 173	" 165	" 160
" 166	" 180	" 177	" 167	" 165
" 167	" 175	" 173	" 163	" 160
" 168	" 176	" 174	" 165	" 160

Paper cap on No. 2, the deflection of which is to the left. No. 1 was in action.

Eyes readjusted on the pads.

Obs. 169	Before light, 210	Impact, 212	Continued, 205	Removal of light, 205
" 170	" 190	" 198	" 190	" 191
" 171	" 178	" 185	" 178	" 182
" 172	" 175	" 180	" 173	" 178
" 173	" 168	" 174	" 168	" 173

VI.—EXPERIMENTS WITH VARYING LUMINOUS INTENSITY.

For this purpose the eye was carefully guarded from extraneous light, as previously detailed, and the source of light was allowed to act on the eye at different distances. The following experiments were made on an eye of a frog that had been removed from the body two hours previously, and had been during that time lying on the clay pads of the electrodes :—

a. *Obs. 174 to 178.*—The gas flame was placed at a distance of *one* foot. Rose eight degrees ; fell eight degrees below original point during action of light ; and rose twelve degrees on removal of light. This was repeated five times with the same results.

b. *Obs. 179 to 184.*—The gas flame was now placed at a distance of *four* feet. The result was that on impact, rose five degrees, during continuance of the light fell five degrees below the original point, and there remained stationary. On removal of light rose to the original point. This happened in six experiments.

c. *Obs. 185 to 188.*—The gas flame was then placed at a distance of *eight* feet. Rose one degree ; fell during continued light five degrees below original point ; and rose five degrees on light being removed. This was repeated four times with same result.

From these observations it results that when the luminous intensity varies in the proportion of 64 to 1, the electrical variation is as 8 to 1, showing that it by no means varies directly as that of the luminous source, but much more slowly. At various times we have made observations on the diminution of electrical effect from a difference of luminous intensity in the proportion of 100 to 1, and as a mean of these observations we have found the electrical variation in different eyes to lie between proportions of 3 to 1 and 6 to 1. That is to say, if a luminous source, at a distance of ten feet, gives an initial impulse corresponding to one degree of our galvanometer scale, then at a distance of one foot the electrical effect would be from three to six degrees. In this comparison, we take the alteration produced in the galvanometer on the impact of light as the measure of the electrical variation, because the eye in this case is always compared under similar conditions, having been previously for a certain time in the dark, so as to allow it to recover from the effect of a previous experiment, whereas if we regarded the electrical effects produced on the withdrawal of light as comparable, we would have to deal with different degrees of

exhaustion of the retina and the action of light during different times. All such questions are left for examination in a subsequent paper on the effects of fatigue. Similar results to the above have been found for the compound eye. The natural query then arises, Are the physical effects we have described and measured really comparable in any way with our sensational differences in light perception when we eliminate all mental processes of association, &c., and leave only perception of difference of intensity? In other words, are these changes representative of what is conveyed to the sensorium? It would appear at first sight that this problem is altogether beyond experimental inquiry. There is, however, a way of arriving at very accurate measures of the variation of our sensational differences in the case of light, and this has been developed theoretically and experimentally by the justly renowned German physiologist FECHNER, and more recently extended and verified by DALBŒUF of the Belgian Academy. FECHNER's law as applied to light may be stated as follows:—If we call I and S respectively the luminous and sensational intensities, and a and k constant quantities, then the following equation is found to express the relation of the above quantities, viz. :—

$$S = a \log I + k \quad . \quad . \quad . \quad (1).$$

From this it follows, if we take any two new values, S_1 and I_1 , we get the following relation—

$$S - S_1 = a \log \left(\frac{I}{I_1} \right) \quad . \quad . \quad . \quad (2).$$

That is, the difference of our sensations is directly proportional to the logarithm of the quotient of the luminous intensities. The formula (1) is, however, defective when considered with regard to small values of I , and gives an imaginary quantity when it becomes zero, viz., infinity. In order to remedy this defect, DALBŒUF has introduced a constant quantity along with the value of the luminous intensity, that may be regarded as representative of the normal condition of the retina in darkness, or the proper light of the eye. Not that there is a certain amount of light when the eye is kept in the dark, only that the receiving organ has a natural susceptibility that must be included in the equation.

The expression (1) becomes when we insert this constant quantity, C ,

$$S = a \log (C + I) + k \quad . \quad . \quad . \quad (3).$$

And as S must be nothing when I is zero, we have—

$$S = a \log \left(\frac{C + I}{C} \right) \quad . \quad . \quad . \quad (4).$$

This formula has been verified experimentally by DALBŒUF, using the method of successive contrasts introduced by PLATEAU, and from it he has succeeded in deducing the value of the unknown constant C . The value of this constant is found in his experiments to vary at different times between the values 0.1 and

0.5. Introducing these values in equation (4), we can find what is the ratio of the sensational intensities for any assigned values of the luminous intensity. In this way, if the amount of light varies as 100 : 1, then the corresponding sensational values may differ as 3 : 1, or 5 : 1. Contrasting the above values with the electrical variations previously given for the same difference of luminous intensity, we observe a close agreement that can scarcely be regarded as accidental. It would thus appear that the psychophysical law of FECHNER is not dependent alone on perception in the brain, but in part on the structure of the eye itself. The effects which occur on, during, and after the action of light on the retina also take place after the eye has been removed from all connection with the brain. Thus the law of FECHNER is not a function of the brain alone, but is really a conjoined function of the brain and the terminal organ, the retina.

VII.—EXPERIMENT ON THE EFFECT OF MOONLIGHT.

Although in previous experiments we endeavoured to eliminate the effects of heat from those of light, we thought it advisable to note the effect of moonlight, which is well known to contain heat rays so few in number as to require the most sensitive arrangement for their detection. The experiment was accordingly made on a clear moonlight night. The troughs, with the eye placed on the cushions, were carried into the open air, and were connected by long wires with the galvanometer. It is sufficient to state that in these circumstances we found the eye as sensitive to moonlight as to the light of a candle placed at a distance of five or six feet.

VIII.—EFFECTS OF CERTAIN ACTIVE SUBSTANCES ON THE SENSIBILITY OF THE RETINA.

The subcutaneous injection into the frog of lethal doses of woorara, santonin, belladonna, morphia, and Calabar bean, does not destroy the sensibility of the retina to light. The change in the electro-motive force was not altered quantitatively to such an extent as to be appreciable. This is most remarkable in the case of santonin, which is well known to produce in man sensations of yellow light.

IX.—EXAMINATION OF OTHER TEXTURES OF THE EYE.

It is manifest that in an investigation such as this, it is of great importance to examine the action of light on the various textures or tissues composing the globe of the eye and the adjacent skin, so as to determine whether or not these are affected by light.

1. *The Skin*.—As has already been shown by DU BOIS-REYMOND and many others, there is no difficulty in obtaining a strong deflection when the clay points are applied to the skin of the frog, one to the inner and the other to the outer surface. From the well-known experiments of LISTER on the pigment cells in the skin of the frog, in which he showed that the pigment molecules changed

position under the action of light, it might possibly have been expected that there would be a change in the electro-motive force. It has been found, however, in this research, that this current is not affected by light.

2. *Pigment Cells of the Choroid*.—A current is readily obtained from the pigment cells of the choroid, when a portion of this membrane is laid on the clay points. This current does not exhibit any sensitiveness to light.

3. *Sclerotic and Nerve without the Retina*.—These structures give a considerable current not affected by light.

4. *Anterior Segment of the Eyeball*.—The eye of a frog was bisected so as to include in the anterior segment the cornea, iris, and lens. This preparation was then placed on the pads, so that the corneal surface was in contact with the one pad while the posterior surface of the lens was in contact with the other. A strong deflection was obtained, which was, however, not affected by light.

5. *Posterior Segment of the Eyeball*.—On bisecting an eye of a frog, so as to remove completely the anterior portion, including cornea, aqueous humour, iris, ciliary muscle, and lens, and on bringing the retina into actual contact with one of the clay pads, we obtained a large deflection, which was as sensitive to light as when the whole eye was employed, thus eliminating any possibility of contractions of the iris under the stimulus of light, having to do with the results previously obtained.

6. *Distribution of the Electro-motive Force of the various Textures of the Eye*.—The distribution of the electro-motive force between the different portions of the eye and cross section of the nerve may be stated as follows:—The most positive structure is the cornea, then the sclerotic, then the longitudinal surface of the nerve; the cornea is also positive to the posterior surface of the crystalline lens, and the retina itself is positive to the transverse section of the nerve.

X.—EXPERIMENTS ON THE EYES OF LIVING BIRDS AND MAMMALS.

At this stage of the inquiry, we examined the action of light on the eyes of living animals. In our earlier experiments on the eyes of rabbits, cats, and dogs, we found that sensibility to light disappeared with great rapidity. This phenomenon is in conformity with the well-known fact, that the sensibility of the nervous system in warm-blooded animals quickly disappears after death from loss of arterial blood. It was important to ascertain, if possible, whether or not the variations in the electro-motive force, by the action of light, which we had noticed in the eyes of various animals removed from the head, also occurred in the eye while it remained in the head, and was nourished by blood. We, accordingly, instituted a series of experiments which were practically very troublesome. We examined the eye—(1), of the living cat; (2), of the living pigeon; and (3), of the living owl (*Strix flammea*, Lin.) In all cases the animals were deeply under the influence of chloroform during the experiments.

1. *The Cat*.—The animal was securely fixed in CZERMAK'S rabbit-holder. The skin around the orbit was reflected. The zygomatic arch was snipped through by bone forceps, so as to expose as much as possible of the side of the orbit. The cellular tissue of the orbit was then pushed aside along the superior and the lateral aspect of the eyeball, so as to reach the optic nerve with as little disturbance as possible to the vascular arrangements of the eyeball. On exposing clearly the optic nerve, and staunching hæmorrhage, the nerve was cut through transversely with sharp scissors. When this was done, the globe could be pulled downwards, inwards, and forwards, so as to expose a clear transverse section of the nerve. With the head firmly fixed, one narrow clay point was now placed on the cornea, while the other was in contact with the transverse section of the nerve. On opening the key a strong deflection was obtained, in several instances so powerful as to drive the spot of light off the scale. In one experiment it was for a long time impossible to obtain sufficiently slight contact to procure a deflection small enough to lie on the galvanometer scale. In several experiments we succeeded in tracing the curve made by the movement of the spot of light on the surface of a horizontal cylinder, and it will be seen, on referring to this curve, as shown in the annexed plate (Plate XI.), that on the impact of light the electro-motive force diminished, during its continuance the electro-motive force gradually rose to a point where it became steady, and on the withdrawal of light the inductive effect, that is, a rise in the electro-motive force, was well marked.

2. *The Pigeon*.—The eye and optic nerve of the pigeon were more accessible than those of the cat, and the head of the bird was more easily held in a stable condition. The eye was found sensitive to light, and followed the same law as that of the cat.

3. *The Owl*.—As this bird is believed to be capable of seeing objects with a very feeble intensity of light, we considered it important to examine its eye in the same manner as we examined that of the pigeon. We found exactly the same results. We could not detect any quantitative differences between the effects of light on the eye of the pigeon and on the eye of the owl.

XI.—EXPERIMENTS ON THE EFFECT OF LIGHT, IN WHICH A PORTION OF THE BRAIN WAS INCLUDED IN THE CIRCUIT.

A number of experiments were made with the view of determining how far into the brain substance it was possible to observe any electrical variation produced by the action of light on the retina, the optic nerve and the eyeball remaining in their natural position.

1. *The Frog*.—On bisecting the head of a newly killed frog with a sharp pair of scissors, it is possible to obtain a longitudinal section of the brain, with

the various parts in their natural position, then to carefully cut away the anterior and posterior portions of the brain, leaving only the middle portion in contact with the optic nerve. On placing this preparation between the clay points, so that the one touches the surface of the cornea, while the other is in contact with the brain substance, a strong deflection is obtained which is sensitive to light, and follows the course observed in the case of the frog.

2. *The Pigeon*.—The effect was also traced into the optic lobes of a living pigeon (deeply under chloroform), the head of which was securely held between the clay points of the electrodes. The optic lobes in the pigeon are easily exposed. The following were the effects of this observation:—*a*, When one pole was applied to the left optic lobe, and the other to the cornea of the right eye, a deflection was obtained which was sensitive to light; *b*, when the pole was removed from the right eye, and applied to the cornea of the left, a smaller deflection was obtained, also sensitive to light; and, *c*, when light was allowed to impinge on both eyes, while the one pole was in contact with either eye, and the other with the left optic lobe, the result was nearly double that produced by the impact of light on one eye alone, either right or left. These effects may be explained by the decussation of the optic nerves in the optic commissure.

XII.—EXPERIMENTS ON THE EYES OF FISHES.

The subject was examined in the case of three fishes—(1), the gold fish (*Cyprinus auratus*); (2), the rockling (*Motella vulgaris*); and (3), the stickleback (*Gasterosteus brachurus*).

1. *The Gold Fish*.—With the eye of this fish a deflection of about 600° was obtained with the smallest possible contact. On the impact of light it rose 50° , remained steady for a short time, and then slowly sank. On removal it fell about 8° or 10° below the starting point.

2. *The Rock Fish*.—Specimens of this fish were obtained from the rock-pools in the neighbourhood of Kinghorn in Fifeshire. The deflection was similar in amount to that of the gold fish, but the variations occurred in a much longer period of time.

3. *The Stickleback*.—The species examined is that frequently met with in brackish pools, a little beyond the level of high-water mark, at the sea-side. The eyes examined were very small. A deflection of 400° was obtained, which was remarkably sensitive to light, the variations occurring in very short periods of time.

On comparing these results, it will be seen that, in the active and alert stickleback, the variations occurred rapidly; in the more sedate goldfish, they occurred more slowly, while, in the case of the sluggish rockling, which hides beneath stones near low-water mark, the variations occurred very slowly.

XIII.—EXPERIMENTS ON THE EYES OF AMPHIBIA AND REPTILES.

Observations have been made on—(1), the frog (*Rana temporaria*); (2), the toad (*Bufo vulgaris*); (3), the newt (*Triton aquaticus*); and (4), a snake (species unknown). Those made on the frog have been already fully described. The action on the eye of the toad was found to be exactly similar to that of the frog, only the variations were much more sluggish. In the case of the newt, the eye was found to be sensitive to light to the extent of about 12 per cent. of the electro-motive force. It was found also to be sensitive to a flash of light. These results were obtained in an eye having the following dimensions :—

Transverse diameter of eye, $\frac{1}{8}$ th of an inch.

Diameter of the pupil, $\frac{1}{30}$ th of an inch.

Diameter of the lens, $\frac{1}{20}$ th of an inch.

The eye of the snake* resembled that of the frog.

XIV.—EXPERIMENTS ON THE ACTION OF LIGHT ON THE COMPOUND EYE.

We have examined the eyes of the following crustaceans, namely—(1), the common edible crab (*Cancer pagurus*); (2), the swimming crab (*Portunus puber*); (3), the spider crab (*Hyas coarctatus*); (4), the hermit crab (*Pagurus Bernhardus*); and (5), the lobster (*Homarus vulgaris*).

1. *The Common Edible Crab*.—In this case it was found that the nerve surface was positive to the other parts of the eye, not negative, as had been shown in the simple eye of various vertebrate animals. A deflection was obtained to the extent of about 300° of the scale of the galvanometer. The effect of light was to increase the current from 5° to 10° of the galvanometer scale, then to fall slightly. During the action of light the deflection remained very permanent, and on the withdrawal of light it fell.

2. *The Lobster*.—With this eye the following points were noted :—

(1.) The direction of the current was reversed, as in the case of the crab, that is to say, the corneal surface was negative, not positive, as was observed in the simple eye.

(2.) A deflection of about 600° of the scale were obtained with the smallest possible contact of the clay points with the cornea and nerve surface.

(3.) The action of light was to diminish the current to the extent of 10 per cent of the total amount of deflection, and the inductive effect, or effect of removal of light, was nearly equal to the depression produced by the impact of light.

* Kindly sent us by Mr BARTLETT of the Zoological Gardens, Regent's Park. Genus and species not given.

(4.) It was found in this eye that at a distance of one foot, on the impact of light, there was a depression of 90° to 103° ; at a distance of ten feet, there was a depression of from 15° to 20° . Here the depression did not at all correspond to the intensity. At a distance of ten feet, when the amount of light was $\frac{1}{100}$ th part of what it was at one foot, the diminution in the alteration of electro-motive force by the action of light was not $\frac{1}{100}$ th, but $\frac{1}{5}$ th of the total amount.

3. *The Swimming Crab*; and, 4, *the Spider Crab*.—These animals, dredged from a depth of twenty fathoms from the Firth of Forth, had eyes which were found sensitive to light.

5. *The Hermit Crab*.—The length of the eye examined in this case was about one millimetre. Its breadth was about one-fourth of a millimetre. Still it was found very sensitive to light.

XV.—EXPERIMENTS ON THE EFFECT OF FATIGUE ON THE EYE.

A few experiments, preliminary to a more detailed examination of the question, have been made as to the effect of fatigue on the eye, as estimated quantitatively by the amount of change in the electrical potential of the visual structures. We may, in this communication, shortly allude to these. Suppose we place the eye of a frog in connection with the clay pads of the troughs of the galvanometer. The box with the draw-shutter, already described, is placed over it, and one candle, giving a steady flame, is put a distance of twelve to sixteen inches from the box. The draw-shutter is now opened, and the amount of deflection on impact of, during the continuance of, and on the removal of, light is noted. If we then light another and exactly similar candle, so as to double the amount of light, the variation in the electro-motive condition of the eye is not much altered, certainly not to the extent of double the amount of that produced by one candle. If we then extinguish one candle, a slight alteration occurs; but when we extinguish the remaining candle, there is at once a great variation in the deflection of the galvanometer. That is to say, the eye becomes less sensitive to any increase in the intensity of light after it has become for some time subjected to the action of a light of certain intensity; or, in other words, it becomes fatigued. This experiment also shows that the eye is more sensitive to variations in light of weak intensity than to variations in light of great intensity.

XVI.—MODES OF GRAPHIC REGISTRATION IN THESE EXPERIMENTS.

During the course of our observations we have found it necessary to construct a true graphical representation of the variations of the electro-motive force occasioned by the impact, duration, and removal of light. It is clear that to register minute galvanometrical alterations from a THOMSON'S instrument, the only plan

that could be employed, yielding perfectly accurate results, would be to photograph on a sensitive surface, covering a cylinder rapidly revolving on a horizontal axis, the alteration of position of the spot of light reflected from the mirror, just as continuous magnetic observations are registered. As the apparatus required to execute these observations is very complicated, and will only become necessary in the second part of this memoir, where we must deal with the time occupied in the transmission of retinal impressions due to different coloured rays, we have, in the meantime, adopted two simple methods of registration. The first plan is to note the position of the galvanometer at equal intervals of time before, during, and after the impact of light on the eye. In these observations we have used a seconds pendulum giving a loud beat. One observer reads aloud the galvanometer, the other marks every interval of two seconds, registers the numbers obtained, and regulates the supply of light. A little practice in the method above described has enabled us to obtain very satisfactory results, agreeing very closely in different observations, and showing in a decided way the salient points of the variation curve (see curves, Plate XI.). The second plan of registration which may be of service in many physical and physiological researches, consists in placing at proper distances from the galvanometer, instead of the ordinary graduated scale, the surface of a cylinder covered with paper, and moving on a horizontal axis by clockwork. The spot of light reflected from the galvanometer mirror is rendered more precise by having the shade of the galvanometer lamp blackened over the entire surface, with the exception of a spot about three millimetres in breadth, in the centre of which a line or cross is made of soot. The image of this line or cross is, of course, reflected by the mirror upon the cylinder. When the cylinder is set in motion by the clockwork, the spot of light may be accurately followed by the hand of the observer after a little practice, with a fine brush moistened with ink. The cylinder we employed performed a complete revolution in eighty seconds. This time was divided into four equal parts, each representing twenty seconds, by four lines drawn transversely at equal intervals across the paper on the cylinder. The first space, between lines one and two, represented twenty seconds, in which the eye was in the dark, and in which the electro-motive force is represented by a straight line; the second space, between lines two and three, represented twenty seconds, during which the effect of the impact of light took place, and in which the variation of the electro-motive force is indicated, either by a curve to the right or to the left; the third space, between lines three and four, represented twenty seconds of continued action of light, during which the electro-motive force gradually rises; and lastly, the fourth space between lines four and one (the point of starting), represented twenty seconds, during which the electro-motive force at first rises on the withdrawal of light, and afterwards sinks rapidly. Illustrations of curves registered by

the two methods described above will be found in Plate XI. appended to this paper.

As this paper embodies only the experimental results obtained from a general survey of the subject, leaving many minute and intricate points of difference between the result of the natural stimulus applied to the nervous structure of the eye and the effect of stimulating a motor nerve (that is, what is called a negative variation of the normal nerve current), we do not enter into any theoretical explanations regarding the cause of these peculiarities, but leave all such matters for discussion and further elaboration in the second portion of this paper, on which we are at present engaged. In the meantime we may succinctly summarize the results of this part.

XVII.—SUMMARY OF PART I. ON THE PHYSIOLOGICAL ACTION OF LIGHT.

We have experimentally proved—(1), that the impact of light on the eyes of members of the following groups of animals, viz., Mammalia, Aves, Reptilia, Amphibia, Pisces, and Crustacea, produces a variation amounting to from three to ten per cent of the normal electro-motive force existing between the corneal surface and the transverse section of the nerve ; (2), that this electrical alteration may be traced into the brain ; (3), that those rays that we regard as most luminous produce the largest variation ; (4), that the alteration of the electrical effect with varying luminous intensity seems to follow very closely ratios given by the psycho-physical law of FECHNER ; (5), that the electrical alteration is due to the action of light on the retinal structure itself, as it is independent of the anterior portion of the eye, eliminating, therefore, the natural supposition that the contraction of the iris might produce a similar result ; (6), that it is possible, by experiment, to discover the physical expression of what is usually called in physiological language, fatigue ; and (7), that the method employed in this research may be applied to the investigation of the special organs of the other senses.

[APPENDIX.

APPENDIX.

18th February 1874.

While this paper was passing through the press, we have been fortunate enough to have obtained information from a Swedish lady that a paper relating to the same kind of work had appeared in a Swedish Journal. We, accordingly, endeavoured to obtain this paper, which we find to be one by Professor HOLMGREN, a distinguished physiologist, published in the "*Upsala Lakere-förenings Förhand lingar*," vol. vi. 1870-71, No. 5, p. 419. It is with pleasure we at once acknowledge that he has the claim of priority in observing an electrical fluctuation by the action of light, and that his memoir is a valuable contribution to science. It is, however, almost needless to state that our work was done in entire ignorance of any previous observations on the subject, and that our methods of experiment, the delicacy of our instrument, and the distinct numerical data obtained, have enabled us to prosecute the matter in various new directions.

EXPLANATION OF THE PLATES.

PLATE X.

This Plate shows the arrangement of the apparatus.

A. Galvanometer.

B. Scale.

C. Key.

D. Du Bois-REYMOND's non-polarizable electrodes having the eye placed between the clay points.

E. Moist chamber, consisting of an annular glass vessel of at least two inches section, filled with water, and placed over the electrodes to keep the temperature uniform, and absorb radiant heat. Over this is placed a blackened box, the outlines of which are marked in dotted lines, having a moveable shutter in front, F.

G. Lamp.

PLATE XI.

This plate shows curves representing variations in the electro-motive force of the eye produced by the action of light; or they represent the path pursued at different times by the image reflected from the galvanometer mirror if the plate were supposed to move vertically upwards. The dotted lines represent the normal position of the galvanometer-image in the dark; and the complete dark lines represent the position of the image during and subsequent to the action of light.

I. and II. Actual curves taken from the living eye of the cat on a cylinder moving horizontally.

III., IV., and VI. Graphical curves representing the general action taken by time observations.

III. and VI. Frog. IV. Toad. V., VII., VIII., and IX. Actual curves taken on a cylinder from the eye of a frog.

X. and XI. Curves obtained from eye of cat removed as quickly as possible from dead body.

The four dotted horizontal lines divide the time of the experiment into four equal intervals, namely, 1st, in darkness; 2d, during action of light; 3d, during continued light; and, finally, on removal of light. A movement to the right of the dotted line indicates an increase, and to the left a diminution, in the electro-motive force.