smaller degree) in the genus *Tinamus**, one or two Parrots, and also in some of the Birds of Prey." I have not, however, met with them in any group except the *Ardeæ* and their allies. I can assert most positively that no traces of these patches exist in the Pelicans, Storks, or Cranes. I have also taken considerable trouble with *Scopus*. This bird is considered by Prof. Reinhardt to be closely allied to *Baleniceps*. I cannot find anything to justify such a belief; the skins and skulls of the two birds are so entirely different, that it is useless to enter into any further details respecting them.

There is one thing, however, that I wish to remark, and I do so with considerable uneasiness lest I should be accused of casting a doubt upon the veracity of the gentleman to whom we are indebted for the first living specimens of this rare bird; and this consideration would have prevented my making the remark, had not my great desire been to call attention to the subject in the hope of obtaining a truthful explanation of what appears to me inexplicable—I refer to the statement, made by Mr. Petherick, that *Baleniceps* runs about in search of food immediately after it is hatched. If this is true, it is one of the most extraordinary facts I have yet met with.

April 9th, 1861.

Dr. J. E. Gray, F.R.S., V.P., in the Chair,

The following extracts from a letter addressed to the Secretary by Robert Swinhoe, Esq., Corresponding Member of the Society, dated British Consulate, Amoy, Jan. 1st, 1861, were read to the meeting:—

"I returned a few weeks ago from the north of China, where I was attached to Sir Hope Grant as Interpreter. Through the kindness of Major Sarel of the 17th Lancers, attached as Aide-de-camp to Sir Hope Grant, I was enabled to procure three skins of a Deer found in herds in the Imperial Parks of the Summer Palace. The three deer are apparently first, a two-year-old buck; the second, a one-year-old buck; and the third a doe. I skinned them, and preserved their skulls. They are, I think, *Cervus wallichii* of Pallas, from Eastern Siberia; and I trust will form an acceptable acquisition to the collection of the Zoological Society. Major Sarel also procured the heads of three fine bucks, which he has sent to his brother-in-law, Edmund Bury, Esq., of Manchester, and which will shortly be

* Since writing the foregoing I have examined *Tinamus*. The structure referred to by Nitzsch appears to differ so widely from the down-patches of the Ardeine family, that I shall describe it in another paper upon this subject, which I hope to have ready shortly.

J. Wolf, lith.

M & N. Hanhart, Imp

HYLOBATES PILEATUS

put into the hands of Mr. Leadbeater for preservation. If, therefore, the Society are anxious to compare the skins I send with the heads of older and finer animals, they will find no difficulty in so doing by applying to Mr. Leadbeater.

"I send also in the same box the skin of a Mole (perhaps *Talpa leucura*, Blyth) from Pekin.

"A Hedgehog, which I take to be new, is also from Pekin. But this species is also found in Shanghai, and has lately been captured at Swatow.

"In a bottle, preserved in spirits, you will find a small Mouse and a Bat, besides several Snakes, Lizards, and Frogs.

"All these specimens have been fastened in a tinued box, and shipped on board the 'Harkaway' bound for London. I enclose you the mate's receipt."

Extracts were also read from a letter addressed to Mr. T. Rupert Jones by Mr. S. R. Pittard, containing remarks relative to the flight of Albatroses (*Diomedea*) and other oceanic birds, as observed during his voyage to Australia.

The following papers were read :—

1. LIST OF MAMMALIA, TORTOISES AND CROCODILES COLLECTED BY M. MOUHOT IN CAMBOJA. BY JOHN EDWARD GRAY, F.R.S., V.P.Z.S., F.L.S.

(Plate XXI.)

M. Mouhot having sent to Mr. S. Stevens a number of Mammalia and other animals from Camboja, a series of them has been secured for the British Museum Collection.

I hasten to lay before the Society a list of all the specimens, and the varieties that they present, with a description of the species or varieties which appear to be new to science.

MAMMALIA.

There are eight specimens of the Long-armed Ape (*Hylobates*). They differ considerably in colour, and yet offer some peculiarities which I have not observed in specimens from other parts of India or the Asiatic islands, so that I am induced to believe them to be a separate species or local variety. M. Mouhot is inclined to regard them as varieties or states of the same species; as he observed that they were all found in company on a small island near Camboja.

1. The oldest specimens, whether they are black with a more or less grey back and white hands, or a white with a more or less long black spot on the chest, generally have a large black spot on the crown of the head, circumscribed by a pale edge; some of the white examples have white and others black whiskers.

2. The two younger specimens are yellowish white, without any black, not even the spot on the head; but this may be only on account of their youth.

From this spot on the forehead I am inclined to distinguish this species or variety by the name of *H. pileatus*, though this is not quite a distinctive mark, as both *H. leucurus* and *H. agilis* have the crown more or less black.

HYLOBATES PILEATUS. (Pl. XXI.)

Male.—Black; back of head, back of body, and front hind legs greyish; forehead and circumference of the black spot on the crown paler grey; hands and tuft of long hair round the organ of generation white.

The three specimens in this state are all nearly of the same size, and appear to be adult. They only vary slightly in the size of the coronal spot, and in the extent of the white colour on the hands.

Female.—White; back brownish white, slightly waved; a large ovate spot on the crown, and a very large ovate blotch on the chest, black.

These specimens are all of one size, and appear to be adult, and are probably female; three have the teats well developed. They vary in the size of the black chest spot, and in the colour of the whiskers, thus:—

a—b. White; spot on the chest moderate, reaching only half-way down the abdomen; whiskers on side of face white.

c. Brownish; spot on chest larger, reaching further down the abdomen; sides of the face black; a few black hairs on the throat.

d. Brownish; side of the face, under the chin, and the whole of the throat, chest, and belly black; teats well-developed.

Young.—Uniform dirty white, without any black spot on chest or head.

PTEROPUS EDWARDSII, Geoffroy, Ann. Mus. xv. p. 92.

Hab. Camboja.

A single specimen.

LEOPARDUS VIVERRINUS, Gray, List Mamm. B.M. p. 43, 1843.

Hab. Camboja.

Three specimens.

PARADOXURUS PALLASII, Gray, P. Z. S. 1832, p. 67.

Hab. Camboja.

Two specimens, adult and young.

VIVERRA MALACCENSIS, Gmel.

Hab. Camboja.

A single specimen.

HERPESTES RUTILUS.

Grisled chestnut-brown, variegated with black and white rings on the hairs; the head and limbs darker chestnut, with scarcely any and very narrow white rings; lips and throat, and under part of the

body, uniform duller brown, not grisled; ears brown, the nape with longer hairs, forming a broad short crest.

Hab. Camboja.

A single specimen.

Browner than any other Indian *Herpestes* I have seen.

PTEROMYS MOMOGA?

Hab. Camboja.

One specimen.

This species is very like *Pteromys momoga*, Temm. Faun. Japon. Mam. t. 14.

There are five specimens of a Squirrel in the collection; they appear to be distinct from any we have before received; but they offer four very distinct varieties, and it is not easy to decide if the animal may not itself be a local variety of some other Indian species.

SCIURUS SPLENDENS.

All the specimens are bright red bay.

Var. 1. All over dark and very intense red bay, with a white spot on each side of the base of the tail. Two specimens, nearly identical in colour and size.

Var. 2. Top of the head and tail, like var. 1, dark and very intense red bay; side of the back, under sides of the body, and tip of the tail paler red bay, without any white spot at the base of the tail. A single specimen.

Var. 3. Uniform pale bay, like the side of var. 2; tail and middle of the back rather darker and brighter; tail without pale tip or white basal spot. A single specimen.

Var. 4. Crown, middle of the back and tail dark intense red bay; throat, chest, and under side paler red bay, like vars. 2, 3; cheeks, shoulders and thighs, and outsides of the fore and hind legs brown, grisled with yellow rings on the hairs; side of the body rather greyish red. A single specimen.

This last state would almost lead one to suppose that this Squirrel may be only a variety of some other species.

SCIURUS JAVENSIS, Schreb.

Hab. Camboja.

Three specimens, one male and two females, very nearly alike in size and colouring.

SCIURUS MACCLELLANDII, Horsf. Proc. Zool. Soc. 1839, p. 152.

One specimen.

SCIURUS SIAMENSIS, Gray, Proc. Zool. Soc. 1859, p. 478.

Hab. Siam.

Two specimens, none now sent.

SCIURUS MOUHOTII.

Grisled grey-brown, with pale rings; lips, chin, throat, and un-

der side of body and inside of limbs white; the upper part of the sides with a longitudinal black streak, edged above and below with a narrow white line; tail blackish whitish washed, hairs elongate, brown, with two broad black rings and a white tip; ears simple, rounded.

Hab. Camboja.

A single specimen.

The species differ from most of the Squirrels of the size, in the three streaks being on the upper part of the back, and in the dark colour between the two colours of the upper and under surface.

RUSA PERONII, Gray, Cat. B. M. p. 211.

Cervus peronii, Cuvier.

C. kuhlii, S. Müller.

Hab. Camboja.

There is a pair of shed horns, evidently of an adult specimen, in the collection; they are small and slender, and similar to the horns of *C. kuhlii* in the Museum Collection, which were received from Holland. They may be the horns of an allied but distinct species; for it is quite impossible to distinguish deer by the horns alone, they vary so much in specimens of the same species.

RUSA — ?

Hab. Camboja.

There is a single horn of the right side of the head, which has been cut from the base, which is very distinct from any horn that I know. It is very thick and strong, curved, covered with very large grooves and warts, and with a short, conical, thick branch, bending rather inwards on the front of the inner side quite at the base.

CERVULUS CAMBOJENSIS.

Hab. Camboja.

There are the forehead covered with hair and the horns of a Muntjack in the collection. It is very much larger than any specimen of that genus in the Museum Collection, and is probably a distinct species.

The horns are thick, nearly straight, with a short, thick, recurved branch on the outer part of the front side near the base, and one of them has a somewhat similar callosity on the hinder side on the same level. Hair of forehead very rigid, close pressed, dark brown, with narrow yellow rings.

TRAGULUS AFFINIS, Gray, List of Mamm. Brit. Mus. p. 173, 1843.

Not yet described. Similar to *T. javanicus* in colour, but rather smaller and much paler, and the side of the neck similar in colour to the side of the body; the belly is white, with a brown streak on each side of the central line; the head is smaller. It is larger than *T. kanchil*, very much paler, and the neck is not blacker and grisdled. A specimen of the species has been in the Museum as above

named for many years: it is said to have come from Singapore; but that probably was only the port of transit. It may be only a small pale local variety of *T. kanchil*.

Six specimens, adult, all exactly similar; one young.

Sus — ?

Hab. Camboja.

There are two lower tusks of this genus in the collection, indicating the existence of a species of the genus; one is very large and thick, the other is elongate and slender. They may be the tusks of the two sexes.

MANIS PENTADACTYLA, Linn.

Hab. Camboja.

Two specimens, adult and half-grown.

REPTILIA.

TESTUDO ELONGATA, Gray.

One specimen, half-grown.

The shields are yellow, with an irregular black ring round the areola; the disk of the areola is sometimes varied with a few roundish black spots; the sternal shields are yellow, with an irregular square subcentral blotch covering part of the areola.

GEOCLEMYS MACROCEPHALA, Gray, Proc. Zool. Soc. 1859, p. 478. t. xxi.

Hab. Siam and Camboja.

Two specimens, one adult, other young.

The adult is nearly twice as large as the specimen first described; the three keels are distinct, but very blunt, and the vertebral shields are oblong, transverse, but rounded at the angles; the margin of the shell is yellow; the under side yellow, with a more or less large black blotch on each shield.

The young shell is brown, with a narrow, pale-yellow margin; the keels are very distinct; the central one is very broad, and wider on the hinder parts of each shield; the lateral keel ends with the third costal plate, and is at the end bent in towards the central line; the under-side is dark brown, with a white streak down the centre, and a white streak on the margin; and on the keel, which separates the flat parts of the sternum from the shelving part of the sides, that shelving part is high, much higher than in the adult shell.

GEOEMYDA SPINOSA, Gray, P. Z. S. 1834.

Hab. Camboja.

Three specimens in different ages, from young to nearly adult. They are all marked with radiating brown lines on each of the sternal and the under side of the marginal shields, which are widest in the youngest and narrowest in the most adult specimens. The older specimen alone has these lines on the dorsal and upper side of the

marginal shields; the areolæ being plain brown, as in the younger specimen.

EMYS CRASSICOLLIS, Gray, Cat. Tort. Croc. Brit. Mus. p. 16, 1844.

Hab. Camboja.

Two specimens, adult and young.

As in *Geoemyda macrocephala*, the sternum of the young specimen is rather higher than the adult, and the centre is separated from the side by a distinct marginal keel, which is not so visible in the more adult specimen.

CISTUDO AMBOINENSIS, Gray, Syn. Cat. Tort. Croc. Brit. Mus. p. 30, 1844.

Hab. Camboja.

One specimen, adult, of the high, oblong variety.

The sternum is white, with an irregular black spot on each shield.

CROCODILUS POROSUS, Schneid.

Crocodilus porosus, Gray, Cat. B. M. p. 58.

C. biscutatus, Cuvier, Oss. Foss. v. 65. t. 2. f. 8.

Hab. Camboja.

There is a young specimen of this species, rather more than 4 feet long, in the collection. It has no appearance of the ridges on the centre of the hinder part of the skull, on which *Crocodilus siamensis*, Schl., and *C. galeatus*, Cuvier (Oss. Foss. v. t. 1. f. 9), is founded; indeed it is very doubtful if these were not accidental malformations in the skull figured.

2. REMARKS ON THE BREEDING OF THE LARGER FELIDÆ IN CAPTIVITY. BY A. D. BARTLETT.

(Plate XXII.)

The Lion (*Felis leo*) appears to breed more freely than any other species of *Felis*, and the number of young at a birth is greater, not unfrequently four and sometimes five being produced in a litter. It is remarkable that these animals breed more freely in travelling collections (wild-beast shows) than in zoological gardens; probably the constant excitement and irritation produced by moving from place to place, or change of air, may have considerable influence in the matter.

The Tiger (*Felis tigris*) has rarely bred in confinement; but there are several well-authenticated instances of the female Tiger breeding with the Lion. The hybrids lived, and in due time arrived at maturity.

Animals of this mixed breed have been exhibited in a travelling menagerie kept by Mr. Atkins; the appearance of the animals at once bespoke their mixed origin.

I have more than once met with instances of the male Jaguar (*F. onca*) breeding with a female Leopard (*F. leopardus*): these hybrids also were reared recently in Wombwell's well-known travelling col-

J. Wolf, del. et lith.

M. & N. HARBER, Imp.

FELIS CONCOLOR *Juv*

lection. I have seen some animals of this kind bred between a male black Jaguar and a female Indian Leopard; the young partook strongly of the male, being almost black.

The Leopard is not unfrequently bred in captivity; many have been bred and reared in the Society's Gardens, and elsewhere.

The Puma (*F. concolor*) has bred frequently in the Society's Gardens. This species appears to produce generally two only at a birth, in some cases only one: they rear their young without difficulty.

The Ocelot (*F. pardalis*) has also bred two or three times in the Society's Gardens.

As far as I am able to ascertain, the period of gestation in the foregoing species is *sixteen weeks*. The young of some of the species bear a great resemblance to each other; thus, for instance, the young of the Lion is indistinctly spotted all over; the young of the Puma is also spotted with large and well-marked patches, which the accompanying figures (Plate XXII.), drawn by Mr. Wolf from young Pumas born in the Gardens, well exhibit.

The Cheetah (*Felis jubata*) never to my knowledge has bred in England; Dr. Günther, however, informs me that this species has bred in the Gardens in Frankfort. From all that I have experienced with reference to this beautiful species, I consider it one of the most difficult of the family to keep, and consequently the chances of its breeding are rare. This animal is generally gentle, timid, and very excitable. I am inclined to think the want of sufficient space and exercise, together with over-feeding, are the cause of convulsions and fits, to which this species is liable; I have witnessed the death of two or three that have died from excitement after a full meal.

The young of the Tiger is striped like the adult, but of course less distinctly. The young Leopard also resembles the adult in its markings. Not only in the large Cats, but in all the smaller species of the genus *Felis*, the spots, stripes, or markings are always present in the young of those species that are so marked in the adult state; and, as far as my knowledge extends, the young of all (except the *domestic* cat) exhibit traces of spots or other markings, although they disappear in the adult animals.

A very extraordinary malformation or defect has frequently occurred among the lions produced during the last twenty years in the Regent's Park. This imperfection consists in the roof of the mouth being open—the palatal bones do not meet, the animal is therefore unable to suck, and consequently always dies. This abnormal condition has not been confined to the young of any one pair of lions, but many lions that have bred in the Gardens, and not in any way related to each other, have from time to time produced these malformed young, the cause of which appears to me quite unaccountable.

Since writing the above, I have the pleasure of adding that the Society's collection has been enriched by a litter of cubs from the female Mexican Jaguar (*Felis hernandezii*) and the male of the common Jaguar (*Felis onca*). I may add at the same time, that the Tigress in the Society's Gardens is now in young.

3. ON A NEW SPECIES OF THE FAMILY BOIDÆ.
BY DR. ALBERT GÜNTHER.

(Plate XXIII.)

PELOPHILUS FORDII.

Head rather narrow; neck slender; body thick, compressed; tail conical, tapering, prehensile. The anterior part of the head is covered with regular shields, symmetrically arranged, the posterior part from the orbits is scaly. There are four pairs of frontal shields, two small shields being intercalated between the posterior pair; then follows a large square vertical shield, the largest of all the shields of the head, situated between the superciliaries. The nasal opening is small, between three shields—namely between the two nasals and the anterior frontal. Three or four loreals, two anterior and five posterior oculars. Thirteen upper labials, the sixth, seventh and eighth entering the orbit; none of them grooved. Pupil vertical, elliptical. Scales smooth, in twenty-eight series on the posterior part of the neck, and in thirty-three on the middle of the body. Ventral shields rather narrow, 253; anal 1; subcaudals entire 70.

The ground colour is a reddish olive, more yellowish inferiorly; a reddish brown streak from the nasal shield through the orbit to the angle of the mouth. A series of about eighty transverse reniform spots from the head to the end of the tail; each spot light reddish brown, edged with dark brown. Another series of similar, but smaller and irregular spots along each side; belly nearly uniform.

Length of the head $\frac{2}{3}$ inch, of trunk 22 inches, of tail 4 inches.

This species comes from Western Africa, but I am not aware from what particular part. I have named it after Mr. Ford, whose merits in herpetology are well known by his truly artistical drawings.

4. ON A NEW SPECIES OF FISH OF THE GENUS GERRES.
BY DR. A. GÜNTHER, FOR. MEMB.

(Plate XXIV.)

GERRES LONGIROSTRIS, Rapp. (Pl. XXIV.)

D. $\frac{9}{11}$. A. $\frac{3}{8}$.

The height of the body is one-half of the total length (without caudal). Præorbital and præoperculum not serrated. *Dorsal fin not notched*; the spines are moderately strong; the length of the second being four-sevenths of that of the head, or two-fifths of the depth of the body. The second and third anal spines are nearly of the same strength, and considerably shorter than those of the dorsal fin. Silvery, with darker stripes along the series of scales.

Hab. Cape of Good Hope.

This species is distinguished from all the others by the dorsal fin, which has the upper margin even, without any indication of a notch. It has been established by Professor W. von Rapp from specimens in

PELOPHILUS FORDII.

GERRES LONGIROSTRIS.

the Museum of Tübingen, and to him I am indebted for the accompanying fine drawing (Pl. XXIV.).

5. DESCRIPTIONS OF A NEW GENUS AND SOME NEW SPECIES OF SHELLS FROM THE COLLECTION OF HUGH CUMING, ESQ.
BY H. ADAMS, F.L.S.

Genus OCANA, H. Adams.

Operculum costa spirali convexa, granosa instructum; valde perforatum; margine externo simplici.

Testa turbinata, solida, lævis, imperforata; spira brevis, conica; apertura subcircularis, latior quam longior; labium planatum, excavatum, antice subproductum, callo tenui extenso.

Operculum with a convex, granular spiral rib; axis deeply perforated; outer edge simple.

Shell turbinate, solid, smooth; axis imperforate; spire short, conical; aperture subcircular, wider than long; inner lip flattened, excavated, scarcely produced anteriorly, with an extended thin callus.

Ex. *Ocana helicina*, H. Adams.

Turbo helycinus, Born.

Turbo cidaris, Gmel.

When this species was included by my brother and myself in *Sarmaticus*, to the shells of which genus it bears considerable resemblance, the operculum was unknown to us. I have, however, recently had an opportunity of examining it, and find it to be entirely different from that of *Sarmaticus*. It approaches more nearly that of *Callopora*, but it is without submarginal ribs. The species is from the Cape Colony, which is also the locality of the two known species of the former genus.

Genus PHYSA, Drap.

Subgenus AMERIA, H. Adams.

Testa anfractibus postice planatis, et angulatis vel carinatis; spira brevi, depressa.

Shell with the whorls flattened, and angulated or carinated at the posterior part; spire short, depressed.

1. PHYSA (AMERIA) CARINATA, H. Adams.

P. testa subovata, tenui, pallida flavo-fusca; spira brevissima, apice planato; anfractibus 3, postice planatis et valde carinatis; apertura subovali, plica columellari mediocri.

Shell subovate, thin, pale yellowish-brown; spire very short, the apex flat; whorls three, flattened and strongly carinated posteriorly; aperture subovate, columellar plait moderate.

Long. $5\frac{1}{2}$, diam. $3\frac{1}{4}$ lin.

Hab. Boyne River, Australia.

2. *PHYSA (AMERIA) TRUNCATA*, H. Adams.

P. testa truncato-ovali, pellucida, fragili, pallido-fusca; spira truncata; anfractibus 3, valde carinatis; apertura subovali, peritremate continuo; plica columellari obsoleta.

Shell truncate-ovate, translucent, fragile, pale brown; spire truncate; whorls three, strongly keeled; aperture subovate, peritreme continuous; columellar plait obsolete.

Long. 4, diam. $2\frac{1}{2}$ lin.

Hab. Calliope River, Australia.

3. *PHYSA (AMERIA) OBESA*, H. Adams.

P. testa truncato-ovata, brevi, tenui, fusca; spira brevissima; anfractibus 3, postice carinatis et planatis; apertura subovali, lata, peritremate continuo; plica columellari obsoleta.

Shell truncate-ovate, short, thin, brown; spire very short; whorls three, keeled and flattened posteriorly; aperture suboval, wide, peritreme continuous; columella fold indistinct.

Long. $3\frac{3}{4}$, diam. $2\frac{1}{2}$ lin.

Hab. Fitzroy River, Australia.

4. *PHYSA (AMERIA) CUMINGII*, H. Adams.

P. testa ovali, tenui, pallido-fusca; spira brevi, anfractibus 4, postice angulatis et planatis; apertura subovata, peritremate continuo; plica columellari mediocri.

Shell ovate, thin, light brown; spire short; whorls four, angulated and flattened at the posterior part; aperture subovate, peritreme continuous; columellar plait moderate.

Long. 8, diam. $5\frac{1}{4}$ lin.

Hab. Port Essington, Australia; New Zealand.

5. *PHYSA (AMERIA) MÆSTA*, H. Adams.

P. testa oblique ovali, tenui, olivaceo-fusca; spira mediocri; anfractibus 5, postice carinatis et planatis; apertura ovali, plica columellari conspicua.

Shell obliquely ovate, thin, olive-brown; spire moderate; whorls five, carinated and flattened posteriorly; aperture oval, columellar plait distinct.

Long. 7, diam. $5\frac{1}{2}$ lin.

Hab. New Zealand.

Genus *PHYSOPSIS*, Krauß.*PHYSOPSIS JUKESII*, H. Adams.

P. testa subglobosa, tenui, albida; spira brevi, obtusa; anfractibus 4, convexis; apertura subovali, peritremate continuo; columella recte truncata.

Shell subglobose, thin, white; spire short, obtuse; whorls four,

convex; aperture subovate, peritreme continuous; columella sharply truncated.

Long. 3, diam. 2 lin.

Hab. Port Essington, Australia.

This species was discovered by Mr. Jukes at Port Essington. The only other species of the genus known is that described by Krauss from Port Natal.

Genus PLANORBIS, Guett.

Subgenus HELISOMA, Swains.

PLANORBIS (HELISOMA) LAUTUS, H. Adams.

P. testa subovata, tenui, pari altitudine et latitudine, pallide alba, supra profunde et anguste umbilicata, infra planata; anfractibus 3, celeriter crescentibus, convexiusculis, superne angulatis et angustis, subtus carinatis, subtilissime striis decussatis; apertura subobliqua, subovata, supra anfractum penultimum surgente, peritremate continuo.

Shell subovate, thin, the height equalling the width, yellowish-white, deeply and narrowly umbilicated above, flat below; whorls three, rapidly increasing, rounded, angulated and contracted above, carinated below, decussated by fine striæ; aperture slightly oblique, subovate, extending above the penultimate whorl, peritreme continuous.

Diam. 2 lin.

Hab. New Orleans.

Subgenus ADULA, H. Adams.

Testa anfractibus convexis et numerosis, supra profunde umbilicata, infra convexa; apertura campanulata.

Shell with the whorls rounded and numerous, deeply umbilicated on the upper, and convex on the under side; aperture campanulate.

Ex. P. (Adula) multivolvis, Case.

This species of *Planorbis*, which is one of the few having campanulate apertures, is so different from the rest in the whorls being numerous and slowly increasing, and in the spire being depressed considerably below the under side, that it appears desirable to separate it from them.

6. REVIEW OF THE VERMETIDÆ. BY OTTO A. L. MÖRCH (OF COPENHAGEN). (PART I.)

(Plate XXV.)

The shells of the *Vermetidæ* are at once to be distinguished from those of the *Serpulæ* by the presence of a spiral nuclear shell and of concave smooth interior septa; but these parts are often lost or con-

PROC. ZOOLOG. SOC.—1861, No. X.

cealed, and it is then exceedingly difficult to decide with certainty whether the shell is the production of an Annelid or of a Molluscos animal.

If the shell is formed of a solid matter strongly sculptured with longitudinal grooves or scales, or of a brownish colour, it is certainly formed by a *Vermetus*; but if the shell is of a soft earthy matter, feebly longitudinally grooved, it is doubtful to which subkingdom it belongs.

The shells of the *Serpulidæ* have an anal opening (except the genus *Cymospira*), and appear only to be composed of two layers, the *Vermetidæ* having three.

There is a striking analogy between the *Vermetidæ* and *Ostreidæ* in respect of colours, sculpture, and growth. As in the latter family, the sculpture of the *Vermetidæ* is much influenced by the bodies to which they attach themselves, the surface often showing the cast of the sculpture of a coral* or another shell.

I have convinced myself by the comparison of numerous specimens, chiefly of *Siphonium nebulosum* and *Vermetus decussatus*, Gm., that one and the same species may vary, from being strongly grooved and scaly to being quite smooth and polished.

The form and growth are not less variable; the same species being found nearly regularly spiral, top-shaped, planorbiform, or as a nearly straight tube. The same species is sometimes affixed and solitary, sometimes free and agglomerated, imbedded in mud (*Spirogyphus glomeratus*, Biv.). Some species (*Siphonium nebulosum*, Dill.) are affixed during the greater part of their lives, and seem then, like *Cacum*, to "decollate" the shell and live afterwards free in the mud; perhaps this may have relation to the sexual functions†.

The thickness of the shell may vary from being thin and nearly pellucid to being very thick and incrustated; if the shell be laterally affixed, the dilated angular sides are filled out with a solid calcareous mass (Chenu, Illustr. pl. 2. f. 2).

In the interior of several species (*Petalococonchus*) are found some very long lamellæ, generally regarded of generic value. I have reason to think that these lamellæ are dissolved with age, like the teeth of some species of *Pupa*‡; or perhaps they are of a sexual nature, like the interior cup of *Begonia* (*Thecalia concamerata*, Chem., from the Cape of Good Hope, which is said to contain the fry.

Daudin and Marcell de Serres thought they had discovered a generic character in the habit possessed by some Wormtubes of burrowing in other shells. Although this character is more developed and constant in some species than in others, it is scarcely of specific value, and is common to nearly all young *Vermeti*. Vertically growing species often show the same dissolving power in removing parts

* Seba, 'Thesaurus,' vol. iii. t. 93. f. 9, and *Ostrea cumingii*, Dkr. in Philippi Abbildungen, t. 1.

† Siebold (Lehrbuch der vergleich. Anatomie, p. 355) denies the presence of an exterior penis; but Rüppell (Atlas, t. 11. f. 3) has figured a species with an exert male organ. It appears that the large fusiform cells containing undeveloped spermatozooids (Siebold, l. c.) are Spermatophores.

‡ V. Martens, Ueber *Pupa Wienlandi* Kur. Mal. Blätter. 1859, p. 209.

of neighbouring tubes which stop their development. *Saxicava* and other genera of burrowing Bivalves afford sufficient examples of the generic and specific value of this character.

The colours are exceedingly variable in one and the same species, and often offer a good guide to determine the genus in the absence of the operculum. The young shells are generally of a brownish colour; the adults vary sometimes from brown or black to pure white.

It sometimes appears that the *Vermeti*, *Balani*, *Ostrea* and *Anomia* not only borrow the sculpture, but even the colours from other shells. It is frequently the case that animals and plants (chiefly sea-weeds) of the most different families and classes living under the same circumstances, are coloured in the same manner. Brewster has shown that the mother-of-pearl colour is received by a cast of melted wax, and consequently is not chemical but purely mechanical, like the different colours of the clouds. A further inquiry will perhaps prove that this fact does not stand quite alone.

The presence or absence of an operculum is of great generic value; but the tube alone does not seem to afford any character by which the operculated species can be distinguished from the non-operculated. The different manner of interception will perhaps prove of use in this question, but further observations must first be made. The shape of the operculum varies in thickness and convexity according to age, but this seems not to be of great specific value, although it is indispensable for the generic determination.

The long bristles of the lid * of the genus *Stephopoma* afford excellent specific characters. The surface of the lid of *Siphonium* and some other genera shows nothing particular, even under the lens; but if softened in water and scratched with a knife, some small spiny bodies are obtained, which under the microscope look very like some cylindrical nodulous forms of the Cactus family (see Plate XXV. fig. 16). All the *Vermeti* seem to be viviparous. The unborn shells, easily found in the dried animals, have the same fixed form and sculpture as shells in general, and are of more consequence for the establishment of species in this family than in any other.

BURTINELLA, Mörch.

Moerchia, Mayer, Journ. de Conch. viii. August, 1860, p. 309, non A. Adams, Ann. and Mag. N. H. April 1860.

Vermicularia, Mantell.

Solarium, Galeotti.

Vermetus, Nyst.

Serpula, Phil.

This genus, which has no interior septa, is perhaps most allied to *Siliquarius*, which it resembles in the peculiar exterior sculpture, but it differs from it in wanting the branchial slit, like the upper whorls of

* The only shell before known to have a ciliated operculum is *Turritella unguina*, first mentioned by Lovén and Forbes and Hanley; but it has been overlooked that the cilia have five small equidistant spines on one side.

that genus. All the species are fossil, unless the shell figured by Humphrey (Conchology, t. x. f. 8 *) is a living species of the genus.

1. BURTINELLA CONTRARIA (*Trochus*), Schröter.

1777. Kräusel, Martini, Neue Manigfalt. iv. p. 404, t. 1. f. 2.

1783. Schröter, Einleitung. i. p. 784.

1786. Fragmentum *Trochi contrarii* petrificati, Chem. ix. p. 134. f. 977.

1788. *Trochus contrarius*, Schröter, Register, p. 107.

1791. *Trochus ferrugineus*, Gm. Syst. Nat. no. 70.

Fossil from Rhänden or Rhendenberge near Schafhausen.

The last whorl is strongly angulated. The original specimen is still preserved in the Spenglerian Collection.

2. BURTINELLA NYSTII (*Solarium*), Galeotti.

1784. Burtin, Oryctogr. de Bruxelles, pl. 8. f. 9 (Mayer).

1837. *Solarium nystii*, Galeotti, Mem. Geol. de Brab.

1843. *Vermetus nystii*, Nyst, Polyp. Foss. de Belg. ii. p. 373. pl. 36. f. 8.

1860. *Moerchia nystii*, Mayer, Journ. de Conch. viii. p. 310.

Inferior Tertiary formation (Parisienne et Bartonienne).

3. BURTINELLA TURBINATA (*Serpula*), Philippi.

1846. *Serpula turbinata*, Phil. Verz. Magdeburg. Verstein. in Palæontographica, 1846, August 1, p. 80. t. x. f. 14.

Inferior Tertiary sand (lingurian) near Magdeburg (*Phil., Mayer*).

4. BURTINELLA SOWERBII, Mantell (sp.).

1822. *Vermicularia sowerbii*, Mant. Sussex, p. 111. t. 18. f. 14, 15. Hornsey marl pits (grey chalk marl), *Mantell*.

5. BURTINELLA PHILLIPSII, Mörch.

1835. *Vermicularia sowerbii* (Mantell): Phillips's Yorksh. i. p. 95. pl. 2. f. 29.

Speeton Clay (*Phillips*).

This species, according to the author, is also found dextral.

6. BURTINELLA NODUS, Phillips, 1835.

Vermicularia nodus, Phill. Yorksh. i. p. 125. pl. 124. f. 34.

Oolite of Cloughton, Bransby (*Phillips*).

STEPHOPOMA, Mörch, Journ. de Conch. 1860, Jan., p. 42.

Vermetus, Quoy & Gaim.

Siliquarius, part, Gray.

Tenagoda, part, Adams.

T. adulta affixa, contorta, solitaria vel agglomerata; apertura

* That the figured shell is in reality *sinistral*, like the species of this genus in general, is proved by the other *dextral* shells being *sinistral* in the accidentally reversed plate.

superne leviter inflexa, inferne obsoletissime effusa; stria incrementi bis arcuatim retroflexæ.

Operculum arctispirale, inferne convexum, superne concavum, setis longis multifidis munitum.

Animal viviparum, descriptum sequente modo in opere dicto 'Voyage de l'Astrolabe,' "En avant du mufle est un appendice median vertical, évasé en entonnoir, susceptible de s'allonger; un peu plus en avant en est un autre qui se recourbe vers l'opercule, et qui n'est peut-être que le pied lui même renflé. Ce sont les analogues de ceux que nous avons vus pairs dans les espèces précédentes, et dont nous ignorons l'usage. Le pied a de plus, sur le côté droit seulement, un petit tubercule pointu, tel qu' Adanson l'indique dans son espèce. La cavité respiratrice est ample, et une branchie, à lamelles jaunes, cylindriques et rigides, se rebrousse par-dessus le manteau et fait saillie à l'extérieur."*

Testa natalis nautiliformis sed asymmetrica, arcuatim regulariter granulosa; operculum nudum membranaceum planum.

Cuvier, in 1830†, was the first to mention *Vermeti* with a spiny operculum, a fact probably communicated to him by the naturalists of 'L'Astrolabe,' just returned the preceding year from their second circumnavigation.

The form of the aperture varies in the same species from being exactly circular to ovate or reniform. A peculiar obliquity in the aperture of some specimens in each group led me to recognize the species of this genus before the lids were found; this obliquity is very likely occasioned by the long opercular bristles often forcing the animal to withdraw the operculum obliquely, and, by their frequently touching the border of the aperture, stopping the growth of the tube at that side. The bristles, giving the lid quite the appearance of a broom, are much worn, and have the interstices always filled up with fine clay. The use of these bristles in the economy of the animal it is difficult to guess, but it may very likely be the same as that of the spiny horns on the lid of *Cymospira gigantea*, Pall., and the pallets of *Xylotrya* (Leach), which in shape are not very unlike the bristles of *Stephopoma roseum*, Quoy & Gaim. (see Plate XXV. fig. 11). The bristles seem under the microscope longitudinally canaliculated *outside*; but I believe this canal in reality is the hollow interior, as I have distinctly seen under the compressor the air-bubbles move in the interior and even *out* into the lateral spines. The sculpture of the unborn shells has a striking resemblance to that of *Argonauta oryzata*, Meusch., and is continued in a part of the first whorl of the affixed shell. In *Stephopoma tricuspe*, Mörch, the granules look under the microscope like holes with a dark border. The small holes described by Quoy and Gaimard as covering the shell of *Stephopoma roseum*, and supposed to be formed by some parasitical animal, will perhaps prove to be an optical deception occasioned by the sculpture.

* Adanson has mentioned a little opening in the border of the mantle, but not in the foot.

† Cuvier, Règ. Animal. 1830, iii. p. 109.

1. STEPHOPOMA ROSEUM, Quoy & Gaim. (Pl. XXV. f. 11).

"*T. minima, spirali porrecta, cylindrica, rugosa et rosea.*

"Long. 6-8 lin.

"Opercule rond, brun, multispiré, recouvert d'une grande quantité de soies cornées, divergentes et multifides."

1833. *Vermetus roseus*, L'Astrolabe, ii. p. 300. t. 17. f. 20-24.

Siliquaria rosea, Gray, Figures, iv. p. 83. t. 56. f. 3; Guide, p. 128.

Stephopoma roseum, Mörch, J. de Conch. viii. p. 42.

"La rivière Tamise (qui est saline) à la Nouvelle Zelande, par d'assez grandes profondeurs."—Quoy et Gaimard.

2. STEPHOPOMA TRICUSPE, Mörch (Pl. XXV. f. 1).

T. agglomerata tenuiuscula, ferruginea hic illic albescens, laxè spirata, longitudinaliter obsoletissime subtiliter-lirulata; striæ et rugæ incrementi irregulares leviter arcuatæ et reflexæ; anfr. primi 2-3 conjuncti helici-vel planorbiformes, sub lente minutissime arcuatim granulosi; apertura circularis vel reniformi-ovalis, faucibus sæpe badiis.

Diam. aperturæ 4 m. (circ.).

Operculum arcte spiratum, superne setis tricuspidibus breviusculis confertissimis instructum; inferne planum, badium, concentrice sulcatum et striatum; disculo centrali subimmerso, fossula declivi circumdato; verruca centrali punctiformi; limbus flavescens, latus, nitidus, margine tenui, reflexo.

Setæ triramosæ, stipite brevi crassiusculo supra basin contracto, cingulis 5-6 spiniferis; ramus medianus erectus, gracilis, longissimus, nudus, stipite quadruplo longior; ramus sinister dimidium longitudinis vix attingens, extrorsum quadrispinosus, spinæ infimæ geminæ; ramus dexter oppositus brevissimus, vertice trispinoso (fig. 2).

Testa natalis nautiliformis (sed asymmetrica) nitida, pallide iridescens, alba vel pallide cyanea, arcuatim punctato-granulosa; granula sub microscopio inspectæ, ocellatæ (centro pellucido annulo obscuro circumdato). Spira subulata, obliqua, vix prominens, ferruginea; sutura impressa, canaliculata; umbilico plano, apertura obliqua subhexagonalis; peristoma superne leviter inflexum; anfr. ultimus carina lata convexiuscula. Striæ incrementi profundæ, unde peripheria crenulata. Inveni circiter x. pullos in specimine desiccato.

Diam. maj. 1 m.

Australia; massa agglomerata (coll. Cuming).

I cannot see any trace of the specimens having been affixed to foreign objects. It is not unlike *Pyxipoma lacteum*; but all the specimens are agglutinated to each other.

3. STEPHOPOMA SENTICOSUM, Mörch (Pl. XXV. fig. 2 & 14).

T. alba, crassa, aperturam versus tenuis, contorta vel interdum

laxe spirata, umbilico angusto; anfr. primi 1-2 modo testæ natalis oblique arcuatim granosi, anfr. sequentes longitudinaliter lirulati, liræ interdum asperæ, sed plerumque omnino evanescentes; interstitia lirula intercalante, anfr. ultimus obsolete biangulatus, teretiusculus; apertura circularis interdum subovalis, superne obsolete inflexa; faucibus interdum pallide castaneis; striæ et rugæ incrementi irregulares, superne et inferne arcuatim retroflexæ.

Diam. aperturæ circ. 4 mm.

Operculum superne concavum setigerum, centro nudo, inferne conico-convexum, badium, limbo nitido castaneo; centro liris concentricis 3-4. Setæ longiusculæ, porrectæ utrinque spinis circ. 12 plerumque suboppositis, spinæ majores inferne spinis 1-2 horizontalibus; scapo nudo basi calcarato; lamina expansa subbasalis, margine setoso, setis quatuor longissimis gracilibus approximatis, basi ramulo conjunctis; seta interna recta longior, tertia arcuata.

Testa natalis (fig. 14). Argonautæ oryzatæ non absimilis, sed asymmetrice, arcuatim dense regulariter tuberculata; peripheria mammillis validis circiter 25 ornata; spira immersa bullata, umbilico plano; apertura obliqua orbicularis superne inflexa, inferne obsolete effusa. Color castaneus vel albescent.

Diam. maj. $1\frac{1}{8}$ m.

Operculum membranaceum, planum, circulare, inerme, liris duabus concentricis. Mus. reg.

Testæ solitariae vel geminae in Tridacna scapha (Meusch.) affixæ.

4. STEPHOPOMA PENNATUM, Mörch (Pl. XXV. fig. 3-8).

Stephopoma pennatum, Mörch, Journ. Conch. 1860, p. 42.

Stephopoma bispinosum, Mörch, Malacol. Blätter, 1860, p. 78. n. 72.

T. solitaria spiraliter angulatim contorta, umbilico aperto; anfr. primi, modo testæ natalis, oblique confertissime verruculosi, anfr. sequentes longitudinaliter conferte lirulati, quadrangulares vel obsolete pentagoni; striæ incrementi sigmoideæ, antiquatæ, irregulares, squamoideæ; apertura circularis vel subquadrangularis, superne interdum inflexa; anfr. primi crassi, aperturam versus tenues.

Diam. ap. 4 m.

Color albus, badius vel mixtus.

Operculum inferne plano-convexum, badium, liris duabus concentricis; disculo centrali ruguloso subprominulo, limbo angusto pallido nitido; superne concavum radiatum setosum, centro nudo læviusculo; setæ confertæ erectæ, utrinque spinis inæqualibus alternantibus subinflexis hic illic ternatis; scapus nudus inferne calcaratus, versus basin lamina expansa cornu Cervi elaphi non absimilis, margine aculeato, spinis quatuor longis erectis inæqualibus.

Testa natalis (et juvenilis) lateraliter affixa planorbiformis, anfr. primi pulcherrime arcuatim granulosi; granulæ in quadrata dispositæ.

Var. BISPINOSA (Pl. XXV. figs. 9 & 10.), an sp. distincta?

Operculum inferne conico-convexum; setæ spinis utrinque circ. 14 sæpe ternatis, calcar basale validissimum erectum (lamina basalis deest?).

Although this operculum is smaller than that of the preceding, the basal spur of the bristles is much larger; if this is not in consequence of its being unworn, it may be a different species, especially as it is nearly conic below.

Hab. Realejo, on *Callopoma saxosum*, Wood, and *Crucibulum scutellatum*, Gray. On the latter shell the specimens affixed to the light-coloured parts are white, and those affixed to the brown parts brown; having, however, seen this on one specimen only, I cannot decide whether it is accidental.

SIPHONIUM, Mrs. Gray, Figures of Moll. Anim. iv. p. 82.

Siphonium, "Browne," Mörch, Journ. de Conch. vii. no. 4. p. 353.

Stoa, M. de Serres, Annales des Sciences Nat. iv. 1855, p. 238.

Operculum magnum, concavum vel infundibuliforme, obsolete spirale, inferne plerumque mammilla centrali.

Patrick Browne comprehended under the generic name *Siphonium* nearly all those different forms of "worm-tubes," which Linnæus afterwards included in his genus *Serpula*. The first species is thus characterized in the 'Civil and Natural History of Jamaica,' "*Sub-æquale angulatum et flexuosum*." The second word very likely signifies triangular (*triqueter*), and by "*flexuosus*," is probably meant creeping horizontally in bends, in opposition to "*contorta*," which applies better to the *Vermeti*. It seems to me therefore likely that the first and typical species is a *Serpula*. The fifth species is without doubt *Vermetus Knorrii*, Desh., a form for which Lister previously had instituted the name *Vermiculus*. The only figured species is a *Teredo*. I have adopted the name after Dr. Gray, as it does not appear to have been used before. *Siphonium*, Link., is very badly described in the 'Catalogue of the Rostok Collection,' iii. p. 9, 1807, thus: "Shell straight, with septa forming rings outside. Here belong the straight *Nautili*. *Siph. fasciatum* is exactly like *Nautilus fascia*, Gm., but larger." The species of Gmelin belongs to the Foraminifera.

M. de Serres has established a genus of "Annelides sedentaires tubicolés," under the name *Stoa*, chiefly on account of their habit of burrowing a bed in the surface of other shells, exactly corresponding to the genus *Spirogyphus* of Daudin. Both authors have mixed together species of two very different genera, the one with a concave, the other with a thick convex operculum; but as Daudin has selected for illustration of his genus a species with an operculum of the latter kind, I regard the represented species as the type, although

it is the second in order. The genus *Stoa* is characterized thus: "Tube testacé contourné en spirale orbiculaire et irrégulière d'un forme discoïde, renflé et convexe; dernier tour détaché du premier, et se prolongeant parfois en un tube droit, ouverture ovulaire, terminé par un *opercule solide*, calcaire, conique et surchargé, formé par de très petites bandes circulaires presque subspirales." A conical operculum, which is solid and calcareous, seems so unnatural and without analogy among the Mollusca and Annelida that I suppose a covering of *Nullipora* or some other calcareous matter, which I frequently have observed, has given rise to the above statement. In the figure of M. de Serres the last whorl of the operculum terminates abruptly at the edge, just like the lid of *Siliquarius*, represented at p. 321 in Chenu's 'Manuel,' which I have never seen so strongly expressed, neither have I seen an operculum of the high conical shape figured. The concavity of the operculum is very different, according to the species. In the centre of the under side there is often a conical wart, sometimes very prominent; but I do not believe it is of generic value, as in a series of species it decreases more and more, and finishes by disappearing entirely. In all the smaller species the operculum is yellow with a red central spot, which, as well as the central wart below, is wanting in the larger species. The whorls are generally few and indistinct; *S. megamastus*, Mörch, has two or three concentric erect laminae, which give the operculum the appearance of being threefold. The surface is covered with some exceedingly small bodies—scarcely to be seen with the naked eye—of a cylindrical nodulous form, invested with numerous spicula of a glass-like appearance (Pl. XXV. fig. 13). These bodies would be of great importance for the distinction of species if their form were not altered by exsiccation, and if they were not, on account of their minuteness, exceedingly difficult to be shown in the same position for comparison. They are at once to be distinguished from the bristles of *Stephopoma* by their microscopic size and, as it seems, the silicious nature of the spicula, which are always broken abruptly like a steel needle; whilst the spines of *Stephopoma* are rarely broken. The axis seems to be of the same substance in both genera. The animal is described by Quoy and Gaimard, and Dr. Gray in the 'Spicilegia.' The propodium is semilunar with projecting corners, with two rather small approximating pedal tentacula, which appear to be rudimentary or wanting in *S. tonganum* and *S. reticulatum*, Q. & G., whilst they are rather large in *S. nebulosum*, Dillw., and considerably exceeding the tentacula in size in *S. platypus*, Mörch; a dried specimen of the latter species showed a flat leaf-like propodium, bringing to mind the under lip of the pupa of the Dragon-fly.

Sect. a. DENDROPOMA, Mörch. *Operculum planiusculum, inferne mammilla centrali validissima, superne laminae 2-3 concentricis suberectis; asperitates vertice trifidæ, spiculis subulatis leviter arcuatis.*

5. SIPHONIUM (DENDROPOMA) MEGAMASTUM, Mörch (Pl. XXV. f. 12 & 13).

T. teres repens varie torta, lactea in fundo ferruginea, liris longitudinalibus subremotis, lira mediana sæpe valida, striæ et rugæ incrementi arcuatæ, intersectionibus squamiferis; interstitia lilarum obsolete foveolata; apertura circularis obliqua; diam. 4-5 m. T. juventute læviuscula liris obsoletis, nitida, tenuis, castanea, plerumque planorbiformis.

Operculum tenuiusculum flavum, superne sat concavum, laminis 2-3 concentricis suberectis, punctis asperis munitum, area centrali aurantia, centro impresso plerumque flavo; inferne planiusculum, limbo nitido pallido convexo; impressio muscularis opaca, cerea, concentrice ruguloso-striata, lirula circulari terminata; mammilla centrali validissima obtuse conica.

Diam. $3\frac{1}{2}$ m.

Asperitates oculo armatissimo inspectæ arborescentes; stirps robustus brevis, ramis tribus porrectis inæqualibus, lateraliter biaculeatis, vertice aculeis 4-6; aculei longi subulati leviter arcuati sed plerumque fracti (fig. 13).

California?: numerous specimens burrowing, but not very deeply, in the surface of a *Haliotis nodosus*, Reeve. The small vaulted compressed scales of the grooves seem to be caused by numerous small holes of perforating animals, chiefly those of *Cliona*, which have made the surface of the *Haliotis* very rough; but it must be observed that the *Vermeti* themselves are also attacked.

Var. CENTIQUADRA.

Aletes centiquadrus, var. *imbricatus*, Carp. Cat. p. 302.

T. brunnea fundo albescens, planorbiformis, semi-immersa, anfr. planati utrinque obtuse angulati, longitudinaliter lirati; liræ subæquales approximatae, striæ et rugæ incrementi arcuatæ, intersectionibus nodulis obsoletis subremotis, vel in locis obtectis, squamulis asperis remotis munitis.

Diam. testæ 20 m.; diam. aperturæ 3-4 m. Operculum ignotum.

California: a single specimen deeply burrowing in an *Haliotis splendens*, Mawe. The sculpture and colour of this shell, which I only regard as a variety, bring to mind the *Aletes centiquadrus*, Val.

6. SIPHONIUM (DENDROPOMA) LITUELLA, Mörch.

Stoa ammonitiformis, M. de Serres, Ann. des Sc. l. c. p. 242, note. *Spirogyphus*, sp., Carpenter, Report, p. 324 (verisimiliter).

T. varie torta plerumque lituiformis, profundissime immersa; anfr. graciles plani, utrinque obtuse angulati, sæpe fossula longitudinali mediana, unde carina vel lira obtusa lateris externi; striæ et liræ incrementi arcuatæ, approximatae, regulares; apertura circularis interdum obliqua, nonnunquam soluta; color albescens vel cinereus, interdum fascia obsoleta fusca, anfr. primi badii. T. infantilis nitida, apice bullata.

Diam. ap. 2 m. (circiter).

Operculum nudum (?) *membranaceum, pallide flavum, superne leviter concavum, area centrali vivide coccinea, lirula circulari mediana circumdata; inferne convexiusculum; area muscularis flava opaca, mammilla centrali conica prominente, limbus pallide flavus nitidus.*

Diam. $1\frac{1}{10}$ –2 m.

California; about ten specimens deeply imbedded in the surface of a young *Haliotis splendens* (coll. Cuming).

This species is to be distinguished from the preceding by its narrower whorls, smaller size, prominent and regular striæ of growth, and the absence of longitudinal liræ. It is difficult without the operculum to distinguish this species from *Spiroglyphus spiruliformis*, Serres. The *Stoa ammonitiformis*, stated by M. de Serres to be found on *Haliotis californiensis*, is very likely the young of this species; but the figured typical form from the East Indies differs in the whorls being broader and cylindrical. If the *Spiroglyphus* from Upper California, mentioned by Carpenter (Report, p. 324), is from the Cumingian collection, it must be this species labelled California. The operculum closely resembles that of the young of the preceding, but wants the microscopical tufts, which are perhaps dependent on age.

7. SIPHONIUM (DENDROPOMA) LEUCOZONIAS, Mörch.

T. planorbiformis, superficialiter corrodens, tequis, castanea, fasciâ pallida vel alba laterali; anfr. teretiusculi, utrinque obtuse angulati, superne (nec lateraliter) liris circiter vi. obsoletissime nodulosi; anfr. ultimus aperturam versus antiquatus; striæ et rugæ incrementi arcuatæ, unde irregulariter reticulatæ; apertura circularis, plerumque subsoluta, rarissime immersa; anfr. ultimus interdum rectus.

Diam. testæ cir. 8 m., aperturæ $2\frac{1}{4}$ m.

Operculum rufescens, superne profunde concavum, opacum, concentrice liratum, periphæria margine angusto obscuro; inferne convexum; area muscularis opaca, concentrice bilirata, centro mammilla valida concentrice subtilissime ruguloso-striata; limbus latus nitidus, lineis duabus concentricis saturate rufis, periphæria margine angusto albo.

Diam. 2 m.

West Africa; on a specimen of *Purpura neritoidea*, L., var. *columnella unimaculata*; about thirty-six specimens with a *Vermetus adansonii*, Daudin? The periphery below the angle is white and without liræ. Most of the specimens do not burrow very deep, except where they meet the nodules or ribs of the *Purpura*.

Var. *datur lituiformis*.

8. SIPHONIUM (DENDROPOMA ?) TEREDULA, Mörch.

T. albescens, in fundo interdum sordide carnea, repens corrodens, varîe torta vel glomerata, anfr. teretes antiquati, liris longitudinalibus alternatim minoribus; striæ vel rugæ incrementi acutæ, subremotæ, sigmoideæ; apertura circularis.

Diam. circ. $2\frac{1}{4}$ m.

On *Haliotis tuberculata*, probably from Morocco. Operculum unknown.

Deeply burrowing, about half of the shell being concealed.

*Vermetus annulatus**, Rousseau, Chenu's Illustr. pl. 2. f. 2 (the smaller figure) looks very like this shell; but it is a gregarious, not burrowing species. *S. teredula* is perhaps most allied to *S. megamastus*, Mörch.

9. SIPHONIUM (DENDROPOMA ?) AFRUM, Gm.

Le Datin, Adanson, Senegal, p. 165. pl. 11. f. 4.

Serpula afra, Gm. S. N. p. 3745, no. 23.

T. solidissima, albida, corrodens, lituiformis, versus aperturam tenuiuscula, anfr. antiquati, utrinque obtuse angulati, superne planiusculi, liris longitudinalibus alternatim minoribus obsoletis; sulci incrementi profundi irregulares, striæ incrementi confertæ arcuatæ, inferne lateraliter reductæ; apertura circularis vel subquadrangularis.

Diam. aperturæ circ. $3\frac{1}{4}$ m. Operculum ignotum.

Hab. Gaboon, Western Africa, on *Chama senegalensis*, R.; two specimens. Closely allied to the preceding, but differs in being very thick; the whorls are nearly regularly quadrangular; the striæ of growth are very prominent, except in the last whorl. According to Adanson, the lid is only one-sixth of a line in diameter; it is therefore doubtful if it belongs to this genus.

10. SIPHONIUM (STOA) POLITUM (Daudin).

Spiroglyphus politus, Daudin, Recueil, 1801, p. 49.

"*S. irregulariter spiralis, politus, apertura rotunda.*"

"Diam. de 3 lignes au plus. Tube blanc, poli, roulé en deux tours de spirale irrégulière, et plus gros à son ouverture, qui est cylindrique. Il se creuse un lit et s'attache sur diverses coquillages bivalves de l'Inde, du genre des jambonneaux et des peignes." The expression, "plus gros à son ouverture," and the above measurement, lead me to consider this species as identical with *Stoa perforans* of De Serres.

a. Stoa perforans, M. de Serres, Ann. d. Sc. N. 1855, v. p. 241. t. 8 c. f. 3-6.

"Cette espèce, à tube court, à spirale raccourcie, dont la dernier tour est sur le même plan que le second, présente dans son ensemble une forme à peu près discoïde. L'ouverture de la bouche ample, arrondie est analogue à celle des Cyclostomes. Les dimensions dépassent peu 0^m·010 à 0·012."

Represented on a young specimen of *Tridacna gigas*, L.; but, p. 239, it is indicated to be found on *Tridacna*, *Hippopus*, and *Serpula*; p. 238, the species of the latter genus is named *Serpula panamensis*, Chenu, which is *V. eburneus*, Reeve.

* It is without doubt *Serpula annulata*, Lam., 1818, v. p. 364, no. 11.

11. SIPHONIUM (STOA) PLATYPUS, Mörch.

T. solidissima, corrodens, spira planorbiformis anfractibus angustis, anfr. ultimus ovalis, aperturam versus planus, utrinque obtuse angulatus, striæ et stratae incrementi arcuatæ remotæ irregulares in latere externo acutæ; apertura circularis obliqua, peritremate superne crasso; color albus, faucibus fusciscentibus.

Diam. aperturæ 7 m.

Operculum tenue pellucidum pallide flavum, superne concavum laminis concentricis circiter 4 obsoletis appressis; inferne convexum, limbo nitido latiusculo; impressio muscularis opaca, centro convexo lato aurantiaco, tuberculo minimo elongato sub-centrali (an morbo?).

Diam. $6\frac{1}{4}$ m.

Animal rostro brevissimo; capite depressiusculo, utrinque acute carinato, inferne fornicato excavato et transversim pulcherrime lirate; tentacula breviter obtusiuscula oculos ad basin externam gerentia; os minutum infundibuliforme; propodium foliaceum antice bilobatum, utrinque cornu tentacula duplo vel triplo superante. Mesopodium? postice uti bituberculatum. Capite superne et tentaculis utrinque maculis nigris irregularibus ornatis. Ex specimine desiccato.

Hab. Sandwich Islands, on a *Chama*. The specimen has burrowed in its own spire. This shell resembles very much the larger figure of *Stoa perforans*, M. de Serres, but is considerably larger, and has the whorls flattened. The operculum is more flat than conical.

12. SIPHONIUM (STOA) DA-COSTÆ, Mörch.

Serpula, Humphrey, Conchology, t. 9. f. 15.

Spiroglyphus da-costæ, Mörch, Journ. de Conch. 1860, p. 46. n. 8.

Since the above name was published I find in my note-book that I have seen this shell, perhaps the very specimen of Humphrey, in the cases of the British Museum, under a more correct name. Although I have not at present any specimen for comparison, I suppose this species is sufficiently distinguished by its crest and want of longitudinal grooves. It appears to be one of the largest burrowing Worm-tubes.

13. SIPHONIUM (STOA) SUBCRENATUM (Lam.).

"T. repente flexuosa, albida; carina dorsali carinisque laterilibus dentato-crenatis, operculo brevissime conico.

"Ocean indien, sur le Spondyle mutique. Elle se creuse un lit sur la coquille" (coll. Lam.).

Vermilia subcrenata, Lam. 1818, v. p. 370; ed. Desh. v. p. 634: Blv. Dict. Sc. t. 57. p. 329.

Stoa? subcrenata, M. de Serres, Ann. Sc. 1855, v. p. 240.

The shell represented in Chenu's Illustrations, pl. 9. f. 3, et pl. 10.

f. 16, has not a single character in common with the description of Lamarck. A *Spondyle mutique* is not mentioned in Lamarck's later publications; but I suppose it is the *Spondylus microlepos*, Lam. (no. 16), according to the expression "cet *Spondyle* semble *mutique*," a species not even mentioned in Chenu's Monograph. A query is added to the locality "Ocean indien." I refer to this species the following varieties:—

Var. α. SQUAMULOSA.

T. albescens, planorbiformis vel repens, sæpe profunde corrodens, longitudinaliter lirata; liræ approximatae, huc illuc alternatim minores, nodulis transversim compressis sat remotis, et in locis occultis squamis parvis acutis ornatis; liræ 1–3 laterales interdum validiores; striæ incrementi obliquæ, obsoletæ, squamiferae; apertura subcircularis.

Operculum membranaceum, diaphanum, pallide flavum, superne sat concavum, vix striatum, inferne convexum, limbo latissimo planulato, flavo, nitido, liris concentricis tribus; mammilla centrali prominente obtusa, obscure purpurea, fascia basali aurantia circumdata.

Diam. $1\frac{3}{4}$ m.

On *Tridacna scapha*, Mensch. (Mus. reg.)

Var. β. BIFUNICULARIS.

T. crassa, profunde erodens, transversim oblique ruguloso-squamosa, longitudinaliter subtilissime lirulata, liris duabus latis subtilissime longitudinaliter liris.

Diam. 4 m.; diam. operculi 3 m.; liræ circiter $\frac{1}{2}$ m. latæ.

On *Turbo margaritaceus*, var.? The two broad longitudinal grooves look like a cast of the furrows of the *Turbo*.

Var. γ. CRISTATA.

T. triquetra, crassiuscula, carina mediana squamis validis remotis; utrinque angulis acutis, squamis latis angulatis ornatis; liræ incrementi pulcherrimæ, regulares, arcuatæ, leviter flexæ in interstitiis; apertura superne rectangularis.

Diam. $2\frac{1}{2}$ m.

On the same shell as the preceding, burrowing in the suture. Not unlike *Vermetus carinatus*, Rouss. (Chenu, Ill. t. 2. f. 4).

Var. δ. SUBDECUSSATA.

T. repens, profunde immersa, obtuse quadrangularis, antiquata, lirulæ longitudinales subtilissimæ, interruptæ, striæ et lirulæ incrementi subtilissime conferte undulatæ, unde obsolete granulosa.

Diam. 2 m. Operculum differt a præcedentis macula coccinea centrali latiore.

This variety, as well as the preceding, is very like *S. lituella*, no. 6. On the operculum was found a bristle with minute distant spines; but I am not sure that it is not quite a foreign body.

Var. *c.* SPINOSA (Pl. XXV. fig. 16).

T. gracilis, candida, irregulariter spinoso-squamosa, vix lirulata, intus lactea politissima.

Diam. 3 m.

Operculum pallide aurantiacum, superne infundibuliforme, centro impresso, spiraliter conferte lirulato, nodulis microscopicis subcylindricis vertice medio inflexo spiculis obsitis; inferne breve conicum, area muscularis parva, margine concentrico elevato terminata, centro acuto submamillato rufo; limbo latissimo nitido, pallide aurantiaco, margine angusto peripherico pallidissimo terminato.

Diam. 2 m.

Hab. Ins. Philippin. (H. Cuming, Mus. Reg.); imbedded in the lower valve of a *Spondylus nicobaricus*. Some parts of the tube are quite free, but near the aperture it is closely imbedded: it seems consequently not to be burrowing except perhaps in the very first whorls. The spines resemble those of the *Spondylus*; it may prove to be a distinct species.

14. SIPHONIUM (STOA?) TEXTUM, Mörch.

T. corallicola, albida planorbiformis, omnino immersa; anfr. utrinque rectangulati acuti, aperturam versus ampliati, superficie planissima vel leviter concava; laminæ incrementi arcuatæ, margine conferte undulatæ, regulares erectæ obtusæ, interstitiis profundis angustis; lirulæ longitudinales 5-6 spurie obsoletæ, aperturam versus lirulis obliquis tenuissimis; apertura circularis, obliqua.

Diam. testæ 15 m., diam. aperturæ $5\frac{1}{2}$ m.

Operculum pallide flavum, pellucidum, superne profunde concavum, laminis concentricis confertis appressis obsoletis; inferne convexum, area muscularis spiraliter obsolete lirulata, centro coccineo prominulo obtuso; limbus latus nitidus.

Diam. $5\frac{1}{4}$ m.

Hab. Ins. Philippin. in *Porite* emortuo, profundissime corrodens. H. Cuming. legit; extat in collectione Dr. Hornbeckii.

Var. *a.* SCAPHITOIDES.

T. scaphitiformis, spira occulta; apertura circularis resupinata æqua, margine postico extus undulato, punctis duabus aurantiis.

Diam. aperturæ fere 7 m.

Hab. Ins. Philippin., in *Porite* emortuo profundissime corrodens (coll. Cuming.).

Testa juvenilis planorbiformis, umbilico lato planiusculo; anfr. primi fusci, rugæ incrementi arcuatæ geminæ, antice insculptæ, cinereæ; intus fascia columellari latissima castanea, marginibus pallidis.

Diam. 4 m.

(Coll. Dr. Hornbeck.)

The sculpture looks very like the knitting of a stocking, and would

be taken for a cast of the coral, if several other specimens of *Vermeti* on the same kind of *Porites* did not show a different sculpture. This species is the deepest burrowing of all the *Vermeti*, the surface being exactly in the same plane as the surface of the dead coral in which it burrows. The young shell closely resembles *Spiroglyphus spiruliformis*, M. de Serres, but the whorls are broader and flatter. The operculum is covered with microscopical bodies similar to those shown in fig. 16 of the plate.

Var. β . UNGUICULATA.

T. corallicola erodens semi-immersa, spira planorbiformis badia, anfr. utrinque obtuse rectangulati, superne plani, liris obsoletis subremotis, versus aperturam saepe linea castanea (interdum undulata) marginatis; laminæ incrementi tenuissimæ appressæ margine irregulari, in locis occultis, præsertim in antico latere varicum, crassiores, conferte undulatæ; apertura circularis libera, faucibus albis, columella badia, colore in latere sinistro perspicuo; pone aperturam varicibus duabus erectis, validis, acutis, antice excavatis.

Diam. apert. $5\frac{1}{2}$ –6 m.

I have no hesitation in referring this shell to *S. textum*, and the presence of the characteristic undulated laminæ of growth in a few concealed places, and the two small ferruginous spots in var. α corresponding to the brown lines in the present variety, seem to be sufficient reasons for this opinion. In the two varices, the form of the whorls, and the manner of burrowing, it approaches *Siphonium maximum*, Sow. The brown lines sometimes bordering the longitudinal liræ point to some affinity with *S. pictum*.

15. SIPHONIUM (STOA) SCAPHITELLA, Mörch.

T. scaphitiformis affixa vix corrodens; spira planorbiformis late umbilicata; anfr. affixi unde planissimi; superne convexi versus aperturam ampliati, longitudinaliter lirulati; liræ plerumque alternatim minores, lira mediana et liræ lateris interni tres validiores squamiferæ; squamæ compressæ, acutæ, remotæ; latus externum oblique varicoso-rugosum; laminæ incrementi brevissimæ, tenuissimæ, regulares, præsertim in regione umbilicali prominentes, aperturam versus obsoletæ; apertura circularis, obliqua, libera. Color albus; stratum internum lateris affixi denudatum, badium.

Diam. $5\frac{1}{2}$ m.

Operculum flavescens pellucidum, superne concavum centro impresso; inferne convexum; impressio muscularis spiraliter conferte tenuissime lirulata; centro aurantio, disculo obtuso coccineo; limbus latus nitidus.

Diam. 5 m.

The lid is much thinner and somewhat flatter than that of the preceding, the centre is also more blunt and flattish.

Hab. Ins. Philippin.; a single detached specimen in Mr. Cuming's collection.

The affixed side wants the thick exterior layer, so that the thin interior brown one is exposed, which seems to indicate that the animal has been slightly burrowing, although it is quite flat, without any trace of convexity. The size and form are those of *S. textum*, M.; but the sculpture and semicylindrical vaulted shape of the whorls are quite different.

16. SIPHONIUM PICTUM, Mörch.

T. tenuiuscula, repens, superficialiter corrodens; anfr. subcylindrici, superne planiusculi, liris noduliferis compressis 4-5, utrinque linea ferruginea terminatis, lira externa interrupta; noduli obsoleti, elongati, remoti (subsquamosi), antice interdum fissi; interstitia lirulis duabus inæqualibus; laminæ incrementi tenuissimæ, appressæ, margine leviter rimosæ, in interstitiis arcuatis; apertura ampla, circularis; intus ubique badia vel castanea.

Diam. aperturæ 10 m.

T. juvenilis planorbiformis. Operculum ignotum.

Hab. India orientalis; in specimen pedale *Tridacna gigantis* leviter corrodens (Mus. Regium).

Humphrey, Conchology, pl. 10. f. 12, 12 (verisimiliter).

This shell is rather thin in proportion to its size. The laminæ of growth are arcuate between the grooves, and are not unlike those of *S. textum*, var. β , which it also resembles in the *liræ* being most strongly coloured at the sides.

Var. α . TURBOIDES.

T. crassa, candida, turbinata; anfr. ultimus superne planiusculus, antiquatus, fulvus, liris 4-6 ferrugineis, nodulis elongatis, subsquamosis, compressis, badiis; interstitia lirulis 1-2 parvis; latus externum inæquale, noduloso-rugosum, internum planum; laminæ et stricæ incrementi arcuatæ, unde interstitia leviter imbricata; apertura subovalis, intus candida, inferne fascia tosta intrante pellucente.

Apert. 8 m. lata, 10 alta; diam. anfr. ult. 12-13 m. (coll. Cuming, sine loco).

Hab. India orientalis.

A detached specimen with the impression of the ribs of a *Pecten*, small remains of a yellow coral, and an agglutinated dark-green spine of *Stella spengleriana*, Chemnitz, to which it was very likely attached.

This variety differs chiefly from the type in its great thickness and in having only a dark basal band inside, while the whole interior of the type is brown. The sides are pure white without longitudinal grooves. The under side shows, close to the mouth, a part of the interior dark band. On the upper side are four or five not very prominent arched varices, gaping a little in the front. From the var. γ of *S. nebulosum* it differs in its proportionately larger calibre, but chiefly in its yellow band and brown-coloured grooves, of which I have never seen any trace in the numerous specimens of its West Indian analogue.

PROC. ZOOL. SOC.—1861, No. XI.

17. SIPHONIUM NEBULOSUM, Dillwyn, 1817.

Tuyaux de mer, Davila, Cat. 1767, i. p. 100. t. 4. f. H (typus); Favanne, i. p. 652. t. 6. f. 3 (copy).

Serpula nebulosa, Dillw. ii. p. 1076. no. 19; Wood, Index, *Serpula*, fig. 18 (copy).

T. juvenilis planorbiformis, corrodens, castanea, aurantiaca vel alba, leviter spiraliter lirulata.

T. ætate media affixa, spiraliter varie contorta, alba pallide aurantiaca variegata, longitudinaliter lirata, superne carinis tribus squamiferis; striæ et rugæ incrementi sigmoideæ, inæquales; apertura circularis, libera, interne fascia basali intrante tosta vel anthracina, ferrugineo terminata.

T. grandæva libera, tubulata, leviter torta, teres, æqualis, antiquata, liris longitudinalibus obsoletis et subobliquis, annulatim rugosa et sulcata, inferne septo convexo obliquo terminata. Color candidus, intus juxta aperturam macula interrupta castanea.

Long. 80 m., diam. 10 m., diam. apert. 9 m.

Operculum pellucidum, flavesceus, superne concavum, laminis approximatim concentricis, inferne convexum; impressio muscularis magna, rufa, concentrice ruguloso-striata, linea coccinea peripherica terminata, vertice obtuso lato prominulo; limbus elevatus, nitidus.

Diam. 7 m.

The concentric *laminæ* seem to form a distinct upper layer, which can be separated with a knife. The spiniferous bodies are clavate, and of the same kind as shown in fig. 16, but more elongate, and dilated at the base. The surface is sometimes entirely occupied by a young specimen or a *Serpula*.

Var. α. SERRATA.

T. liris destituta, carina laterali squamis 6-7 subremotis, compressis, acutis, inæquilaterali-triangularibus; latere externo plano; fascia interna nigra ferrugineo terminata.

Diam. 6-7 m.

Hab. Honduras; on a branch of dead coral (coll. Cuming.).

Var. β. ACULEATA.

T. carina submediana, squamis 7-8 compressis, angustis, longis, liris duabus aculeato-squamosis in regione umbilicali, unde apertura superne trispinosa, latere externo liris obsoletis nodulosus, inferne transversim rugosa, latere affixo glauco et castaneo variegato; intus fascia nigra tosta.

Diam. ap. 5 m.

Hab. Honduras (coll. Cuming.).

Var. γ. TURBOIDES.

T. liris 6 subæqualibus, leviter nodulosus.

Serpula bicarinata, Sow. Man. f. 4; Humphrey, Conchology, pl. 10. f. 9 & 10.

Vermicularia arenaria, var. *a*, *b*, *c*, Gravenh. Tergestina, p. 51 (ex specim. orig.?).

Var. *δ*. IMBRICATA.

Serpula? *dentifera*, Sow. Gen. f. 6.

Vermetus dentiferus, Gray, Philos. Trans. 1833, p. 816.

Var. *ε*. PLANORBOIDES.

T. affixa, carina submediana acuta, squamis remotis munita; regione umbilicali bilirata, latere externo lirulis 6-7, interstitiis lirula intercalante.

Diam. 20 m., diam. ap. $5\frac{1}{2}$ m.

Hab. St. Thomas, on *Avicula columbus*, Bolt.

Var. *ζ*. RUGOSA.

T. antiquata et annulatim rugoso-striata; anfr. primi liris granulosis, aperturam versus evanescentibus.

Diam. 7 m.

Humphrey, Conchology, t. 10. f. 8.

Var. *η*. ANAULAX.

T. lris carens, lamina variciformi pone aperturam ampliatam, intus fascia basali castanea.

Diam. apert. 9 m.

Hab. Honduras (coll. Cuming.).

Var. *θ*. FISSURATA.

Anfr. ultimus conferte et regulariter annulatim rugosus.

This variety looks very like a *Siliquarius*.

Serpula dentifera, Lam., no. 24, var. *b*, testis majoribus subsolitariis (Mus.), and var. *d*, testis subrugulatis, glomeratis, belong very likely to this species; it would at least be very remarkable if this exceedingly common shell were not known to Lamarck.

18. SIPHONIUM GÆDEROPI, Mörch.

Serpula (an *Vermetus*?) *turboides*, Chiereghini, Venez. 1847?; Chenu, Monogr. du Spondylus, pl. 12. f. 3?.

T. solitaria, contorta, spira affixa; anfr. ultimus solutus, costulis spiralibus 1-3 squamis compressis remotis munitis, interstitio dorsali bilirato; stricæ incrementi subtilissimæ, arcuatim flexæ, huc illuc antiquatæ vel rufæ. Color albus vel sordide carneus.

Operculum superne profunde concavum, rufum, laminiis spiralibus confertis brevissimis appressis flavis, unde annulis angustis alternantibus flavis et rufis, centro profunde immerso fundo lato; inferne convexum; impressio muscularis obscura, concentrice rugoso-striata, sulco lato plano pallido terminata, centro obtuso lato prominente pallido, limbo latissimo nitido, lineis 4 flavis concentricis.

Diam. $2\frac{3}{4}$ m.

Var. α , *costulis æqualibus, circiter 12-14, approximatis, asperis, oblique tortis, præsertim in regione umbilicali, ubi squamifera; apertura circularis leviter constricta.*

Var. β , *carina mediana prominente.*

Diam. apert. 5 m.

T. juvenilis candida crystallina, pallide fusco tincta, verrucis cylindricis munita (Algir. Guyon, Mus. Reg.).

On *Spondylus gæderopus*: probably from Spain.

The young shell is of a white, chalcedony-like appearance.

19. SIPHONIUM LURIDUM, Mörch.

T. contorta, tenuiuscula, subdiaphana, pallide lurida; anfr. teretes, rapide crescentes, soluti, angulatim contorti, longitudinaliter striati et lirati; liræ prominulæ, circiter 20, squamis fornicatis compressiusculis, præsertim aperturam versus, validis, interdum transversim confluentibus; striæ incrementi in interstitiis costarum leviter arcuatæ; anfr. primi graciles, laqueati, albi, sculptura evanescente; apertura circularis, soluta, faucibus strato tenuissimo sericeo obtectis.

Diam. apert. 18 m.

Hab. Society Islands (coll. Cuming.).

The colour and sculpture remind one of certain varieties of *Pinna pernula*, Ch. The longitudinal flat and feeble *liræ* are almost obsolete on the scales; there are about three to six between the keels. The inner thin layer scarcely conceals the outer brown colour. Some of the keels show small indistinct white spots. The sculpture on the infero-marginal side is entirely rubbed out, and shows the cast of some foreign body: on one of the first whorls are distinctly to be seen the cast and rests of the cells of a *Bryozoon*. The lid is unknown, and the genus of course a little doubtful.

20. SIPHONIUM MARGARITARUM, Val.

Vermetus margaritarum, Val. Voy. Venus, pl. 11. f. 2 (the larger).

Vermetus margaritaceus, Rouss., Chenu, Ill. t. 4. f. 2 (the same).

Aletes margaritarum, Carp. Cat. Reigen, p. 303.

Siphonium margaritarum, Mörch, Journ. Conch. 1859, p. 359.

Hab. Panama (Val.); Mazatlan (Reigen).

I refer this species to the genus *Siphonium*, on account of the circular mouth, and some likeness to the much smaller *S. nebulosum*, Dw. The lid is unknown. In Mr. Cuming's collection there is a shell, without locality, very like the quoted figure; but the aperture is much higher than broad, of an oblong trapezoid shape, and the outer basal angle of the aperture is very sharp, while the upper is obtusely rectangular (alt. max. obliq. 22 m., lat. 14 m.). In the interior is a broad blackish basal band spreading on the sides like that of *S. nebulosum*, but broader. This shell will very likely prove to be generically distinct, as all the specimens of *Siphonium* I have seen have a circular mouth, depending on the large operculum.

21. SIPHONIUM SUBGRANOSUM, Mörch.

Siphonium costale, "Lam.," Mörch, Journ. Conch. 1859, p. 358.

T. crassiuscula, spiraliter contorta, spira affixa, alba vel flavescent; anfr. teretes vel latere inferno dilatati, longitudinaliter conferte lirulati; liræ subæquales, sæpe 5-6, prominentiores, lira mediana plerumque valida prominens; interstitia lirula intercalante; striæ et lamellæ brevissimæ, incrementi sigmoideæ, regulares, unde liræ subtiliter granulosa et in locis occultis minutissime undulato-squamosa; apertura circularis, libera nec soluta, intus candida.

Diam. circ. 5 m.

Operculum superne concavum, laminis confertis concentricis lateris, rufum, zona submediana flava, centro nudo excavato; inferne convexum, impressione musculari parva, cruentata, concentricè ruguloso-striata, linea coccinea terminata, centro conico valido prominente, apice obtuso leviusculo pallide flavescente; limbus convexiusculus, nitidus, bicolor, dimidio interno flavo, externo coccineo.

Diam. fere 7 m.

Hab. India orientalis (Tranquebar?), on *Malleus vulgaris* (L.), *Delphinula distorta* (L.), *Voluta lapponica* (L.), and *Margaritifera inflata* (Schum.).

Var. α. CARINATA.

T. affixa, spiraliter torta, umbilico lato, subtilissime conferte lirulata; liræ æquales, subtilissime granulosa, carina valida compressa mediana, squamis remotis obsoletis, lira prominula in regione umbilicali. Color flavescent.

Diam. ap. 8-9 m. (Mus. Reg.).

Var. β, varice erecta intus incrassata.

Var. γ, triquetra.

T. tenuiuscula, decliviter contorta, umbilico clauso lira prominula circumdata, unde anfr. subtriquetris, longitudinaliter pulcherrime lirulatis et granulosis, interstitias lirula intercurrente; anfr. primi teretes, planorbiformes; apertura circularis, contracta.

Diam. ap. $3\frac{1}{2}$, diam. anfr. ult. 5-6 m.

Operculum pallidum unicolor (desiccatione verisimiliter), superne laminis solutis concentricis, unde uti duplicatum, limbo obscuro, centro infundibuliformi fundo excavato; inferne mammilla valida, conica, prominente acutiuscula.

Diam. $3\frac{1}{2}$ m.

Several specimens on an *Avicula* resembling *A. ægyptiaca*, Chemn. xi. f. 2019, but posteriorly deeply inflected.

This species is not very unlike *V. costalis*, Chenu, t. 10. f. 6; but I have never seen any specimen of that size with such broad grooves, nor with a protracted last whorl.

22. SIPHONIUM CARINIFERUM, Gray, 1843.

Vermetus cariniferus, Gray, Dieffenbach, New Zealand, p. 242.

"Shell thick, irregularly twisted, opaque white, with a high compressed wavy keel along the upper edge; mouth orbicular, with a tooth above it, formed by the keel. Operculum orbicular, horny."

New Zealand, Parengarenga, North Cape, North Island (*Dr. Dieffenbach*).

According to the description of the operculum, *Dr. Gray* is no doubt correct in referring this shell to the *Vermetidæ*. From the dorsal keel I suppose it must be a *Siphonium*, which is the only genus among the *Vermetidæ* which has this character, most common among the *Serpulidæ*.

23. SIPHONIUM MAXIMUM, Sow., 1825.

Serpula maxima, Sow. Tank. Cat. App. p. 93. n. 23.

"*T. irregulariter contorta, læviuscula, carina dorsali antice in spinam nonnunquam desinente; apertura expansa, testarum juniorum subtrigona, operculo corneo.*"

Several specimens exist in collections; they are attached to, and some of them deeply imbedded in coral. It is a very large species, the tubes being almost an inch wide. The aperture in the older shell is round (*Sow.*).

Vermetus maximus, Gray, Spicileg. 1829, 1. p. 3, cum fig. anim.

"*T. junior adhærens, irregulariter contorta, subtriangularis, dorso carinato, superius erecto, rotundato; apertura orbicularis; peristomate tenui*" (*Gray*).

Pacific Ocean, with coral, *Stutchbury* (*Mus. Brit.*).

Vermetus gigas, Gray, Fig. of Moll. iv. p. 28. t. 128. f. 1.

Siphonium maximum, ib. p. 82.

Serpula maxima, Brit. Mus. Wood, Index Supp. t. 8. f. 2 (8 inches).

I have never seen a triangular young shell with a prominent keel running out into a spine; such a shell is, however, represented by *Wood*, creeping on the old individual; perhaps it is a *Serpula*. The figure of *Wood*, from a specimen in the British Museum, is probably not only the very specimen of *Gray's*, but also of *Sowerby's* description.

Var. α. *T. grandæva, repens, aperturam versus ampliata, superne undulato-varicosa, varice præcipitante juxta aperturam; striæ incrementi membranaceæ, confertissimæ.*

Diam. aperturæ 33 m.; diam. cavitatis postice 14 m., long. cavitatis circiter 230 m.; diam. max. tubæ antice 40 m., postice 20 m. (coll. Chr. viii.).

Vermetus gigas, A. & G., Chenu, Ill. t. 5. f. 6 a (but smaller).

Var. β, *teres; læviuscula.*

*Karang surumbung**, Javanorum, *Junghuhn*, Java, 1854, iii. p. 68. (tabula ultima).

* *Karang* signifies a tube or quiver, and *surumbung* a coral reef or rock; the former is probably the same as the Malayan *cappang*, according to *Rumphius*, used for *Cuphus* and *Tenagodus*.

T. adulta elongata, tubulosa, recta aut subflexuosa, e partibus lateralibus septorum concreta, inde concentrice lamellosa et stratificata, crassa, extus transversim striata et subannulatim rugosa et aspera, sæpius subtuberculosa et inæqualis, intus lævissima, nitida (testa junior ad basin infimam attenuata et spiraliter torta); septa transversa, remota, imperforata, tenuissima, tubuloso-hemisphærica (cupuliformia) se invicem amplectentia (superiora inferioribus incumbentia); partes laterales in testam concretæ; basis septorum libere prominula, clausa, hemisphærica, papyraceo-tenuissima, fragilis, rarius hemisphærico-conica et subobliqua; habitus orthoceratitoides, sed siphon nullus (Junghuhn).

Diam. testæ 8-14 lin., cras. testæ $\frac{3}{4}$ -2 lin.; septæ internæ diam. 6-7 lin.

According to the inhabitants this shell is only found in Java, at Karangitam, which forms a part of the broad reefs at the mouth of the small river Kalapatjondong, in the district Djampong, on the south coast of Java. The tubes, which at flood-tide are covered with 8 feet of water, stand upright in slightly bent positions, like organ-pipes, so closely together on the rock which they form, that it is only by touching the many small holes of the apertures that their presence is recognized. To this variety belongs very likely the shell brought from Sumatra by Capt. Griffiths, and described by Home, Phil. Transact. 1806, pp. 276 & 287 (Woodward, in Proceed. 1857, p. 245).

Vermetus gigas, Owen, Mag. of Nat. Hist. 1838, Aug., p. 408, f. 20.

Vermetus gigas, A. & G., Chenu, Ill. pl. 5. f. 6.

Siphonium gigas, Chenu, Man. p. 320. f. 2304 (copy).

Serpula gigantea *, or great Oriental Worm-shell with its operculum: a young specimen (Seba, t. 94), Port. Cat. 1786, p. 6, no. 97.

The figures of Owen and Chenu seem to unite Junghuhn's shell with the following form, which is the typical:—

Var. γ . SCANDENS.

T. contorta, inter ramos Heteroporæ scandens; anfr. teretiusculi superne et inferne subplanati, rugis transversis latis obsoletis arcuatis, varicibus crassis præcipitibus 1-2; striæ incrementi membranaceæ, confertæ, sæpe ferruginæ; apertura circularis, intus candida, peritrema intus carneum.

Diam. 21 m.

Operculum pellucidum castaneum; superne profunde concavum, area centrali opaca, periphæria elevata; limbo lævissimo, margine angusto flavo, uti cote attenuato; inferne elevato-convexum; impressio muscularis rotundata, prominula, concentrice interrupte rugulosa; limbus planissimus, latus, nitidus, margine interno canali plano angusto terminato.

Diam. $18\frac{1}{3}$ m., lat. limbi 5 m., alt. operc. circiter 7 m.

* The middle figure on pl. 97 of Seba is very likely meant, which perhaps is a *Cuphus*. The specimen was obtained by Sykes for 9s. at the auction.

Hab. Lord Hood's Island (coll. Cuming.); a specimen twisted among the branches of a dead *Heteropora*, agreeing very well with Wood's figure, which distinctly shows close to the mouth one of the characteristic abrupt feebly excavated varices. The operculum of *Vermetus giganteus* from Guam, represented by Quoy and Gaimard, pl. 67. f. 14, differs in having an impressed line close to the periphery, and in having the muscular impression concentrically furrowed with lines of alternately different sizes. The operculum in Chenu's 'Illustrations' has a central red spot, which I have not observed in any of the present specimens. This shell shows a tendency to grow vertically, and unites thus the form described by Junghuhn with the other varieties, which are all creeping.

Var. δ . OPERCULO MEGACENTRO.

Operculum inferne convexum (nec elevato-convexum), limbo convexiusculo prominulo angusto; impressio muscularis magna, subtiliter concentrice striata.

Diam. $17\frac{1}{3}$ m., lat. limbi 4 m., alt. operc. 5 m.

Hab. Lord Hood's Island (without shell).

This lid is flatter than that of the preceding, and, although the diameter is only a little smaller, the central area is notably larger, and the brim consequently narrower; the exterior yellow peripheral band is marked by an elevated margin as if scratched thin. The concentric lines of the muscular impression are more feeble, sharper, and more regular towards the periphery; they are more distinct, and form small elongate areas.

Var ϵ . BIANGULARIS.

Karang surumbung, var., Junghuhn, Java, f. 3 (optime).

T. repens, corrodens, semiimmersa, aperturam versus lente crescens, utrinque obtuse angulata, lateribus planatis, superne planiuscula vel leviter concava varicibus arcuatis præcipitibus; striæ incrementi membranaceæ, plerumque ferrugineæ; apertura circularis (vel in specimine incompleto utrinque dilatata ex angulis). Color anfr. primi ferrugineus.

Spec. majoris long. cavitatis int. 120 m.; diam. ap. 15 m., alt. 14 m.

Spec. minoris long. cavitatis int. 90 m.; diam. ap. 15–16 m., alt. 13 m.

Operculum tenue, convexum, pallide castaneum, superne plano-concavum, laminis appressis concentricis obsoletissimis, uti farinosum; area centrali magna, obscura peripheria elevata; limbo obsoletissime radiato, peripheria pallide flava; inferne plano-convexum; impressio muscularis opaca, concentrice ruguloso-striata, læviuscula; limbus nitidus, castaneus, linea obscura interna terminatus, peripheriæ margine angustissimo pallido.

Diam. 15 m., limbi lat. $3-3\frac{1}{4}$ m., alt. circ. $2\frac{1}{2}$ m.

The upper side is covered with a farinaceous layer, which, seen under the microscope, seems to show some bodies of the same

character as those seen in fig. 16. In the brim of the under side are here and there some of the kind, shown by small sharp points, which look like imbedded sand-corns.

Hab. Lord Hood's Island; two associated specimens half sunk in the surface of a dead coral (coll. Cuming.).

The smaller specimen has the aperture incomplete, and shows on both sides an auriculated production formed by the lateral angles; the upper side of the aperture is inflected like the striæ of growth, whilst that of the adult specimen is straight. The first part of the whorls is entirely concealed by the coral which, perhaps, has overgrown it. This, however, is not clear, because a saw has been used to diminish the size of the coral.

Var. ζ. PRÆLONGA.

T. repens, corrodens, gracilis, fere per totam longitudinem æqualis, utrinque obtuse rectangulata, inferne elevato-convexa, tenuis, superne crassa, plana, lævis, sebacea, sordide alba, varicibus huc illuc parvis acutis oblique arcuatis; striæ et rugæ incrementi obsoletæ, arcuatæ, granulis oblongis obsoletissimis; anfr. primi laqueati conjuncti, lateribus inferne dilatatis; septa frequentissima, cylindraceo-convexa, tenuia; apertura parva, superne arcuata reducta, lateribus rectis.

Long. 64–82 dm., long. cavitatis 12 dm., diam. transv. aperturæ 10–11 m., crassitie superne 6 m.

Hab. Lord Hood's Island, a detached specimen; the exterior layer of the shell is wanting on the under side, which proves it has been burrowing. This enormously long and very narrow shell differs chiefly from the preceding by its smooth tallow-like, slightly shining surface; the striæ of growth are only a little foliaceous in unexposed places; it has nearly the same diameter throughout its length from the mouth, whilst var. ε increases somewhat more rapidly. The transition between the above-described varieties seems so evident to me, that I do not dare to distinguish any of them specifically, not even the last. An allied species from the island of Tubal, in the Red Sea, collected by Rüppell, is indicated by Leuckart, 'Neue Wirbellose Thiere,' 1827, p. 39.

VERMICULUS, Lister, Hist. Conch. 1688, t. 548.

Tubulus, Schwammerdam, Bibl. 1738, p. 50. t. 7. f. 5, 6.

Siphonium, no. 3, Browne, Jamaica.

Vermicularia, Lam. Prod. 1799; System, 1801; Bowd. Elem. 1827.

Vermetus, Gray, Gould, Adams, Stimpson, Carp., non Adanson.

Testa varie torta, juventute turritelloidea.

Operculum magnum, superne concavusculum, laminis concentricis suprapositis margine laciniato, limbo reflexili; inferne convexusculum, impressione musculari confertissime concentrice lirulata.

Stimpson has described the animal thus:—"Mantle fringed at its

margin with short filaments. The foot is very short and broad, dilated into rounded auricles anteriorly. The muzzle is broad, not cleft; the tentacula are short, conical, having their eyes at their exterior base. An elevated ridge runs along the back, becomes flattened into a membrane at the head, and passes round under the right tentacle, forming a kind of canal, near which is the anus. Its colour is a light brown, with spots and patches of black; viviparous. The young shell is helicoid and reversed. The operculum is corneous, black and hard on the inner, and lamellated on the outer surface. It is surrounded by a thin, membranous, flexible portion, about one-fourth its diameter; thus it is enabled to close its shell perfectly at the aperture, and yet to retreat far into the narrower whorls." This latter assertion is not quite correct; an operculum of *V. lumbricalis* β , has a diameter of $6\frac{1}{2}$ m., whilst the aperture of the shell has a diameter of 7 m.

Sowerby, in his 'Genera,' was the first to give a representation of the lid, but this is not very satisfactory. A drawing of *V. eburneus* from life by Prof. Oersted shows considerable differences, as will be described below.

24. *VERMICULUS TORTUOSUS*, Soldr. 1784.

T. laze contorta; *anfr. latere externo plano vel etiam concaviusculo, lævissimo, superne et inferne rectangulato, costa marginato, spira conica abbreviata (ex icone).*

1784. *Le Quadrilatère*, Fav. Cat. de Latour d'Auvergne, no. & f. 1412; Humphrey, Conchology, t. 11. f. 4 (typus).

Serpula tortuosa, Sol., Port. Cat. p. 184. no. 3939.

Vermetus costalis, Rouss., Chenu, Ill. t. 3. f. 1 & 1 b; Man. f. 2295.

Vermiculus tortuosus, Mörch. Journ. Conch. vii. p. 30.

Var. α . *UNICOSTALIS* (ex icone).

Vermetus costalis, Rouss., Chenu, Ill. t. 3. f. 1 a & c.

T. latere externo convexiusculo, liris duabus æqualibus, inferne costa prominula marginato, superne nudo convexo; aperturam versus strigis aurantiis transversis.

Var. β . *NIDIFICANS*.

T. tenuiuscula, fusco-lurida, strigis ferrugineis obsoletis transversis, spiraliter torta; anfr. uno latere contigui, obtuse quadrangulares, longitudinaliter lirati et confertissime striati; latere externo declivi, inferne acutangolo, carina pallida albescente maculis ferrugineis notata; superne liris tribus, lirula intercalante, angulo obtuso mutico; latere interno angulo juxta umbilicum lira gemina munito, liris duabus tribusve, lirula intercalante; regio umbilicalis conferte plano-lirata; apertura ampla, orbicularis, obtuse quadrangularis; spira gracilis, ferruginea; anfr. ultimus carinis obtusis tribus interstitiis striatis, anfr. cæteri lævigati, nitidi, medio carinis duabus subæqualibus, interne lirula suturali obsoleta.

Diam. ap. 11, alt. 12 m.; long. spiræ circ. 9 m.

V. costalis, Chenu, pl. 3. f. 1 *a*, quoad sculpturam; Humph. t. 10. f. 5, quoad formam.

Hab. Ins. Philippin. (*Cuming*).

Three specimens affixed to the branches of a furcate coral, not unlike *Trymohelia eburnea* in shape, but very likely allied to *Leptastrea*.

This species is not quite without doubt distinct from *V. lumbricalis*. The *V. indicus* (Rous. t. 3. f. 2 *b*) seems to be a transition. The spire of var. β is entirely different from that of fig. 1 in Chenu, but agrees with that of *V. lumbricalis*, var.

25. VERMICULUS SOLARINUS, Mörch.

T. solidissima, alba, strigis transversis pallide aurantiis in intersectionibus costarum saturatioribus; anfr. laxi, subquadrangulares, latere externo rectangulato, carinis tribus æquidistantibus crenulatis; interstitia fundo lævi nitido, longitudinaliter lirulata, liris duabus validis obsolete nodulosis; latere interno semitereti, conferte grosse-lirato, interstitiis profundis hic illic lirula intercalante; liræ incrementi regulariter subremotæ, intersectionibus litarum incrassatæ, unde superficie rudi; apertura latere externo rectangulato, interno arcuato.

Long. fere 27 dm.; diam. ap. 9–10 m.

Spira turritelloidea, nitida, alba, basi plana, concaviuscula, carina prominente marginata, superne carina acuta approximata; anfr. angulo inframediano carina acuta munito, inferne lirula suturali subocculata.

Long. fere 6 m.

Hab. Ins. Philippin. (*Cuming*). One specimen.

The first whorls are planorbiform. The spire is very like that of *Turritella carinifera*, Lam., and agrees in this respect with the vars. δ and $\delta\delta$, of the following species.

26. VERMICULUS LUMBRICALIS, L.

“*T. quasi cornea, teres seu versus apicem obsolete angulata, sensim angustior.*” L. Mus. L. Ulr. 689. no. 431.

Serpula lumbricalis, Linn. S. N. x. p. 787, no. 698; xii. p. 1206, no. 801.

Professor Liljeborg of Upsal has favoured me with a drawing of the original specimen in the Museum of the Queen. I do not know any figure corresponding exactly with it, except perhaps *V. indicus*, Chenu, Ill. t. 3. f. 2 *a*. It agrees very well with the var. *a* in size and form; but the keels are not continued to the mouth, and the longitudinal grooves, rather strong between the keels, are not represented in the drawing.

Var. *a*. DIAPHANA.

T. tenuis, diaphana, dilute castanea; anfr. laxè contorti, inferne carinis duabus leviter crenatis, maculis vel punctis obsoletis albis et castaneis, interstitio plano vel leviter excavato, liris

3-4 *approximatis*; *latere externo plano, læviusculo, obsoletissime lirulato, superne obtuse angulato, liris tribus approximatis, latere interno convexo subtilissime lirulato*: *spira elongato-conica, turritelloidea* (figura in Gen. Sowerbyi non absimili sed minore), *ferruginea, basi lata plana vel concaviuscula, carina prominente marginata, superne carina compressa approximata, versus suturam lira sursum evanescente*; *anfr. primi inferne carinis duabus subæqualibus*.

Diam. ap. fere 7 m.

Buonanni, f. 20 ? (aggregata?).

Operculum superne concaviusculum, flavescens, laminis concentricis suprapositis margine laciniatis; inferne convexiusculum, badium; impressio muscularis conferte concentrice sulcata, centro prominulo convexo; limbus bipartitus, parte externa flava flexili, interna nitida badia, juxta aream canali circulari plana.

Diam. circ. $6\frac{1}{2}$ m.

Coll. Cumingii, sine loco.

Var. β . CORNEA, Knorr, Vergn. t. 2. t. 13. f. 1 quoad colorem.

T. solidissima, badio et castaneo variegata, laxè contorta, longitudinaliter subtilissime lirulata; anfr. graciles (præsertim superne) *latere inferno plano, utrinque carina irregulariter subnodulosa, interstitio lira gemina mediana; latere externo inferne lira acutiuscula carinam approximante, superne angulo obtuso, liris 3-4 approximatis subæqualibus; latere interno pallido convexo, lirulis inæqualibus; striis et rugis incrementi sigmoides, aperturam versus antiquatis; spira deficiente sed verisimiliter ultra solitum parva; apertura circularis inferne incrassata.*

Diam. 7 m.; diam. septi (fracturæ) circ. $\frac{3}{4}$ m.; long. testæ $22\frac{1}{2}$ m.

Coll. Cumingii, sine loco.

Differs from the preceding by its great solidity, and chiefly in the outer side being of a very dark, nearly black-brown colour.

Var. γ .

"*Apertura biangulata.*"

Vermetus bicarinatus, Desh. Anim. s. Vert. 1843, ix. p. 67.

Var. δ . AMPLIATA.

Vermetus bicarinatus, Desh., Chenu, Leçons Elém. p. 182. f. 602.

V. indicus, Rouss., Chenu, Ill. t. 3. f. 2 *b* (idem).

T. tenuiuscula, dilute castanea, laxè contorta; anfr. approximati, subtilissime lirulati, superne graciles versus aperturam ampliati, oblique quadrangulares subgibbosi, latere interno declivi planiusculo, liris geminis in medio, superne carina acuta pallida, inferne carina obtusa subnodulosa terminato; latere externo superne obtusissime angulato; apertura transversa, ovata, latere interno angustato; spira turritelloidea et ei V. solarini simil-

lima; anfr. infimus inferne bicarinatus, anfr. 6 primi angulati, carina acuta, lira suturali subocculta.

Diam. ap. circ. $9\frac{1}{2}$ m.; alt 9 m.

A specimen very like the quoted figure is in Mr. Cuming's coll. (sine loco).

Var. ♂♂. RUGULOSA.

T. parva, alba, versus spiram pallidissime castanea, laxè contorta, subtilissime lirulata, latere inferno planiusculo livato, præsertim aperturam versus, angulo externo carina acutiuscula, punctis ferrugineis fere æquidistantibus notata, angulo interno obtuso; rugæ et striæ incrementi validæ, sigmoideæ, approximatae; spira uti præcedentis.

Diam. ap. circ. 5 m.

Hab. Ins. Philippin. (Cuming).

A specimen not unlike Gualt. t. 10. f. Q, attached by the aperture to another specimen chiefly differing in its uniform pale-brown colour, more slender whorls (diam. 4 m.), and very feeble sculpture.

The spire is of the form of *Turritella carinifera*, Linn., like the preceding. This var. and the preceding approach to *V. tortuosus*. Buonanni, f. 43 (spira parva) verisimiliter.

Var. ζ. TERES.

T. laxè contorta, latere interno albescente, externo badio, subtilissime longitudinaliter striata; anfr. 4 primi inferne plani, carina acutiuscula externa longitudinaliter lirata et conferte striata, latere externo longitudinaliter lirati et striati, superne obtuse angulati; anfr. ceteri teretiusculi, inferne obtusissime bingulati, rugis incrementi sigmoideis; apertura triangulari-ovata.

Alt. 10 m., diam. 9 m.

Spira turritelloidea gracilis; anfr. convexi, trilirati; liræ duæ medianæ in anfractibus primis æquales, lira superior minor in anfr. cæteris, evanescens in ultimo anfractu, lira tertia suturalis.

Long. 10 m.

Hab. Ins. Philippin. (Cuming), from the specimen represented in Reeve's Conch. Syst.

Vermiculus longus eleganti clavicula tortili desinens, List. p. 548. f. 1.

Tubulus anguinus, Mart. i. t. 2. f. 12 B (copy).

Serpula lumbricalis, Wood, Index, t. 38. f. 1, Amboina (copy).

Vermetus lumbricalis, Reeve, Conch. Syst. t. 152. f. 1.

Serpula, Humphrey, Conchology, t. 10. f. 9.

Vermetus lumbricalis, Desh., Cuv. Regn. An. t. 62. f. 3.

Vermetus indicus, Rouss., Chenu, Ill. t. 5. f. 2 a.

Hornschlange, Rumph. t. 41. f. 1.

Serpula lumbricalis, Burrow, Elem. t. 22. f. 2.

Vermiculus, D'Argenv. t. 29. f. 1.

Hab. Ins. Philippin. (H. Cuming).

Notwithstanding the very different turritelloid portion, I doubt

whether this is specifically distinct, the var. α showing a transition. A part of the keel shows some small, equidistant brown spots like those of the preceding variety.

Serpula vermetus of Sowerby's 'Genera,' seems to be most nearly allied to var. α , chiefly by the spire. The operculum represented in the mouth of the shell, and the much smaller free specimens, are very likely rather inaccurate.

27. VERMICULUS SPIRATUS, Phil. 1836.

Var. α . MELANOSCLERA.

T. solida, badia, strigis transversis atris; anfr. obtuse quadrangulares; latere externo medio angulato, carina pallida, longitudinaliter lirulata; latere interno fortiter lirato, apertura subquadrangularis.

Diam. 8-9 m.

Hab. Vera Cruz (*Rathsak*).

Var. β . QUADRANGULARIS.

Rufescens.

Vermetus quadrangulus, Phil. Mal. Zeitschr. 1848, p. 17. no. 77.

Hab. Yucatan (*Philippi*).

Var. γ . BICARINATA.

Hornschlange, Hühnerdarm, Knorr, iv. t. 17. f. 2.

V. spiratus, Phil. Wieg. Archiv. 1836, p. 244. t. 7. f. 5 a-c.

V. knorrii, Desh. Anim. s. vert. 1843, ix. p. 68.

V. lumbricalis, d'Orb. Hist. Nat. Cuba, p. 234.

V. lumbricalis, Bowd. Elem. p. 23. t. 9. f. 17. (1822).

V. d'adanson, Blainv. Man. 1825, t. 34. f. 1 (copy).

V. lumbricalis, Sow. Man. f. 345.

V. indicus, Chenu, Ill. t. 3. f. 2 (verisimiliter).

Hab. Havanna (*C. B. Philippi, d'Orb.*).

Var. δ . CINEREA.

Vermetus lumbricalis, Gould, Invert. Massach. p. 287.

Vermetus radícula, Stimps. Shells of New Engl. p. 38.

Vermetus radícula, Gray, Guide, p. 126.

Hab. New Bedford Harbour (*Gould*); Buzzard's Bay southward (*Stimps.*).

Var. ϵ . UNGULINA.

Spira magna, turritelloidea, spiraliter striata et trilirata; liræ duæ infimæ subæquales, majores; umbilicus perivius; color castaneus.

Long. 7 m., lat. bas. 8 m.; diam ap. circ. 4 m.

Vermetus lumbricalis, Woodward, Man. t. 9. f. 7, simillima.

Hab. Ins. St. Thomas (*A. H. Ruse*).

The spire of this shell is still larger, as in the quoted figure. I have never seen a depressed aperture, as in Woodward's plate; but a somewhat similar one is to be seen in Schwammerdam (t. 7. f. 5, 6).

It is very like in sculpture the last variety of *V. lumbricalis*. I have seen a specimen in which the three first whorls are angular and smooth, and the rest of the spire flat and trilirate.

Var. ζ. SCALARIS.

T. castanea, tenuiuscula, laxæ spiralis; anfr. teretiusculi superne et inferne subplanulati unde obtusissime biangulati, longitudinaliter lirulati et striati; rugæ incrementi regulariter approximatae, conferte undulatae; striæ incrementi interdum membranaceae; spira parva, turritelloidea, pallida; anfr. primi obliqui, infra medium acutanguli, anfr. ult. carinis duabus æqualibus, basi concaviuscula.

Diam. apert. 6 m.

Serpula, Humphrey, Conchology, t. 10. f. 5 & 7.

Hab. Ins. St. Thomas.

All the specimens show remains of sponges; they are all twisted in the same manner as the spongiolate *Tenagodi*. The transverse undulated rugæ remind one very much of the varices of *Cirsostrema*. To this variety belongs, according to the original specimen, *Vermicularia glomerata*, var. 2 c (Gravenh. Tergestina, p. 61 c).

Var. η. TERES.

Vermetus lumbricalis, Pictorial Museum, f. 2820.

Lumbricoid Vermetus, Knight, Encyclop. 1856, iv. p. 1134. fig. (copy).

Vermetus lumbricalis, Illustrirte Naturg. f. 3743. p. 193 (copy).

A group affixed on a *Chama* (probably *C. macrophylla*) is very likely a variety of this species.

28. VERMICULUS ANGUIS, Forbes.

Vermetus? anguis, Forbes on Fossil Invert. from Southern India; Geol. Trans. vii. 3. p. 124. t. 13. f. 1 (1845-56).

Uncoiled spiral of two loose turns of angulated rugose whorls, spirally striated and semiplicated.

Hab. Pondicherry, cretaceous (*Grant*).

The sculpture of this shell is very like that of *V. spiratus*, var. ζ.

29. VERMICULUS COCHLEIFORMIS, Jos. Müller.

Vermetus cochleiformis, Müll. Mon. Aachener Kreide, 2. p. 6. t. 3. f. 3.

Hab. Greensand of Königsthorn on Lusberg and near Vaels (*Müller*).

Not unlike in shape some forms of *V. spiratus*, var. ζ. This and the preceding have very likely lived in sponges.

30. VERMICULUS? CIRCUMCARINATUS, Stoppani.

Serpularia circumcarinata, Stop. Pal. Lombard. t. 6. f. 4-6.

Fossil in triassic formation. This shell is perhaps more allied to *Eccyliomphalus*, Portlock.

31. VERMICULUS TURRITELLA, Rousseau.

Vermetus turritella, Rouss., Chenu, Ill. t. 4. f. 3.

Vermetus turritella, Rouss., Chenu, Man. p. 319. f. 2298 (copy).

This shell is perhaps most allied to *V. lumbricalis*, var. β ; it is very likely not full-grown.

32. VERMICULUS ROUYANUS, D'Orb., 1842.

Vermetus rouyanus, D'Orb. Ter. Cretac. 2. pl. 233. f. 5-7.

V. rouyanus, D'Orb., Chenu, Man. p. 319. f. 2296.

Hab. Couches aptiennes ou terrain néocomien sup. des bassins méditerranéen et parisien (*D'Orb.*).

33. VERMICULUS ALBENSIS, D'Orb., 1842.

Vermetus albensis, D'Orb. Ter. Cret. 2. p. 386. pl. 233. f. 8-9.

Hab. Couches aptiennes des environs d'Evry (Aube) (*Dupin*).

The descriptions of the last two species do not seem to correspond with the quoted figures; but whether the names on the plate or the diagnoses are displaced I cannot decide. It is doubtful if these two species are not *Turritellæ*, as neither of them is known in a distorted state.

34. VERMICULUS CARINATUS, Hörnes.

Vermetus carinatus, Hörnes, Geol. Reichsanst. 3. p. 486. t. 46. f. 17.

Vermiculus triqueter, Dujard. Mém. Géol. ii. p. 283.

Vermetus adansonii, Deffr. Dict. Sc. Nat. lvii. p. 326 (nec Daud.).

Serpula vermeti aff., Sow. Gen. 1824, f. 4.

T. trochiformi-acuminata, irregulari, anfractibus lævibus medio acute carinatis, supremis trochleiformibus, ultimo ad aperturam coarctato, basi linea elevata instructa, apertura rotundata.

Long. 27 m., ap. 7 m. (*Hörnes*).

Hab. Tertiary formation of Steinbrunn and Lapagg (*Hör.*); Touraine (*Sow. et Duj.*); Thorigné near d'Angres (*Deffr.*).

Sect. B. *Animal tentaculis longis subulatis; pallio simplici.*

Operculum magnum planum. Testa anfractibus ampliatis.

Professor Oersted's drawing of *V. eburneus* differs from the description of Stimpson in the long subulate tentacula, and in not showing any filaments at the brim of the mantle.

35. VERMICULUS DIMORPHUS, Mörch.

Serpula, Humphrey, Conchology, t. 10. f. 6.

T. solida, spiraliter contorta, cinnamomeo cinereoque variegata; anfr. laxi, lateraliter connati et affixi, ampliati, teretiusculi, longitudinaliter confertissime striolati, latere externo convexiusculo, liris tribus inæqualiter remotis, latere infero convexo lirato, liris regionis umbilicalis approximatis, interstitiis lirula parva intercalante, latere superiore plano albo liris destituto; rugæ et striæ incrementi rectiusculæ, unde indistincte cancellata; apertura orbicularis vel subtriangularis.

Diam. 11 m.

Spira turritelloidea, candida, anfr. planis confusis, triliratis, lira mediana valida compressa acuta.

Hab. Ins. Philippin. (*H. Cuming*).

A specimen affixed to the tube of a *Rocellaria* (?). The sutures of the spire are very difficult to distinguish; it looks as if the two equal liræ between the two carinæ were situated on one and the same whorl.

? Var. α. LITUINA.

T. irregulariter planorbiformis, anfr. latere affixo plano per totam longitudinem; latere libero tereti, liris inæqualibus planis, lira peripherica lata; latere interno liris confertis, interstitiis lirula intercalante; rugæ incrementi irregulares, approximatae; anfr. ultimus flavus, strigis confertis confluentibus lividis; apertura orbicularis, subangulata.

Diam. 12–14 m.

Hab. Ins. Philippin. (*H. Cuming*).

A detached specimen, showing the cast of the denudated layers of a pearl-oyster, and exhibiting a feeble pearly lustre. Spire typical. I have found two specimens with about three detached whorls in a mass of *Petalococonchus flavescens*, Carp., said to be from Sicily; they only differ in the colour being of a dull white, and not glossy white as in the two forms described.

36. VERMICULUS PELLUCIDUS, Brod. & Sow., 1829. (Pl. XXV. figs. 17–20.)

T. pellucida, longitudinaliter striata, apicem versus carinata.

The striæ, which become carinated towards the apex, are best seen in the young shell. The species in texture has somewhat the appearance of *Magilus*, Sow. & Brodr.

Vermetus pellucidus, Brodr. & Sow. Zool. Journ. iv. p. 369.

Var. α. PLANORBOIDES.

Serpula regularis, Chenu, Ill. t. 10. f. 14.

T. tenuiuscula, irregulariter planorbiformis, lateraliter affixa; anfr. longitudinaliter pulcherrime dense striati et plano-lirati, liræ in anfr. primis fortiores, latere interno plano; apertura obliqua orbicularis, latere interno affixo.

Diam. 9 m.

Spira turritelloidea, anfr. vix convexiusculi, carina valida compressa acuta submediana, superne juxta suturam lira prominula et inferne lirula suturali in anfr. primis minutissima.

A specimen without locality in Mr. Cuming's collection, a little smaller than the quoted representation; the affixed side shows some agglutinated shell-rudiments, and two or three flat marks, very likely, from the lustre, those of a *Margaritifera*.

Var. αα. LAQUEARIS.

T. ut in præcedente, sed anfractus connati teretiusculi; anfr. ultimus

PROC. ZOOL. SOC.—1861, No. XII.

ampliatus, tenuis, resupinatus; apertura orbicularis latere interno connato; rugæ incrementi irregulares, confertæ.

Diam. ap. 15 m.

Hab. West Columbia (*Cuming*).

This shell differs from the preceding chiefly in having a whorl more, which is bent back and attached by the aperture to the penultimate whorl, giving it the appearance of the figure 8 or letter S.

Var. β . CINNAMOMINA.

T. irregulariter planorbiformis; anfr. hic illic soluti, latere affixo plani, longitudinaliter subtilissime lirulati, et regulariter lirati, lira peripherica latiuscula; rugæ incrementi irregulares, unde obsolete nodulosæ in intersectionibus; anfr. ultimus aperturam versus pallide cinamominus, strigis transversis obscuris: spira ut præcedentis.

Diam. ap. 14 m.

Hab. West Columbia (*Cuming*).

On the affixed side are agglutinated rudiments of shells and parts of an ochraceous stone, &c. This variety is very like the var. α of *V. dimorphus*, M.; it differs chiefly from the latter in its greater calibre and thinner shell. It is not impossible that the var. *lituina* belongs to this species, and that the locality Philippines is a mistake, particularly as the two small liræ on the spire are not quite equal, as in the three typical specimens of *V. dimorphus*.

Var. γ . VOLUBILIS (Pl. XXV. figs. 18 and 19).

T. eburnea, subobesa, laxè volutata, longitudinaliter costata, costis distantibus, subobsoletis.

Long. $\frac{3}{16}$, diam. $\frac{1}{16}$ poll. (*Reeve*). "Mus. *Cuming*."

Vermetus eburneus, *Reeve*, Proc. Zool. S. 1842, p. 197; *Conch. Syst.* 2. p. 46. pl. 152. f. 2.

Lumbricoid Vermetus, *Pictorial Museum*, f. 2820.

Vermetus lumbricalis, *Knight*, *Encycl.* p. 1134; *Illustrirte Naturg.* f. 3741.

The three latter representations from the aperture are from the same specimen shown from the dorsal side in *Reeve's Conch. Syst.*

Var. δ . VOLUBILIS (ADULTA) PICTA.

Vermetus eburneus, *Carp.* Cat. p. 304.

T. candida, in parte anfr. ultimi flavescens, strigis lividis (6-7) transversis confertis; anfr. ultimus aperturam versus ampliatus.

Long 35 diam., diam. testæ 18-20 m.

Hab. West Columbia (*Cuming*). Two aggregated specimens. *Puntarenas (Oersted: Mörch, Journ. de Conch. viii. p. 30).*

Var. ϵ . CRASSA.

Serpula panamensis, *Chenu*, Ill. t. 10. f. 5.

Vermiculus eburneus, *Mörch*, J. de Conch. viii. p. 30.

Liræ prominentes: animal viride, albo maculatum, unde reticulatum,

tentacula longa subulata, interrupte-lineata; proboscis brevis rotundata.

Hab. Puntarenas (Oersted).

The diameter of the aperture is 10 to $10\frac{1}{2}$ m.; the thickness of the shell 2 m. Fossil at Newburn, N. Carolina (Carp.).

Var. ζ. TIGRINA.

T. solida, sordide flavescens, strigis transversis badiis confluentibus; anfr. obtuse quadrangulares, longitudinaliter expresse lirati, latere externo liris latioribus, latere superiore albo, liris destituto; lira peripherica lata; rugæ incrementi prominentes, confertæ, unde interstitiis hic illic foveolatis; apertura circularis, faucibus fuscis: spira sic ut præcedentis.

Diam. ap. circ. 16 m.

Hab. West Columbia (H. Cuming).

Four agglomerated specimens of different size, showing on the flattened attached side remains of a grey sandstone. The colour, very like that of *Turritella tigrina*, Kien., gives this variety a very strange aspect; but the nearly white var. δ shows on a part of the whorl the same colour, but feebler.

Var. η. CASTANEA.

Versus aperturam strigis albis.

Diam. ap. 7 m.

A planorbiform specimen on a spine of *Murex melanoleucus*, Mörch. The numerous minute striæ which cover the liræ as well as the interstices, is nearly the only constant character.

The operculum is well described by Carpenter (Cat. p. 304) from specimens in Mr. Cuming's collection; but I have only found one specimen, labelled West Columbia, without shell; perhaps another may be found in the specimen figured by Reeve. I have not been able, even under the microscope, to discover the punctated lines mentioned by Carpenter.

Operculum concaviusculum, laminis 3-4, latis, confertis, laceris, radiatim striatis, area centrali parva arcu spirali leviuscula; inferne planum, badium; impressio muscularis lira elevata peripherica terminata, confertissime concentricè lirulata; limbus nitidus tripartitus: zona interna lineis tribus concentricis, zona mediana latior convexiuscula, externa pallida tenuis flexilis.

Diam. 9 m.; diam. impress. musc. $2\frac{3}{4}$ m.

Hab. West Columbia (H. Cuming).

Var. ι. from var. δ.

Hab. Puntarenas (Oersted).

Diam. apert. 15-16 m.

Operculum superne planum, laminis confertis (detritis?) brevissimis; impressio muscularis liris concentricis confertis subirregularibus, lira elevata terminata, centro immerso; limbus nitidus, tripartitus; zona interna angustior excavata, externa pallida angusta subreflexa.

Diam. 12 m., impress. musc. $4\frac{1}{2}$ m.

Vermetus hindsii, Gray, Adams, Gen. t. 38. f. 8 a, b?

Var. 2. *DISCIFER*, from var. δ .

Hab. Puntarenas (*Oersted*): a smaller specimen.

Operculum superne laminis brevibus detritis, centro disco pallido prominulo plano, concentricè confertissime striolato; inferne planum, badium; area centralis irregulariter concentricè lirata.

Diam. 11 m., diam. impress. musc. fere 5 m.

Var. 3, from var. ϵ , of which the animal is described, fig. 17.

Diam. 10 m., diam. impress. musc. $3\frac{1}{2}$ m.

The muscular impression is proportionately much larger than that of the first described, which is much thinner.

Var. 4. *SUBGRANOSA*, from var. η .

Operculum superne concaviusculum, lamina concentrica juxta peripheriam, laminis medianis oblitteratis, inferne badium; impressio muscularis conferte granuloso-rugosa (an morbo?), canali irregulari circumdata; zona interna rugis obliquis approximatis.

Diam. 6 m., impress. musc. $2\frac{1}{2}$ m.

On the surface are some cylindrical detached bodies like blunt spines, which look as if they were broken off at the base, but I believe they scarcely belong to the operculum.

Operculum testæ juvenilis.

Diam. aperturæ 3 m.

Hab. Puntarenas (*Oersted*).

Op. inferne spiraliter liratum; impressio muscularis indistincte terminata, annulo centrali.

Diam. $2\frac{1}{2}$ m.

The operculum is very variable, it varies from black to pale brown; the channel surrounding the muscular impression becomes deeper with age, whilst it is very indistinct in the young state. The radiating striæ of the *laminæ* form small denticles in the border if looked at under the microscope. The operculum of *Serpula panamensis*, represented by M. de Serres, Ann. Sc. Nat. 1855, v. p. 243. t. 8. f. 7, is perhaps of this species; but fig. 8 is very likely erroneous.

Sect. C. *Spira minuta helicoidea*.

The two following species are referred to this genus on account of the operculum, notwithstanding the very different form of the shell.

37. *VERMICULUS RASTRUM*, Mörch.

T. lateraliter affixa, laqueatim contorta, alba pallide flavescens; anfr. semiteretes, inæquales, marginibus dilatatis, superne liris 10-12 compressis, squamas confertas fornicatas amplexentibus, interstitiis lira plana interdum duplicata, squamis confertissimis arcuatis brevibus, anfr. ultimo juxta aperturam libero, cylindrico, coarctato; apertura circularis.

Operculum superne concaviusculum, laminis laciniatis confertissimis, centro conferte lirulato læviusculo; inferne planiusculum, badium;

impressio muscularis conferte lirata, opaca, periphæria leviter impressa; limbo nitido tripartito; zona interna sulcis 2-3 spiralibus, mediana parva vix sulcata, externa tenuis flavescens reflexa.

Diam. $9\frac{1}{4}$ m., impress. musc. fere 3 m., apert. testæ 9 m.

Aletes not unlike *A. squamigerus*, Carp. Cat. p. 304, lin. 28 (coll. Cuming.), sine loco.

The operculum is very like that of *V. eburneus*; but the muscular impression is not limited by an elevated line and flat channel. The affixed side shows four or five diverging channels, very likely the impression of the branches of a *Gorgonia*: on a part where the outer shell is broken is to be seen a very long oblique septum, perhaps pathological. The first whorls are lost; but a specimen from Puntarenas (*Oersted*) seems to correspond with it very well, only the scales are much feebler; its nuclear shell is helicoid and glossy. Another specimen from the same locality, perhaps even the latter whorls of the same specimen, has six to eight marked liræ, with three close-set small liræ in the interspaces. The scales are very numerous, small, and close-set; the colour is ferruginous, variegated with a foreign green hue.

Diam. of aperture 6-7 m.

The striæ of growth of *V. eburneus* are sometimes raised; but I have not seen them forming scales.

38. VERMICULUS EFFUSUS, Valenc.

Vermetus effusus, Val., Chenu, Ill. pl. 5. f. 4 a-c; Mörch, J. de Conch. viii. p. 38.

Siphonium effusum, Chenu, Man. p. 320, f. 2301.

Fig. 4 of Chenu is taken from the very same specimen represented as *V. centiquadrus* in 'Voy. de Vénus'; the specific name seems best to suit fig. 4 a. The lid, if represented of the natural size, seems too large for the shell.

EXPLANATION OF PLATE XXV.

- Fig. 1. *Stephopoma tricuspe*. Spine of operculum.
2. *Stephopoma senticosum*. Spine of operculum.
- 3, 4, 5. *Stephopoma pennatum*. Shell.
- 6, 7. *Stephopoma pennatum*. Operculum.
8. *Stephopoma pennatum*. Spine of operculum.
9. *Stephopoma pennatum*, var. *bispinosa*. Operculum.
10. *Stephopoma pennatum*, var. *bispinosa*. Spine of operculum.
11. *Stephopoma roseum*. Spine of operculum.
12. *Siphonium (Dendropoma) megamastum*. Operculum.
13. *Siphonium (Dendropoma) megamastum*. Spine of operculum.
14. *Stephopoma senticosum*. Young shell.
15. *Siphonium (Stoa) subcrenatum*, var. *spinosa*. Operculum.
16. *Siphonium (Stoa) subcrenatum*, var. *spinosa*. Spine of operculum.
17. *Vermiculus pellucidus*, var. *volubilis*.
- 18, 19. *Vermiculus pellucidus*, var. *volubilis*. Shell.
20. *Vermiculus pellucidus*, var. *volubilis*. Operculum.