neck, extending beneath the clavicle, and an incision in the esophagus, in which he found the teeth embedded deeply in the walls with granulation tissue around them. The teeth were extracted and for a while the patient's food passed through this esophageal opening, but it gradually filled up and the man to-day is practically well without evil consequences. If he had pulled with more force, the patient might have been injured. Last year a man came to him suffering from great pain and difficulty in swallowing, with edema of the posterior part of the larynx and esophageal opening, a history of having swallowed half of the wishbone of a chicken. By simply making the patient say o, a, e, and moving the larynx, he succeeded in seeing a white substance, and with the forceps seized the bone. It was two inches long, with sharp edges. It had penetrated the tissues between the opening of the esophagus and the interarytenoid space and extended through the neck almost to the skin. The man had edema of the larynx, and for several days it was a question whether tracheotomy would not have to be performed, but he finally recovered. About a week afterward a woman, a friend of the last patient, while sitting at the table joked about his swallowing the chicken bone, when she swallowed one herself. Her physician sent for Dr. Myles and he removed part of a breast bone from the upper part of the esophagus. It was about as large as a 25-cent piece, with sharp edges. In both cases the swallowed bones were hidden in the ordinary position of the larynx, but in the act of moving the throat, singing and making noises, they could be seen. A few years ago he had an interesting case in a patient who swallowed a piece of the joint of a bone that had been slashed with a knife and was part bone and part gristle, about as big as a 50-cent piece. He engaged it with an umbrella bougie, but could not turn it loose. He held on to one end of the instrument until finally she jerked and pulled bone and instrument out together. It was too large to come without tearing its way out. Fortunately, she got well. He supposed it was high enough up not to have done much damage. It would seem better to rely on general surgery when teeth or anything of that kind is swallowed, because if too much force is used, it may easily happen that the parts are torn irreparably.

DR. C. M. ROBERTSON, Chicago, spoke of spasm of the larynx in cases of foreign bodies in the esophagus. The lower end of the esophagus belongs to the general surgeon, but the laryngologist's territory extends down to the bifurcation of the bronchus. He had several cases in elderly people who had swallowed large pieces of food, meat or something of the kind, which usually lodges in the first ten inches from the teeth. As soon as the bolus of food gets into the esophagus there is a contraction of the muscles, both above and below the foreign body, and these muscular contractions are so strong that it is almost impossible to remove the foreign body. That has given rise to a so-called stricture of the esophagus when no stricture really exists. In these cases a large bougie will do better than a small one as the contraction of the muscle is overcome by greater pressure. If it does not resolve in some such way spasm of the larynx occurs as a secondary condition and patients sometimes succumb to spasm of the larynx from a foreign body in the esophagus. He thought the surgery in these cases should be limited to dealing with the diverticula. If the foreign body is small the use of the tube with illumination would be the proper treatment. If the foreign body is too large to remove by forceps through the tube then the use of the bougie is indicated.

DR. H. P. MOSHER, Boston, said that he tried not to mix general surgery with special work on the upper end of the esophagus. A good deal of encroaching on general surgery may be legitimately done in the upper end of the esophagus; for instance, in carcinoma. He had the good fortune to be able to curette down a polypoid growth at one sitting with esophageal tubes. He does more in one sitting than can be accomplished in many sittings by the other methods. If anything can go through a stricture a string can; it has been found that a string will find its way where nothing else will. It takes from three or four days to a week or more to find its way down. The fact that coins usually lie transversely in the esophagus emphasizes the fact that the esophagus is flat at the beginning and dilates best transversely. In removing coins many old

instruments will work well. That is illustrated by an amusing experience Dr. Mosher had with a foreign body, a coin in the upper end of the esophagus. He passed a tube directly by it. His house officer later took an old-fashioned corn catcher and presented the coin to him. He had had no personal experience with electrolysis, but was glad to know that it has worked in the cases spoken of. It should be remembered that in sword swallowing the sword has six inches to go before it gets to the esophagus. Somebody might take an x-ray picture and see how far it really goes. He did not, by any means, advocate the use of the speculum for passing the tube. The tube can be passed under cocaine very readily, in the majority of cases. His instrument is to be used when it is necessary to see the pyriform sinus and the beginning of the esophagus, and to do definite work in this locality.

MODIFICATION OF THE SIMPLE MASTOID OPERATION WHICH SHORTENS CONVALESCENCE BY FACILITATING WOUND REPAIR.*

DRS. E. M. PLUMMER AND H. H. GERMAIN.

Professor of Otolaryngology and Assistant Professor of Anatomy, respectively, in Tufts College Medical School.

BOSTON.

It is the purpose of this paper to draw attention to a new method of operative procedure on the mastoid process, which we have employed with most successful results. We will briefly describe its technic, consider its peculiar problems, and state in detail such personal experience as will give a clear idea of its theory and practice.

For a long time we had been convinced that the simple mastoid operation now in vogue for the mechanical relief of acute suppurative mastoiditis, with its accompaniment of unremitting pain, was greatly inadequate, and that there must be some means of improving on it. We refer especially to the long and tedious convalescence that invariably follows it, as shown by the statistics of the duration of wound repairs in our large aural clinics. Even if the case goes well, the time before cicatrization takes place varies from 4 to 8 weeks, and very frequently we are called on to do a secondary operation.

Under any circumstances, prolonged convalescence must be regarded as a hardship; but when the patient is poor it may be a serious calamity. Whether or not he be a worker himself, an additional burden falls on the working members of his family; worse still, if it be the chief breadwinner who is thus incapacitated, the result can be only suffering and privation for those who are dependent on him.

An operative process, then, that can mitigate such conditions, or the possibility of such, is unquestionably a blessing. Fortunately for humanity, the duration of wound repair in the field of general surgery is much shorter than in the field we are considering. We now observe complete recovery after removal of the breast, with extirpation of the contents of the axillary space, in 14 days, and in cases of amputation of the thigh the surgeon deplures having to detain the patient in the hospital over the third week.

Believing that the unfortunate contrast to these conditions furnished by all cases of mastoid surgery was due mainly to a defect in the accepted method of operation, we set ourselves the task of studying its problems

*Read in the Section on Laryngology and Otolaryngology of the American Medical Association, at the Fifty-seventh Annual Session, June, 1906.

with a view to remedying this defect if possible. Accordingly, we made a series of experiments on the cadaver in the anatomic department of Tufts College. Its result was an operation which seemed so rational that there appeared to be ample justification for performing it on the living subject. Following is a brief description of its technic.

TECHNIC.

The usual cutaneous incision is made posterior to and parallel with the general direction of the auricle, extending from the extreme tip of the mastoid process upward to a point $\frac{1}{2}$ inch above the upper insertion of the auricle. If possible, we cut at once down to the bone, and make the entire incision with one sweep of the knife, the exception to this one sweep being in cases of babies and young children with large subperiosteal abscesses, in which there may be great destruction of the

ued by means of large strong curettes and Rongeur forceps. Every portion of the mastoid process that seemed to be affected to a serious degree was removed. After removal of the extreme mastoid tip with the Rongeur, the cells as far back as the posterior border of the sigmoid groove, together with the diploic cells at the posterior root of the zygoma, were carefully removed with the curette.

The next step in the operation was the removal of the parallel with the general direction of the auricle, extending the roof and floor of the canal. This is a distinct feature of the modified operation. Under a brilliant illumination the bone is removed with the Rongeur forceps as deeply as the facial ridge, after which the electromotor burr is employed to smooth the facial spur. There is practically no danger of wounding the facial nerve by this procedure if performed carefully.

Fig. 1.—Modification of the simple mastoid operation, illustrating a radical departure from the classical operation of Schwartz. The exposure of the facial nerve in its osseous canal is incidental and not a necessary accompaniment of the operation.

Fig. 2.—The original Schwartz operation.

outer table of the mastoid. In these cases we work our way carefully through the abscess with the blade of the knife held horizontally so as to avoid injury to important structures.

To avoid mutilating the periosteum and thereby destroying its value as an agent in the repair of bone, it is now carefully pushed forward and backward with Langenbeck's periosteotome, so that the planum mastoideum is freely exposed and the hemorrhage controlled. The soft parts are now retracted with a self-retaining retractor, thus exposing the underlying bone and the operative field and at the same time the superior posterior segment of the bony meatus with the spina suprameatum and fossa. The exposed surface of bone is now carefully examined for a carious spot or sinus and for the minute dark-blue points over the cortex in the vicinity of the antrum—the so-called "shaven beard" appearance. The anatomic and surgical landmarks, as the spina suprameatum and fossa, linea temporalis, the squamomastoid suture, etc., are carefully noted. The primary opening in the bone is made with a gouge of large width, driven by blows from a suitable mallet.

Within the mastoid cavity the operation was contin-

It is presupposed that the surgeon, before attempting to perform a mastoid operation on a living person, has rehearsed it many times on the cadaver and is familiar with the anatomic and surgical landmarks of the temporal bone. It is also highly important that the surgeon should familiarize himself with the use of the chisel and gouge before attempting to perform such an operation. There is probably no other surgical feat which requires a more perfect knowledge of the technic of chiseling than does this one.

The final step in the process consists in packing the posterior membranocartilaginous canal wall firmly backward into the osseous cavity, in close contact with its walls, where it is securely held in position by a tampon of iodoform gauze inserted into the auditory canal. This is an extremely important step, since by this procedure we have a cavity, one-half of which is composed of soft tissues, instead of a cavity in the mastoid whose walls are entirely of bone, as is the case in the classical operation.

ADVANTAGES OF METHOD.

By this method bony cavity is practically disposed of, and hence we have union by first intention, instead of healing by granulation. When the gauze packing has

been completed, the mastoid incision is closed with sutures and the usual dressings are applied; the small gauze wick being allowed to remain in position until the soft parts are adherent to the bony walls. Adhesion is usually looked for at the first dressing subsequent to the operation.

It is evident that this process differs materially from Schwartze's original operation, first, from the fact that the posterior osseous canal wall, together with a part of the roof and floor of the bony canal, is removed, and, secondly, from the fact that the soft structures are carefully pressed back into the cavity thus made. In the simple mastoid operation, as performed to-day, the osseous cavity is allowed to fill up by granulation tissue from the bottom, or with a blood clot, according to the method of Dr. Clarence J. Blake. But in the modified operation, as before stated, the wound cavity is oblit-

erated by the soft structures, which are snugly fitted back into it in much the same manner as a hand fits into a glove, thus leaving no space in which to granulate.

The objection has been raised repeatedly that the antrum is not properly drained by this method, to which we answer that the wound is drained by inserting a small gauze tape or cigarette drain into the antrum. We drain the middle ear by a free paracentesis, beginning in the attic, so as to drain that cavity as well as the aditus ad antrum, thence sweeping through the posterior fold of the membrana tympani, and terminating on a level with the floor of the annulus tympanicus. With the exception of the tampon in the external auditory canal and gauze pad within the concha, the dressing of the wound differs in no material way from the dressing employed in the Schwartze operation. It is often advisable to remove the outer dressings on the following

Fig. 3.—The radical mastoid operation (tympanomastoid exenteration or Stacke-Schwartz operation). The mastoid cells, antrum, tympanum, attic and auditory canal are converted into one large osseous cavity with smooth and healthy walls.

Fig. 4.—The Panse flap operation, with flap of skin dissected from the posterior edge of the mastoid opening and turned into the osseous cavity, thus covering the floor of the mastoid and entire facial ridge with skin.

CASES ILLUSTRATING MODIFIED METHODS.

Age.	Patient.	Sex.	Diagnosis.	Duration of Disease, Days.	Time of Subside of Tympanic Inflammation After Operation, Days.	Time of Complete Healing of Primary Mastoid Wound, Days.	Condition of Hearing.	Remarks.
58.	F. G., female....		Otitis media suppurativa subacuta sinistra complicated with acute mastoiditis....	112	6	8	Unimpaired.	Patient refused operation for long time.
38.	J. M., female....		Otitis media suppurativa acuta sinistra complicated with acute mastoiditis....	49	9	9	Unimpaired.	Patient refused operation for some time.
28.	C. L., female....		Otitis media suppurativa acuta dextra complicated with acute mastoiditis....	6	12	10	Slightly impaired..	
28.	M. F., male....		Otitis media suppurativa acuta sinistra complicated with mastoiditis.....	60	17	17	Slightly impaired..	Patient presented himself for exam. on 8th week of disease.
12.	P. G., male....		Otitis media suppurativa acuta sinistra complicated with mastoiditis.....	6	4	10	Unimpaired.	
26.	H. S., male....		Otitis media suppurativa acuta sinistra complicated with mastoiditis.....	80	28	16	Slightly impaired..	Patient first came for treatment in 8th week of disease.
14.	D. G., male....		Otitis media suppurativa acuta sinistra complicated with mastoiditis.....	7	7	7	Unimpaired.	
40.	L. A., female....		Otitis media suppurativa acuta sinistra complicated with mastoiditis.....	25	8	8	Unimpaired.	
40.	H. S., female....		Otitis media suppurativa acuta sinistra complicated with mastoiditis.....	18	3	10	Unimpaired.	
38.	B. L., female....		Otitis media suppurativa acuta dextra complicated with mastoiditis.....	35	6	12	Unimpaired.	

day if they have become saturated and stiffened with secretions from the wound. Unless the patient has pain, with a sharp rise in temperature, or there is a foul odor about the wound, or other unfavorable manifestations, the tampon in the auditory canal and the gauze tape in the mastoid should remain undisturbed for three or four days. When the gauze tampon is removed from the external auditory canal at the first dressing, it will generally be found moistened with secretions. If the secretion is profuse, the canal should be douched with a warm corrosive sublimate solution (1-5000). If slight it may be dried out with sterile cotton on a cotton-tipped probe, after which the iodoform gauze tampon is again introduced. At the third dressing, in from 10 to 12 days, the tampon is removed from the canal and the middle ear is dusted with thymoliodid (aristol), when the patient is generally discharged cured.

The great benefit derived from this legitimate operative procedure is the unquestioned shortening of convalescence, and this advantage alone entitles it to careful consideration. As to its cosmetic effects, no deformity whatever results from it.

We report 10 cases illustrative of the practical results of the operation described. Of these cases the best results as to time of recovery, including complete healing of the wound, were obtained in 7 days in one instance. The longest period of convalescence was 17 days, demanded in only one case. There were three cases of recovery in 8 days, two in 10 days, two in 12, and one in 16.

DISCUSSION.

DR. F. B. SPRAGUE, Providence, R. I., said that Dr. Plummer's operation is certainly useful if healing can be obtained in the short time Dr. Plummer says. Last year at the American Otological Society, Dr. Sprague said he reported 200 operative cases of mastoid and gave the length of time of healing of the different cases. He did not remember the exact figures, but in round numbers about one-third were healed in less than 14 days; one-third between 14 and 20, and the other third beyond that time. When he began mastoid work the method in vogue was to leave the wound open to heal by granulation. By this method many wounds would heal slowly and fill with granulation tissue which needed frequent curetting, and occasionally a secondary operation. The principal thing which induced him to close the mastoid was the large area exposed to infection. The open wound is long and gaping, according to the length of the incision, and perhaps three-eighths to one-half an inch on both sides and consequently a constant secreting surface open to infection. His first mastoid operations in private work were in 1892, and he sewed up the wound and put in drainage, using the blood clot method as described. He has also used the blood clot method Dr. Blake described. This, however, is applicable only to a limited number of cases and under certain restrictions. Some things in the use of the blood clot which must be regarded are the preparation of the patient, the technic of the operation, and the conditions found on the inner wall. If there is an inflammatory condition, or necrosis of the inner wall which shows an inflammatory condition of the outer or sinus wall or meninges, it is not safe to close the wound for primary blood clot because there is secretion which is infectious and will invade the blood clot, which will become infected and break down. On the other hand, the modified blood clot is perfectly safe, because, after the wound has been thoroughly cleaned, the cigarette drain is placed in it, and the blood clots around the wick. The wick is removed in three or four days, and if it is clean it is left out. If there is any pus it will show, and it may be necessary to treat by drainage. In these cases the blood clot must not be wet, the mastoid wound must not be irrigated, or the blood clot will be broken down and healing delayed. The use of so-called antiseptics at the operation is one of the hindrances. A solution of 1/2,000 bichloride, or strong carbolic is apt to

cause necrosis and interfere with the healing of the wound. Dr. Sprague has given up these antiseptics and uses nothing but a sterilized saline solution. He believes in aseptis to keep the organisms away. Saline solution is non-irritating and does away with the danger of causing irritation and soreness around the wound, as do most antiseptics. Another thing; after the wound has been closed there will be a discharge of blood serum from the blood clot when the modified blood clot is used and this must be wiped away with sterilized cotton. This will continue for two or three days, when the wound does not need to be touched again. In this condition the wound is perfectly open and, if any infection takes place in the tympanum, drainage through the mastoid can be taken care of easily. It is rare to have anything further to do with operative procedures. An important thing in mastoid operation is to open the tympanic membrane first, no matter how many times it has been opened previously, and slit it from bottom to top, getting thorough drainage of the tympanum.

DR. J. F. BARNHILL, Indianapolis, referred to Dr. Plummer's statement that he sometimes removes the bone very widely, in one case he even included the zygomatic cells. It is well known how extensively within the mastoid structure the infection may spread, and how extensive must be the operation in order to cure the case. Hence, in such cases as Dr. Plummer described in which there is a large postoperative wound, Dr. Barnhill said he could not understand how it is possible for Dr. Plummer, by means of gauze tampons inserted into the external meatus, to place the posterior wall of that meatus into a position so far posterior to its normal situation. The difficulty sometimes met in the postauricular flaps, when cut as in radical mastoid operation, in keeping them in place against the denuded bone until adhesion takes place, is well known. This difficulty arises because the flaps contain resilient pieces of cartilage, and on this account it is becoming more and more the practice to remove all portions of the flaps with the exception of the skin itself. Indeed it seems to Dr. Barnhill that it would be impossible (in view of the fact that Dr. Plummer's operation requires no incision of the posterior meatal wall), even when only a small opening in the bony structures of the mastoid has been made, for the operator to crowd the posterior meatal wall backward far enough to bring it into close opposition with the posterior wall of the mastoid cavity, and to hold it there until a union of the two takes place. Since Dr. Plummer reports the successful performance of this feat which seems to Dr. Barnhill to be entirely impossible, he would appreciate it greatly if Dr. Plummer would explain this step of the technic more fully.

DR. G. L. RICHARDS, Fall River, Mass., objected to Dr. Plummer's use of the word paracentesis. Incision of the drum membrane is meant. He would like to have the term incision of the drum membrane take the place of the word paracentesis.

DR. E. M. PLUMMER, Boston, said that in the 10 operations, which he had performed, no deformity of the auricle was noticed. Neither did he notice any facial paralysis. It has been only five or six months since he commenced performing the operation in this way, but so far he had not been called on to do a secondary operation. The value of this procedure in cases of acute mastoiditis is well established. In many of the chronic cases on the other hand, in which there is caries of the ossicles and attic, complicated with cholesteatomatous masses, he should prefer the tympano-mastoid exenteration or radical operation. The average time of healing of the middle ear has varied from three to four days. Unless the patient has pain or there is decided rise in temperature, the cigarette drain should remain undisturbed for three or four days. In reply to Dr. Richards, he said that by the word paracentesis he means a free incision of the membrana tympani, beginning in the attic and terminating on a level with the floor of the annulus tympanicus. The posterior membrano-cartilaginous canal wall is pushed backward into the osseous cavity, where it is held in position by a tampon of iodoform gauze inserted into the auditory canal and a gauze pad is applied to the concha and a pressure bandage applied. So far he has had no difficulty in securing adhesion between the cartilaginous canal and the osseous cavity.