readily to this drug, but to be successful it must be given in large doses. So I have found 30 grains a day necessary to produce marked effects. I have found also that it produces its effects only up to a certain point; I have yet to see a patient entirely cured. I have also found the local application of alcohol, which I have used for the last year or two, gives good results. With these applications I have accomplished better results than before in the treatment of lupus erythematosus. With regard to the treatment of syphilis by hypodermic injection of mercury, I agree with Dr. Baum, using it deeply injected, and I do not think it is applicable to all cases. I do not use it as a routine. The old methods by the mouth or inunctions—which in private practice is objected to by many—are better measures. But there are some cases which require quick treatment, and muscular injection is the best because in a few days one can get a decided effect. In many cases inunctions are not objectionable, and where the patient can be induced to follow out that treatment, it is also of value.

Original Articles.

HYPERACIDITY A CAUSE OF SKIN DISEASES.*

W. R. INGE DALTON, M.D.

NEW YORK CITY.

A year ago I wrote: "I have demonstrated to my own satisfaction that some skin affections are due to a nutritive disturbance, a defective degenerative metabolism, owing to which there ensues deficient utilization of nutritive substances conveyed to the tissues; producing a lethargic condition of the skin, an inefficient stimulation of the nerve twigs supplying the sudoriferous glands; that bacterial agencies cause septic and putrefactive changes in the alimentary canal and that inanimate toxins maintain selective affinities for the tissues, exerting their action upon the primordial protoplasmic groups of cells. I am inclined more and more to think that the bond which connects the diatheses of the dermatoses with gout, rheumatism, arthritis, diabetes, and asthma, is generated by indiscretion of diet—and *hic est muco thesis meæ*": *that the chyme passing in a hyperacid condition from the stomach, through the pylorus, into the second stomach, entails such increased labor upon the duodenum that its contents can not be rendered sufficiently alkaline for physiologic metabolism.*

Here is the starting-point—this acid dyscrasia, as I believe, the inevitable *fons et origo* of the pathologic lesions, which lead certainly to nearly all diseases of the skin, save the contagious exanthemata. The nidus is here furnished for the elaboration of the pabulum upon which the micro-organisms feed and propagate, thereby favoring organic fermentative changes throughout the ileum and large intestine, culminating in stasis of cellular action, cutaneous obstruction, and peripheral atrophy; visceral, cerebral, pulmonary, or cardiac lesions. The hyperacidity also inhibits the functional action of the glands of Lieberkühn and Brunner. The duodenum is the paramount reflex center for the entire intestinal tract; the pancreatic and biliary fluids pour into it; the par vagum connects it closely with the central nervous system; the solar plexus associates it with the sympathetic; the splanchnic, with the filaments of the spinal cord, as well as the thoracic center, and with the plexuses of Auerbach and Meissner—the one of the muscular coat, the other with the mucosa. It is here that iron is absorbed and taken up by the transport

cells, combined with albuminous bodies, to be carried into the circulation, and it is the receptacle for an immense amount of blood. These physiologic and anatomic endowments all demonstrate the supreme importance of keeping a watchful supervision, so that the functional action of the duodenum may not be impaired. It is rather coincidental that whenever there are extensive excoriations or lacerations, or burns, involving a large amount of space, there follows ulceration of this organ.

It is contended that the infection caused by bacilli goes from the tonsillar crypts directly into the circulation, but I hold to my theory as being more tenable—it is that the chyle is emulsified while undergoing pancreatic digestion in the alimentary tract, and these minutely-divided products transport the pathogenic germs immediately through the lacteals, without causing infection locally into the vena cava; the lymph of the lymphatic system, with the micro-organism enveloped in the chyle, goes directly through the thoracic duct into the left innominate vein, and from the innominate into the venous circulation, through the right heart to the lungs, setting up infection there, by retarding oxidation, or they pass on with the blood, to find access to the cutaneous system, where, if they find suitable culture-media (the nidus upon which they feed) their deleterious action commences; leading to an aberrant cellular activity, caused by changes in contiguous surrounding fluids, unless the phagocytes succeed in devouring them.

We meet with brilliant clinical results in the treatment of summer diarrhea in children—those cases where the stools emit a characteristic odor, showing albuminoid decomposition or acid fermentation, when we feed them upon a low percentage, pasteurized, highly alkaline fresh milk, *without exhibiting even one grain of medicine.*

There are other etiologic factors, such as the physiological process of reflex irritation, of dentition, hard fecal accumulations, pregnancy, epileptiform convulsions, shock, worms, phimosis, adenoids, etc., causing, temporarily, tendencies to morbid conditions of the skin, but the *causa causam*—the main etiologic element—I proclaim it boldly, is hyperacidity of the chyle in the duodenum; to this is due many ills to which the skin is heir. I have based my treatment upon this etiologic theory and have met with phenomenal confirmation of its efficacy, in my service at the Metropolitan Hospital, and in private practice, in the various forms of eczema, pemphigus, etc. I prescribe:

R. Naphthalin gr. i
Ipecac gr. i
Charcoal gr. iss
Calomel
Strychnia
Pilocarpin
Arsenious acid, aa. gr. 1/100

The naphthalin causes antisepsis and inhibits the action of micro-organisms through the ileum and large intestine, thereby arresting organic fermentative changes. The charcoal, besides being a splendid antiseptic, converts the anaerobic condition of putrefaction into compounds which are not destructive to the tissues: the mercury, whether it acts upon the liver or not, destroys the bacterial forms in the duodenum and jejunum and lowers the blood pressure, thereby accelerating capillary circulation, thus favoring digestion and obviating putrefaction; the pilocarpin and ipecac exerting, at the same time, their influences upon the sweat

*Presented to the Section on Cutaneous Medicine and Surgery, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.

glands and lymphatics; and finally the strychnin toning up the vasomotors and the whole cutaneous nervous system. I do not claim that my remedy will kill the pathogenic microbes in every instance, but it *does* limit their morbid energies, and by so doing gains time for the conservative remedial agencies of the organism to perform their normal function. Some pathogenic bacteria, which elaborate toxins, or ptomaines in septic or putrefactive conditions, are otherwise entirely innocuous. We can unquestionably lessen their virulent activities by altering the environments of the pathogenic organisms. The stomach must be in proper condition, suitable diet enjoined, otherwise, no matter how specific the remedy may be, if the sympathetic nerve-centers, governing metabolic processes, standing for assimilation, are held in check from ptomain intoxication, no good effects can follow. Obstipation of bowels must in every case be combated, and as uric acid accumulates in the blood, as proved by Horbaczewski and Weintraud—that it is formed wherever living matter exists by the dissimulation of albuminoids—and as Kossel demonstrated, passing through the intermediate stages, nuclein, xanthin to uric acid. I generally follow my tablets by giving sulphate of magnesium, carbonate of lithium, and phosphate of soda in an effervescent form, the next morning.

It is only recently that the term *eczema parasiticum* has been tentatively admitted to the nomenclature, but not yet added to the classification of dermatology—*causa latet: vis est notissima*.

101 Convent Avenue.

DISCUSSION.

DR. W. S. GOTTHEIL, New York City—I am sorry that I did not have a chance to hear the first part of Dr. Dalton's paper. There is a marked tendency now on the part of dermatologists to pay more attention than formerly to the influence of general systemic conditions on their etiology, and it is a tendency in the right direction. The paramount influence of the Vienna school led to the very general acceptance of their ideas of the exclusively local nature and origin of most dermatoses; but that influence seems to be declining, and we are turning more and more to the opinions which have always maintained their ground among the French dermatologists, and are paying more attention to the general systemic condition or diatheses which, in some instances, at least, are the basic facts of their etiologies.

DR. S. SHERWELL, Brooklyn, N. Y.—I am in perfect accord with the general sentiment of the paper, while perhaps not entirely so with the theory of the Doctor, as to the way it is involved. I do most thoroughly believe, however, that the hyperacidity of the blood, caused in my mind by defective assimilation and equally defective metabolism, is at the root, and acts as *fons et origo* of many cutaneous diseases, as notably in eczema and psoriasis, etc. I am glad that there seems to be—the Doctor's paper being one evidence—a reaction against the dogma of a certain school that everything pertaining to skin diseases should be almost entirely regarded as local manifestations and be so treated. I am glad to see that there is more common sense and a larger degree of attention paid to systematic alterations and visceral complications today in general dermatology, diagnostic as well as therapeutic.

DR. L. DUNCAN BULKLEY, New York City—I have long maintained that we are wrong in looking so wholly, as has been done in time past by many, for purely local causes for skin diseases. Perhaps some will remember that in my paper yesterday I quoted Dr. George Elliot, who, five or ten years ago, was almost a pure localist, very full of foreign views, and very strong in his advocacy of purely local causes, and last year he expressed the opinion that the causes of pustular lesions were more sympathetic than local. The Doctor is getting nearer the truth when he calls attention to hyperacidity—another name for the "gouty state," etc.—and we will have greater success if we take into consideration internal conditions. Each one of these remedies named by Dr. Dalton has certain qualities, which combined will accomplish what he claims. The prescription offered is a valuable suggestion.

DR. W. R. INGE DALTON, New York City—Each and every

article in that prescription has been thought out from what has been said by eminent dermatologists. I have been giving that remedy for the last three years, and in 162 tabulated cases of eczema, acne, pruritus and pemphigus, taking together all the cases I have had in my own practice, and in my service in the Metropolitan Hospital and Dispensary, in all these cases I have not had one single failure. Dr. Sherwell has asked me if I employ topical applications. I use them in every case, such as zinc stearates, with carbolic acid or salicylic acid or menthol, to allay itching, in eczemas, with appropriate dressings; also in all forms of dermatitis, but I do not rely upon them as formerly, only as a palliative to relieve distressing symptoms.

ACUTE SUPPURATIVE FOLLICULITIS OF THE SCALP.*

WM. S. GOTTHEIL, M.D.

NEW YORK CITY.

The inflammatory affections of the hair follicles are still, it must be admitted, an unsatisfactory chapter in our dermatologic text-books. Some hardly mention them at all; and those who do practically admit, with James Nevins Hyde, that "there exists a whole series of interesting and important inflammatory affections of the hairy parts which have not as yet been distinctly differentiated from each other, or the whole series from all others."

Periodical literature, on the other hand, contains the records of many cases of follicular inflammation of the scalp; and the great variety in their terminology is a good index of the confusion that still prevails upon the subject. Quinquaud has described a "Folliculite Destructive"; Brocq a "Folliculite et Perifolliculite Décalvante"; Besnier an "Alopécie Cicatrisante Innominée"; Lailier an "Acne Décalvante"; Leloir an "Conglomerate Pustular Perifolliculitis"; Unna an "Ulerytnema Sycosiforme"; Sack a "Dermatitis Perifollicularis Atrophicans"; Lukaszewicz a "Folliculitis Exulcerans"; and Tommasoli a "Bacillogenous Sycosis." And this by no means exhausts the list. The American cases of follicular inflammation have been recorded by Fox, Holsten, Jackson and others.

There can be no doubt that several distinct affections characterized by follicular or perifollicular inflammation of the glandular structures of the scalp, and leading, with or without suppuration, to loss of hair and atrophic changes, are included in this list. The analysis and classification of these cases is a task that must await a wider experience and more exact records. Certain prominent characteristics are, however, noticeable in almost all of them. They are chronic, with a duration extending over months and years; they affect localized or isolated areas of the hairy skin; and they are rarely frankly suppurative, the inflammatory process being subacute, and the resultant atrophy interstitial in character.

If I now venture to add another to the recorded cases it is because it presents a clinical picture of a folliculitis of extreme acuity, pyogenic in character, universal in distribution, causing complete loss of hair, and not resulting in atrophy of the pilous structures. In the literature accessible to me I find mention in but two instances of a similar affection. R. Sabouraud,¹ under the designation of "Impetigo Pustulosa Peripilaris," describes an affection which he believes to be identical with the "Impetigo Pustulosa" which Boeckhart first recorded in 1887. It occurs usually upon the head, and always in connection with the hair. It begins

*Presented to the Section on Cutaneous Medicine and Surgery, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.

1. Annales de Dermatologie et de Syphilographie for 1890.