were men and three women. In eight cases the ulcer preceded outbreaks of tabetic symptoms and in ten they followed. In three cases the ataxia and ulcer appeared at the same time, while in one the fact could not be ascertained. The longest time noted by Hinze in which the ulcer preceded the tabes was ten years. It will be seen that in the case under consideration this length has been exceeded, some twelve years having intervened between the appearance of the perforating ulcer and the first symptom of tabes.

A number of different theories have been propounded to account for the trophic changes in tabes. Charcot believes it to be due to an atrophy of the cells in the anterior horns of the cord, while Virchow simply referred it to lowered nutrition due to perverted nerve influence. Oppenheim and Siemerling demonstrated a change in the peripheral nerves, but it is apparent that such conditions may be only secondary. Volkmann pointed out the relation of traumatism and analgesia in these cases. Undoubtedly, in advanced forms, these factors play an important rôle. It does not account, however, for those cases in which the joint troubles precede by months or even years the earliest tabetic symptoms. As a matter of fact, our knowledge of the trophic centers of the nervous system is yet too slight to throw much light on the pathology of these cases.

An interesting case of perforation of the aortic valves was reported by Teissier¹⁸ in a tabetic. The case presented a close resemblance to the ordinary perforating ulcer of the foot, and was thought by him to have an identical pathology.

Terrillon¹⁹ has reported a unique case of multiple perforating ulcers of the hands in a case of tabes of syphilitic origin.

- ¹ Arch. de Neurol. vi, 18, p. 305, 1883.
- ² Tr. Clin. Soc. Lond., 1882, xvi, p. 158.
- ³ Revue de Médecine, iv, p. 473, 1884.
- ⁴ Rev. de Méd., i, v, p. 487, 1884.
- ⁵ Gaz. hebdom. de Méd. 2, 5, xxii, p. 408, 1885.
- ⁶ Bull. et Mem. Soc. Méd. d. hôp., 1885.
- ⁷ Union Méd. 3 S., xli, p. 493, 1886.
- ⁸ St. Petersb. Med. Wochenschr., 1886, N. F. III, p. 243.
- ⁹ Prag. Med. Wochenschr., 1887, xii, p. 283.
- ¹⁰ Gaz. d. Hôp., 1889, lxii, p. 265.
- ¹¹ Rev. Méd. de la Suisse, 1890, x, p. 581.
- ¹² St. Petersb. Med. Wochenschr., 1892, x, p. 269.
- ¹³ Centralb. für Chir., 1881, viii, p. 660.
- ¹⁴ Tr. Path. Soc. London, 1884, xxxvi, p. 660.
- ¹⁵ St. Petersb. Med. Wochenschr., 1886, III, p. 243.
- ¹⁶ Soc. cit.
- ¹⁷ Centralb. f. Nervenheilk. u. Psych. 1891, II, p. 97.
- ¹⁸ Teissier, Lyon Med. LV, 1887, p. 49.
- ¹⁹ Bull. et Mem. Soc. de Chir. de Paris, 1885, N. S., xi, p. 408.

APPENDICITIS.

DEDUCTIONS FROM TWO HUNDRED AND SEVEN CASES OPERATED, WITH TABULATED REPORT.

Read before the Chicago Pathological Society, Dec. 10, 1894.

BY JOHN B. MURPHY, M.D.

CHICAGO.

Professor of Surgery and Clinical Surgery, College of Physicians and Surgeons, Chicago; Professor of Surgery, Post-Graduate Medical School and Hospital; Attending Surgeon to Cook County Hospital; Attending Surgeon to Alexian Brothers' Hospital; Consulting Surgeon to Hospital for Crippled Children, etc.

In many of the recent periodicals we find these ideas expressed: "The pendulum of operative appendicitis is now swinging in the reverse direction." "The frequency of operation on the appendix is going through the same course as did removal of the uterine appendages." The frequency of operation has not been irrationally swung, nor carried to its present position so-called excessive, but has found the strongest opposition from the very onset and is only now by

the sheer force of pathologic necessities approaching its proper position in frequency of performance. The expressions quoted are used almost exclusively by men who have not yet realized the pathologic conditions nor the true importance of early operation. Their intellectual vision is still in the shadow of the "grape seed," "cherry pit," "catarrhal inflammation," and "rheumatism" of the appendix. They have not the slightest idea of the destructive effect of an infection of the appendix. They do not realize that with an infection, the appendix may become completely gangrenous, may perforate or may produce a fatal peritonitis all in a short period of time (forty-eight hours), from the initial symptom. Again, it is said by the same class of men: "I am opposed to the indiscriminate removal of ovaries and appendices when they are not seriously diseased." So is every rational operator and this idiotic expression should not be tolerated in medical societies without appropriately stamping its author.

There has been no controversy in the history of medicine in which the struggle has been so intense between the surgeon and the physician as on that of appendicitis. Both parties have been honest in their convictions and both equally positive that they are right.

We naturally seek an explanation for this diversity of opinion as well as action, and the explanation is not difficult to discern. Differences of opinion as to the clinical course and probable results with operative and non-operative treatment have been almost entirely due to erroneous notions of the pathologic conditions. Up to five years ago we had on the pathology of appendicitis practically no other teaching but that of the post-mortem table; that has to-day been almost entirely superseded by the teaching of the operating table. To-day we observe on the operating table pathologic changes almost from the moment of their beginning until the destruction of the parts concerned is complete; whereas formerly we made deductions, and these very erroneous ones, as to conditions that existed before the pathologic changes ensued which produced death.

From results found post-mortem we inferred the cause and course, and here was the source of error; to-day we observe the disease in the various stages of its progress from its inception to the final result. We are therefore preëminently qualified to arrive at a correct conclusion, inasmuch as our opportunities are so much greater for observing pathologic changes as they progress.

Now, on what points have the surgeon and the physician differed and what has led to the diversity of opinion?

From 1886 to 1888, when Gaston and Kraft wrote their excellent monographs on appendicitis, it was the prevalent opinion of the profession all over the world that the vast majority of diseases in the right sacro-iliac region were due to cecitis, an inflammation of the cecum or its surrounding tissues as a result of disease of that viscus, and it was variously known as typhlitis, perityphlitis and cecitis. Then, after these elucidations, the physicians gradually yielded point after point until finally, to-day, it is agreed that all except 2 per cent. of the inflammatory pathologic lesions about the caput coli are due to primary lesions of the appendix.

We may now regard the first point as practically settled; the next argument advanced on theoretical

considerations was that the appendix in inflammatory infections becomes swollen, and relieves itself by the discharge of its inflammatory products through its normal opening into the intestine, thus effecting a cure.

What have we learned from observations at the operating table? That in inflammatory lesions of the appendix, the canal is rarely relieved of its inflammatory products in the way suggested, but is relieved by a necrosis of its wall and perforation. This is followed by the formation of an abscess and a subsequent necrosis of the attached wall of the viscus to which it is adherent, the contents of the abscess escaping into the intestine.

The next contest was over the presence or absence of pus outside of the appendix, in the majority of acute cases. Observation in operations has shown us that in the acute stage, in 94 per cent. of the cases, pus is found outside of the appendix; that the existence of pus outside of the appendix does not necessitate a perforation of that organ for its production. That an infection of the peritoneum and the formation of a peri-appendicial abscess may occur with an internal ulceration of the appendix and no perforation, as I have shown by cases reported in the JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION,¹ March 3, 24, 1894.

While the profession was not prepared to believe that pus was present in the form of a circumscribed peri-appendicial abscess in practically all of the cases of acute appendicitis, it had been acknowledged for many years that in some cases abscesses occurred around the caput coli, which emptied into the intestinal tract and the patient recovered. We know now that this is the course in the very great majority (about 82 per cent.) of cases that recover without operation.

The last and final questions, and the ones on which the controversy is still spirited, are: *a*, the pathologic changes produced by perforation of the appendix into the peritoneal cavity; *b*, the likelihood of that perforation, and also perforation of the circumscribed abscesses about it, into the peritoneal cavity; *c*, the immediate and remote symptomatic and physical manifestations of the latter; *d*, the ultimate result of such rupture; and *e*, the danger of infection of the peritoneum from an endo-appendicitis without perforation.

The pathologic changes produced depend: 1, on the character of the material admitted into the peritoneal cavity, *i. e.*, whether the contents of the appendix or abscess consist of, *a*, pus or *débris* in which the staphylococcus predominates; *b*, whether the streptococcus predominates; *c*, whether the bacillus coli communis exists in its virulent or inert condition; and *d*, whether the pus, in which any or all of these may have existed, has become innocuous; 2, the quantity of material admitted at one time into the peritoneal cavity; 3, on the condition of the peritoneum at the time of admission of pus.

I have observed in patients, and demonstrated experimentally, that pus in which the staphylococcus and the inert bacillus coli communis predominated, could be free in the peritoneal cavity for a long

period of time and produce but very slight changes in the peritoneum, *i. e.*, that the intestine may be bathed in pus of this character for days without excoriating the peritoneum. I have observed a pint of pus of this character in the peritoneum of a patient where the history showed conclusively that it must have been present for five days, and upon opening the peritoneal cavity the normal gloss of the peritoneum of the bowel was still maintained. I have taken this pus and produced cultures with it which demonstrated its character, and injected one drachm of the pus as removed from the peritoneal cavity into the peritoneal cavity of animals, both with and without mechanical abrasion of the peritoneum, without producing any symptoms, and no pathologic lesions were found on making an autopsy.

In cases of general suppurative peritonitis, as a result of appendicitis, that have been under my observation, in which the normal glistening appearance of the peritoneum was present at the time of operation, the patients recovered, and inoculations of the pus into the peritoneal cavity of animals produced no symptoms or lesions. When I find that the pus has not abraded the peritoneum, even though the surface has been covered by it for a length of time, I consider the case a favorable one.

Note how this contrasts with the appearance in the abdomen where the streptococcus predominates. Here we find, within a short time after the rupture into the peritoneal cavity, the peritoneum in contact with the infectious material is entirely denuded of its endothelium, resembling the condition produced by a cantharides blister. The quantity of pus is always small, rarely exceeding a few drachms, with here and there flakes of lymph. The bowel is distended and paralyzed; peristalsis has ceased. Inoculations with this pus produced the same condition; the red and angry appearance of the intestine I now recognize as a signal of the gravest importance, as the majority of my cases in which this condition was present terminated fatally.

The septic conditions of the bacillus coli communis produced about the same physical manifestations as that of the streptococcus. The quantity of material discharged into the peritoneal cavity is of great significance; if it be small the intestines and omentum rapidly circumscribe it and protect the general peritoneal cavity from invasion. If it be large, and if it be irritating and poisonous, it may abrade the peritoneum of its endothelium as boiling water would, quickly, and be followed by a rapid absorption, causing death in a few hours. I have had only one case of this class in which an abscess of considerable size ruptured into the peritoneal cavity; within two hours the extreme symptoms of shock were present and in twelve hours the patient succumbed. If a large quantity of pus of a low septic character be admitted into the peritoneal cavity, it may remain in that cavity for days and produce but slight changes; finally, from some cause, this material which had been innocuous, suddenly produces grave conditions and symptoms.

It is a matter of deep-rooted belief that a peritoneum that has been irritated repeatedly by the presence of a tumor, by the close proximity of an accumulation of infectious material finally attains an increased resistance against infection, *i. e.*, the peritoneum of a patient who has suffered from the presence of a large and movable tumor will tolerate much

¹ Dr. Fowler also made the same observation and in the *Annals of Surgery*, January, 1894, expresses himself as follows: "From the first appearance of an endo-appendicitis, channels of infection are open which may lead to the occurrence of any one or all of the forms of septic peritonitis without the occurrence of perforation, formation of adhesions, or rupture of a sero-purulent collection of an abscess cavity."

more manipulation than a perfectly normal peritoneum, and that the peritoneum in the neighborhood of an old circumscribed abscess is more difficult to infect than a healthy peritoneum.

There has been a great difference of opinion as to the symptoms of appendicitis and the relation of the symptoms to the pathologic condition within. This diversity will continue until our power of differential diagnosis is very much increased. From a train of symptoms and signs in the early stage it is impossible to say just what is the exact pathologic condition. We can say, however, that it is a disease of the appendix in some stage of its development.

The symptoms of appendicitis are: 1, a sudden pain in the abdomen; 2, shortly followed by nausea and perhaps vomiting; 3, local tenderness over the site of the appendix, and most frequently in the right iliac region; 4, elevation of temperature.

These symptoms occurring in this order without a previous history of genito-urinary infection, lesions of the gall tracts or Pott's disease, indicate appendicitis with almost uniform regularity. They do not indicate that the appendix is gangrenous or that it is a simple catarrhal appendicitis. They do not indicate whether the appendix has perforated or has not perforated; nor whether the cause is an infection with the staphylococcus, streptococcus, bacillus coli communis or the presence of a fecal stone or a foreign body. They do not indicate whether it is stenosis of the appendix or an appendicitis obliterans but merely that it is an appendicitis, a disease of the appendix.

The physical manifestations, the presence or absence of induration, tumefaction, edema of the wall, tympanites, etc., aid in the differentiation of the various pathologic conditions present in the more advanced stage, but all of these are absent or comparatively worthless in the early stage.

The etiology, which was for a long time a subject of contention, is now practically agreed upon. It was very difficult to eradicate from the professional mind the deep-rooted erroneous belief that in a great majority of cases, appendicitis was due to foreign bodies admitted into and retained in the appendix, as grape stones, cherry stones, fragments of bone. We had four cases of foreign body and 38 per cent. of fecal stone. To indicate the causes of appendicitis, I will use the same classification employed in my article of March 3, 1893:

1. Simple pus infection, producing the catarrhal variety.

2. Extensive infection by bacillus coli communis or pyogenic microbes, producing gangrene of a greater or less portion of the appendix.

3. Pressure atrophy with infection of the appendix: *a*, by fecal concretions; and *b*, by foreign bodies.

4. Retention accumulations: *a*, from cicatricial contractions, stenosis and obliteration; and *b*, from occlusion by enterolith or foreign body.

From the reports of autopsies collected, I find that in 70 per cent. of the cases there was a perforation of the appendix. Of my own cases in which the appendix was removed there was about 80 per cent. perforated. Simple catarrhal primary appendicitis, if it exists, is rarely brought to the attention of the physician and still less frequently to that of the surgeon. Of 207 cases reported we had but one of this variety. A catarrhal inflammation where the more serious form of appendicitis had previously existed is not uncommon.

The question of greatest practical importance to the surgeon and physician is the diagnosis; can we determine when an appendicitis is present, and can we determine the pathologic conditions that exist in the abdomen at any given time in the progress of the disease? To the former we must with positiveness answer in the affirmative; to the latter an equally forcible negative must be given, that is, we can say from the symptoms that a lesion of the appendix exists but we can not say, in most cases, how extensive, how dangerous and how far-reaching the effects of that appendicitis may be, and still less can we say in any given case how soon the life of the patient may be greatly jeopardized by the disease which at the moment has apparently no grave symptoms.

If we were able to determine in the early stage from the symptoms, physical signs or clinical history of a given case that it was going to take a favorable or unfavorable course, we could regulate our treatment accordingly. Unfortunately, there are no symptoms present in the dangerous cases, premonitory of their fatal termination, until the patient has passed beyond the stage where an operation offers reasonable hope for a recovery.

The question to be answered is, When should we operate?

The so-called conservative answers: "Operate on the collapsed cases; the cases that present symptoms of fulminating peritonitis; the cases that are taking an unfavorable course; the cases that appear to be progressing rapidly to destruction under medical treatment."

The surgeon must take a positive stand and answer, No, to each of these rules. It is admitted that 50 per cent. of the fatal cases terminate before the end of the sixth day, many on the fourth and a smaller number on the second.

The surgeons who wait till the sixth day must therefore expect to have a mortality of 50 per cent. of all the cases that would die without operation. I believe the proper position for the true, earnest and advanced medical man is expressed by the eminently scientific and truly conservative physician, Dr. Norman Bridge, in the following words: "Appendicitis is one of the most frequent and dangerous of the inflammatory diseases in or about the peritoneal cavity. We do not know the line between proper medical treatment and the demand for surgical interference, and rather despair of finding it. The medical man is practically powerless to control the destiny of the patient. It is always a surgical disease and the mortality should be materially lessened by skillful surgical treatment."² Would you delay operation in a case that is progressing favorably? By that I mean a case in which the temperature is not above 99 degrees, the pulse not above 80, the expression good, the abdomen presenting no alarming symptoms. No; I have seen cases of this class go to the third, fourth and fifth day with all of the most favorable symptoms that could well be imagined and on the sixth day die. From what? From a suppurative peritonitis that had existed all that time, as shown by the pathologic changes found in the peritoneum.

We have no sign, symptom or combination of signs and symptoms which indicate with any degree of certainty, suppurative peritonitis. It may exist and the free

² Prof. Wm. E. Quine, whose name is synonymous with rational conservatism has said: "I am convinced that early operation is the most scientific treatment for appendicitis with our present knowledge of the pathologic condition."

peritoneal cavity contain drams of pus with the patient about town as the following case illustrates. The case occurred in the practice of Dr. Jas. G. Berry:

Patient, G. F., aged 19, weighed 220 pounds. On the morning of September 8, after breakfast, was attacked with a severe pain in abdomen, shortly followed by vomiting, and the pain and discomfort were sufficient throughout the day to prevent him from attending to his work. He remained about the same during the night, and the following morning, at 10 o'clock, twenty-six hours after the onset of pain, he walked a couple of blocks to the drug store and consulted the druggist concerning his trouble, who advised him to see a doctor, and he was seen by Dr. Berry at 3 o'clock; diagnosis of appendicitis made; his pulse when seen by Dr. Berry was 140 and temperature 102 degrees; no tympanites, increased local tenderness over the region of the appendix, peristalsis absent, patient's countenance good and expressed himself as feeling tolerably comfortable. At 4 o'clock, assisted by Drs. Berry and McQuaig, I did a laparotomy. On opening the peritoneum, pus flowed out; there were no adhesions whatever between the intestines; the appendix was free; a fecal stone protruded from an opening in the side of appendix, where it had produced a pressure atrophy perforation; appendix amputated; abdominal cavity sponged; pus found down in the pelvis; gauze drained. The normal gloss of the intestine was not disturbed in the least by presence of the pus and the surface was not even red. Patient made an uninterrupted recovery. It will be noticed that the operation was performed thirty-three hours after the onset of symptoms and but a few hours after the patient had been walking about in the street with a suppurative peritonitis. What would have been his condition if the operation had been delayed twenty-four or forty-eight hours? The answer is self-evident.

A retention appendicitis, on the other hand, may produce a temperature of 105 degrees or even greater, enormous tympanites and pulse 140, an anxious facial expression and all of the so-called classical symptoms of a peritonitis, and the peritoneal cavity be free from infection. (See report of case which occurred in practice of Dr. E. W. Lee, JOURNAL OF AMERICAN MEDICAL ASSOCIATION, March 3, 1894. These are extreme pictures, but they do occur.

When should we operate? As we are unable from the signs and symptoms to determine the exact pathologic condition, are we justified in allowing the probability of fatal conditions to continue for such a period of time without such action as would place the patient beyond the possibility of rescue even by operative procedure? No; until such time as the physician is able to determine the exact pathologic condition and danger in the individual case, and indeed we must now consider that time far distant, he is not justified at the peril of the patient's life in restraining the surgeon from acting.

It must be conceded by all, that an operative procedure, in competent hands, is in itself one involving very little risk, while the continuation of the pathologic processes in many cases greatly jeopardizes the life of the patient.

There is only one safe position to take, and that is in every case in which a diagnosis of appendicitis is made, the operation should be immediately performed. It is true that many cases can recover without operation, but we can not differentiate, in the early stages which cases are going to be favorable ones. It is further true that the earlier the operation is performed, the less the danger to the patient and the greater the ease of removing the appendix.

In a majority of cases, if an operation be performed within forty-eight hours after the onset of the symptoms, the appendix is not yet ruptured and can be removed without pus infection of the peritoneum, an advantage of which even the boldest surgeon is pleased to avail himself. The following case in the

practice of Dr. Hoelscher is a beautiful illustration of this class:

Date of operation, Nov. 10, 1894. Operator, Dr. J. B. Murphy with Dr. J. H. Hoelscher. Diagnosis, appendicitis. Patient, Miss T., aged 16, was perfectly well up to 4:30 p.m., November 9, when she was attacked with pain in the abdomen, which persisted and increased in severity, compelling her to go to bed. She began to vomit at 8:30 p.m., and vomiting and pain continued through the night and up to the time of operation. At 11:30 of that night, when seen by Dr. Hoelscher, temperature was normal; the diagnosis of appendicitis was made. At 8 a.m., November 10, temperature 99.5 degrees and pulse 90; at 11 a.m., time of operation, temperature 100 degrees, pulse 110. Very tender over region of appendix; not sensitive in any other portion of the abdomen; no tympanites; peristalsis absent in close proximity to the appendix; expression good. Dr. Hoelscher is to be congratulated on his early diagnosis.

Operation: Lateral incision. Appendix covered entirely by peritoneum, adherent to the posterior surface of cecum up on to the ascending colon. End enlarged to half an inch in diameter; contained no fecal stones, but was full of pus; gangrene of mucous membrane beginning: lymph flakes in outer wall of appendix. Appendix ligated and removed without rupture; peritoneum sewed over; gauze drainage. Patient made an uneventful recovery.

In the later stage, if the surgeon is unfortunate enough to be called at the time that the peritoneum is flooded with pus and the patient collapsed, it is his duty perform the operation and give the patient the advantage of the small percentage in his favor in this forlorn condition.

The rule first, last and always should be: Operate in every case of appendicitis, promising or unpromising, at the earliest possible moment. What operation shall we perform? The incision should be made a little toward the median line from the highest point of induration, parallel to the rectus; the muscular layer should be separated with the handle of the scalpel parallel to the course of the fibers, as advised by McArthur and McBurney, and thus lessen the liability of hernia. The opening must be ample to allow the surgeon to perform the work within with ease.

In the early cases the appendix should be removed, as the adhesions are friable and permit of the appendix being readily separated and drawn into the field. The parallel band of fibers on the cecum, opposite its mesentery, can always be followed to the position of the appendix and is an infallible guide to its location.

If the non-infected portion of the peritoneal cavity be opened before the abscess is reached, *as is always desirable in the early stage*, a thorough packing of gauze should be made around the induration to prevent the gas from entering the unaffected portion of the peritoneum. This I consider of very great importance, and I make an effort to enter the peritoneal cavity just to the side of the adhesions, so as to open the unaffected portion of the peritoneal cavity before I open the abscess; in that way the adhesions may be located and the packing well placed, so as to thoroughly protect the peritoneum before the adhesions are torn.

In the later cases, that is, from the seventh day on, I consider it advisable to open directly into the abscess where adhesions to abdominal wall are present, drain the abscess, remove the fecal stone, if one be present, and make no effort to remove the appendix. The appendix in the acute cases should be simply ligated with chromicized catgut; no effort should be made to top-sew it or bury it in the cecum. Occasionally there will be a fecal fistula, but that closes in a very few days itself.

In the cases of appendicitis operated on in the intermediate stage *i.e.*, between the attacks, a cuff should be made of the peritoneum of the appendix, peeling the peritoneum back one-half inch or more; the mucous and muscular layers ligated and the cuff drawn back over the stump of mucous membrane and ligated. The pus should be carefully sponged out of the cavity and no water or disinfecting solutions used; do not waterlog the peritoneum as it is sufficiently taxed in taking care of the pus without this additional labor. A glass and gauze drain should both be used in all cases in which pus is present at the time of operation or in which there has been a recent attack of appendicitis. The drain can be removed in eight or ten days as the cavity fills up from the bottom.

This, in short, is an outline of the procedure I follow. All of the minor details can not be mentioned here. There is no operation in the peritoneal cavity where the patient's life is so completely in the hands of the operator, where the errors of doing too much or too little work would be so likely to result fatally as in the operation for appendicitis, with its multi-form and complicated pathologic conditions. The results depend in a very great degree on the extent of the invasion, and the latter upon the period of time that has elapsed since the onset. In speaking of results, I do not mean the results in recurring appendicitis, the simplest and least dangerous variety of all; I mean the results in all classes of cases, from the general suppurative peritonitis as a sequence of appendicitis, to the simplest catarrhal variety.

There is a growing tendency in the profession to shirk the responsibility of operating where the dangers are great, *i.e.*, to defer operation on the most dangerous cases and advocate operation on the obliterated or stenosing recurrent varieties in which there is the least danger to the life of the patient from the disease. We must not swerve from our obligations to our patients, but make every effort to rescue them, regardless of the praise or condemnation bestowed upon us.

In reporting the cases, it is the duty of every operator to classify or mention in detail the important or common symptoms of each individual case in a tabulated form and give an abstract of the history and pathologic conditions, that the analytical mind in reviewing the results obtained by the individual operators may be able to explain the cause of their success or failure when the operator himself is not. Therefore it is the duty of every surgeon, in writing on this subject, to give sufficient details of his cases to enable the reader to judge why they were of the grave or mild variety; whether they were all cases of recurrent, simple, adhesive appendicitis, or whether many of them were of the gangrenous, perforative and malignantly infectious variety with the peritoneum extensively involved.

(To be continued.)

A MEDICINE MAN SLAIN.—A medicine man of the tribe of Puyallup Indians, named Jim Bonchette, was killed by Jerry Dominic, another Indian belonging to the Muckleshoot Reservation, in the State of Washington, for the reason that the former had failed to cure the sickness of three of the children of the latter.

ORIGINAL INVESTIGATIONS ON THE NATURAL HISTORY, (SYMPTOMS AND PATHOLOGY) OF YELLOW FEVER. 1854-1894.

BY JOSEPH JONES, M.D., LL.D.

NEW ORLEANS, LA.

(Continued from page 404.)

CHAPTER VII.

MICROSCOPIC AND CHEMIC EXAMINATION OF BLOOD.

The color of the venous blood was purplish, between that of arterial and venous blood. When a drop of blood was allowed to fall upon a sheet of white bibulous paper, a central bright red spot remained with a surrounding bright golden aureole of serum, which spread around the central corpuscles by capillary action. The blood coagulated very slowly and formed a large loose coagulum, or clot, which contracted slowly and imperfectly. Thus in the 1,000 grain specific gravity bottle the coagulum filled the whole bottle, and from this amount of blood not more than 150 grains of golden-colored serum could be collected at the end of forty-eight hours. The blood corpuscles tended to rapid dissolution in the serum, and upon long standing the serum changed from this cause to a bright red. The reaction of the blood was most carefully determined as it flowed from the veins, and was found to be slightly alkaline, as is usual in healthy blood. I regarded this observation with interest as in several cases, in which I had abstracted blood from the cavities of the heart, after death, it gave a decided acid reaction; but the present observation would seem to show, that the acid reaction was due to post-mortem changes. Immediately after its abstraction, the blood was subjected to a careful and rigid microscopic examination. Under a magnifying power of one-fifth of an inch (Smith & Bucks, of London, best objective glass), many of the blood corpuscles presented an irregular stellate outline. When viewed under high magnifying powers, as the one-eighteenth inch immersion lens of G. & S. Merz, of Germany, with eye glasses, to magnify 1,050 diameters, the crenated and stellate blood corpuscles were found to be studded upon the surface, with nodular rounded projections. The colored blood corpuscles appeared to be undergoing changes of form, as if nodular transudations of the globulin were forming upon the surface. These changes were most marked and frequent, upon the surface and outer portions of the clots, and resembled in some respects the amoeboid movements of the coalescing corpuscles; the nodules were, however, uniformly diffused over the surface of the corpuscles. When the blood was examined, from the interior of the clot, the corpuscles were found conglomerated together forming rolls or piles, adhering together by their flat surfaces, like the rouleaux of the blood of inflammatory diseases and of the horse. The corpuscles which had been joined and agglutinated together by their flat surfaces, were normal in shape, and presented no stellate or nodulated outline as was the case with the corpuscles from the surface of the clot, and from the surrounding golden serum. It appeared as if the nodulated exudation had formed the bond of cement between the opposing flat surfaces. Upon standing twenty-four hours and longer, the colored corpuscles tended to dissolve and lose their outline, and the serum became colored from the escape of the coloring matter of the red globules.