On June 16, 1905, the following was received: "Your patient is fine. Her eyes have lost their prominence so much that it is hardly noticeable. She weighs 141 pounds" (weight at time of operation, 118). Her other symptoms have been so relieved that she had not called on the doctor for attention for over three months.

CONCLUSIONS.

1. Every irregular tumor of the thyroid, no matter how small, should be removed, as only by following this line of procedure can we hope to save the lives of those having malignant growths.
2. Every regular tumor of the thyroid causing symptoms other than deformity should be resected when, after a few months' medical treatment, its growth is not checked, and no improvement is shown.
3. Avoidance of trauma, care of the stump, free drainage and salt solution to keep full blood and lymph vessels, to prevent as far as possible wound absorption.

MUCOCELE OF THE APPENDIX.

WITH REPORT OF A CASE POSSIBLY CARCINOMATOUS
IN NATURE.*

ALFRED STENGEL, M.D.

Professor of Clinical Medicine at the University of Pennsylvania.
PHILADELPHIA.

The somewhat indefinite term mucocoele may be retained because it sufficiently suggests the general characters of a condition, the exact nature of which is probably not always the same. In some cases the pathology seems clear; in others there is much uncertainty as to the process of formation; but in all there is a cyst-like appendix with mucoid contents. Such cysts, and in fact all kinds of cysts, of the appendix are comparatively rare. According to Kelly and Hurdon, Ribbert found six cases among 400 autopsies; Bryant, one case among 124 autopsies; Steiner, three among 2,286; Bordy, one among 528, and they themselves found sixteen cases recorded among 3,770 autopsies at the Boston City Hospital. I have seen but one well-defined case, that herewith reported, among about 2,000 autopsies.

The appearance of the diseased appendix is that of a rounded or pyriform cyst with a more or less definite constriction of the proximal portion of the organ. Sometimes the distal part or tip alone is dilated; in other cases the whole appendix is enlarged. The lumen may be completely occluded at the cecal end or merely narrowed. In some of the cases the obstruction was due to a kink produced by a bend or by a fibrous band. In one, at least (Treves and Swallow), the cyst contents flowed out when the appendix was straightened.

The contents of the cysts are either of a muco-purulent, sero-mucous, serous or purely mucoid character. Occasionally they are blood-tinged. A review of the reported cases will show that in all probability the cysts are of different pathologic character. Some are undoubtedly merely inflammatory retention cysts, caused by occlusion of the mouth or proximal part of the appendix. A number of such have come under my observation, all being of moderate size and quite evidently of inflammatory origin. Exceptional cases, however, may be of such size and may have lost the characters of inflammatory retention cysts to such an extent that they may be classified as cysts and may admit of much doubt as to the mode of formation.

LITERATURE.

Leube,¹ in referring to these inflammatory retention cysts, states that the contents at first consist of tenacious mucous, and that later the sac contains only a watery serum, because when the appendix becomes much distended its wall becomes thinned and its vessels more superficial. Thus the watery portion of the blood may escape more easily, and the formation of mucus is reduced to a minimum (*hydrops processus vermiformis*).

Sonnenberg,² in referring to hydrops of the appendix and to retentive cysts, after detailing the results of obstruction of the appendix, states that the retained secretion does not always cause a severe disease of the appendix. If the obstruction occurs at a time when the lumen of the appendix is free, or when virulent micro-organisms are not present, there is no decomposition of the secretions. A continued accumulation of secretion takes place and overdistension of the appendix in the form of a retention cyst may occur. The mucous membrane becomes compressed, epithelial cells are thrown off and the connective tissue elements hypertrophy. The muscular and serous coats of the appendix also thicken. Hernious protrusions of the mucous membrane through the muscular layer are occasionally met with, as in the case described by Kelynack and Ribbert. Such protrusions are also found in cases reported by Shoemaker and Kelly and in my own.

Among others the following reports refer to cases supposed to have had this character of retention cysts:

Wenzel-Gruber³ describes a case in which the appendix was 10 cm. long, the spherical sac at the top being 4.2 cm. long, the constricted neck 1.8 cm. and the pedicle or proximal portion 4 cm. in length. The diameter of the sac was 5 cm., of the neck 2 cm. and of the pedicle 1 cm. The wall of the sac was from 2 to 3 mm. thick, as were the walls of the proximal portion. The serosa was thickened, the muscularis hypertrophied and the mucosa convoluted. The cyst was filled with viscid glassy mucus. The patient had tuberculosis of the lungs and intestines. The author notes that the opening of the appendix into the cecum, 1 mm. in diameter, was very small, but must have been thus narrowed by the chronic inflammation.

Féré⁴ presented a mucocoele of the appendix, found at autopsy on a man 55 years of age, dead of disease of the kidneys. The appendix was sigmoid and fusiform, with a circumference of 9 cm. at the central dilated portion. The length was 10.5 cm. The appendix was quite free from adhesions and normal in color. On opening the cecum there was seen a fluctuating tumor the size of a walnut, covered with reddish mucous membrane, which was almost ecchymotic in spots. The contents of this tumor were continuous with the contents of the appendix. On section of the unthickened walls of the appendix a colorless gelatinous mass, resembling boiled starch, was turned out. This was found to be composed of mucin and a few cells. The mucocoele was probably consequent on an obliteration of the appendiceal opening, due to ulceration of the mucous membrane at this point. The tumor distended the wall formed by the mucous membrane of the cecum and the appendix, and thus produced the intracecal tumor. The condition had caused no symptoms.

* From the William Pepper Laboratory of Clinical Medicine, Theobald A. Hearst foundation.

1. Ziemssen's Cyclopaedia of the Practice of Med., vol. vii, p. 361.
2. Path. u. Therap. der Perityphlitis. Leipzig, 194.
3. Arch. f. Path. Anat. u. Physiolog., 1875, vol. lxiii, p. 97.
4. Progres Med. Paris, 1877, vol. v, p. 73.

Other cases of cysts or of dropsies of the appendix are referred to by Bossard, Kraussold and Ribbert; and Wölfler describes a cyst found in an operation for herniotomy and which, on consideration of its characters, seemed to be a cyst of the appendix.

Bristowe⁵ calls attention to dilatation of the appendix which may occur when its orifice is obliterated or obstructed. "Then the appendix becomes elongated and plump (perhaps as thick as the little finger), presents often false diverticula (resembling, on a small scale, those of a sacculated bladder), and is distended with a glairy transparent fluid, the secretion of the mucous membrane."

Treves⁶ states that it is common to find the fore end of the appendix greatly enlarged and distended from the retention of mucus and other matters in the distal end of the canal.

Kelynak says that this has not been his experience. He refers to one case under his own observation and to Fenwick's case,⁷ in which "the appendix was distended by a milky fluid, the communication with the cecum being obliterated."

Maylard⁸ describes a retention cyst of the vermiform appendix from a patient who had died of chronic Bright's disease. The appendix was free, four inches in length, normal for its proximal two inches, "but the distal portion was dilated into an egg-shaped, tense, cyst-like structure." The dilated part was shut off from proximal portion which communicated freely with the cecum. A mass of clear gelatinous substance was turned out of the cyst, leaving behind some "white, creamy-like material, in which was embedded a small calcareous concretion." The peritoneal surface was deeply injected with blood vessels. Maylard notes that Coats⁹ has described a similar specimen.

Shoemaker¹⁰ describes a cystic condition of the appendix, discovered postmortem, not having given rise to symptoms during life. The cyst of the appendix was 4 inches long and $1\frac{1}{4}$ inches in its broadest diameter. "A small portion of the distal end was not dilated, and at about the broadest portion of the cyst was a thinned area seemingly about to rupture." The cyst contained a "clear jelly-like mass."

Biggs¹¹ reports two cases of stricture of the appendix. One appendix was 9 cm. long and 2 cm. in diameter, and the lumen was occluded .5 cm. from the origin by firm fibrous bands. "The contents of the appendix measured 2 drams, were of a slightly pinkish color, and consisted of fat globules, granular matter, and cholesterol crystals." The second appendix was obstructed by a kink near the tip, produced by traction of the peritoneum.

Hawkins¹² reports 2 cases of a condition which he calls "cystic appendix," which were cases of empyema of the appendix. He thinks these cysts may be caused by obstruction of the lumen due to kinking of the tube by an acute bend. He refers to a case reported by Treves and Swallow¹³ in which the cyst was 2 inches in length, very hard, and filled with mucus, which escaped when the appendix was made straight after its removal. The bent position had been maintained by old adhesions.

It is somewhat remarkable that in almost all the cases the condition was discovered accidentally at operations or autopsy or at least had occasioned no pronounced symptoms. If the accumulating liquid is of inflammatory nature, one might expect, in a greater number of the cases, active clinical symptoms, especially as the cysts often reach considerable size and not rarely have been attached by adhesions to surrounding structures. There is little doubt, however, that most of these cases were probably of inflammatory origin.

In some cases, a tumor was recognized in the right iliac fossa or the growth was of such size as to have made it probably palpable during life.

Coats¹⁴ mentions a case in which the appendix had been converted into a large cyst, measuring 5 inches in its long diameter. The cyst contained a tenacious material and the wall was thick and firm.

Wilks¹⁵ saw a case in which the appendix was dilated to the size of the ileum and distended with 3 or 4 ounces of white odorless mucus.

Wien¹⁶ reports a case of cystic degeneration of the appendix vermiformis, which caused a tumor in the right iliac fossa. There was great swelling along the crest of the ilium, and an opening near the posterior spine of the ilium, from which 3 or 4 ounces of quite clear mucus, occasionally stained with pus, were discharged daily. The discharge was said to look almost exactly like vitreous humor.

Guttman¹⁷ reports a specimen of enormous hydrops of the appendix, found on section of a woman 70 years old. The vermiform appendix was 14 cm. long and 21 cm. in circumference, and the opening into the cecum was completely constricted by strong fibrous bands produced by chronic inflammation of this part of the appendix. The secretion of the mucous membrane had become quite watery.

Montgomery¹⁸ describes a cyst of the appendix found in a woman during the operation of panhysterectomy. The cyst was $5\frac{1}{2}$ inches long and $4\frac{3}{4}$ inches in its largest circumference. The contents were "apparently thin and watery." The cyst was removed.

Deaver's¹⁹ case presented a cystic dilatation of the appendix, the size of a small orange, which was adherent both to the neighboring coils of small intestine and to the right broad ligament. The lumen of the appendix was entirely occluded $\frac{1}{4}$ of an inch from its cecal end. The contents of the cyst were quite clear and somewhat tenacious. The author says that the mucous membrane is usually smooth and generally greatly atrophied as a consequence of mechanical pressure. There is also marked atrophy of the lymphoid follicles. If the distension be but moderate, it is not uncommon to find the wall much thickened as a result of compensatory hypertrophy of the muscular coats and also of some connective-tissue hyperplasia.

Occasionally, as in a case reported by Fowler,²⁰ there are active symptoms which necessitate operation. In Fowler's case the dilated portion, about the size of a small walnut, was at the distal end. There were no adhesions.

Though accurate studies of the pathology of the appendix have until recently been wanting, and especially in the class of cases above referred to, it is very probable

5. Reynold's System of Medicine.

6. Lancet, Feb. 9, 1889, p. 268.

7. "Clinical Lectures," 1889.

8. Trans. Glasgow Path. and Clin. Soc. vol. iv, 1891-3, p. 111.

9. "Manual of Pathology," 2d ed., p. 753.

10. Occident. Med. Times, 1892, vol. vi, p. 387.

11. Med. Rec., 1893, vol. xliii, p. 536.

12. "Diseases of the Vermiform Appendix," London, 1895.

13. Lancet, Feb. 9, 1889.

14. Glasgow Med. Jour., 1875, vol. viii, p. 126.

15. Quoted by Fagge, vol. ii, p. 174.

16. Med. Rec., 1880, vol. i, p. 44.

17. Deutsch. med. Wochft., 1891, vol. xxii, p. 260.

18. THE JOURNAL A. M. A., vol. xxix, p. 174.

19. "Treatise on Appendicitis," Philadelphia, 1900.

20. "Treatise on Appendicitis," Philadelphia, 1894.

that most of the cases cited were retention cysts with inflammatory contents. Every one who has studied series of appendices must have encountered minor grades of cystic distension with contents of mucopurulent character. More advanced stages may cause empyema, and hydrops may occur in the appendix, as in the gall bladder.

There is, however, some reason to believe that there exists a quite different sort of dilatation of the appendix with which the inflammatory form may readily be confused. Long ago (1867) Rokitsky reported four cases which he regarded as instances of colloid cancer and in which the gross appearance was that of cysts of the appendix. Virchow described these cases as colloid degeneration and figures one in his "Krankhafte Geschwülzte."

In Rokitsky's series of cases²¹ the appendix was transformed into a thick-walled cyst, varying in size from 2 to 6 cm. in length and from 1 to 2 cm. in diameter. The wall was made up of fibrous tissue, which replaced the coats of the appendix. The inner portion of this capsule was a loose areolar layer, with brownish pigment here and there, and in places it was lamellated. His observations pointed to the fact that this layer might be the remains of an atrophied reticulum. The contents of these appendices consisted of a whitish-yellow gelatinous substance, supported in two cases by a fine reticular network, with small rudimentary blood vessels visible in places. After considerable discussion of the subject, Rokitsky comes to the conclusion that this affection is a colloid-cancerous degeneration of the appendix, and that the tension of the mass of gelatinous carcinomatous material causes a dilatation of the walls of the vermiform process, with a consequent transformation into a fibrous capsule.

At the time of Rokitsky's report little was known regarding the occurrence of carcinoma of the appendix, only two cases, those of Merling (1838) and of Prus (1865), having been reported. Recently, however, a considerable number of cases have been recorded, and Kelly and Hurdon tabulate a total of 52, including 2 which were designated endothelioma. They refer to Rokitsky's cases, though erroneously stating that these were not examined histologically.

In 1884, Draper²² reported a case of colloid cancer of the appendix vermiformis. "The upper third or head of the appendix was enlarged and dilated to such a degree that externally it was of the size and shape of a large plum; its cavity, deeply injected, was irregular in its surface, and would admit the little finger. The opening to the intestine from the appendix was also dilated. The thickened wall of this enlargement presented the characteristic appearances of colloid disease." The tumor had caused symptoms and was probably the cause of death.

The features of this case differed somewhat from those usually described. The thickening of the walls of the diseased appendix and the dilation of the cecal opening contrast strongly with the usual conditions.

Vimont²³ reports a case of mucocele of the appendix, found at the autopsy of a woman who had been operated on for bilateral gelatinous cysts of the ovary, which had ruptured into the peritoneal cavity. The tip of the appendix was found attached to the uterine cornus. The shape of the appendix resembled that of the stomach, with the small end attached to the cecum by a

small pedicle. The appendix was 11 cm. long and 12.5 cm. in circumference at the larger extremity. The contents were gelatinous, like boiled starch, and weighed 38 grams. The walls of the appendix were thickened, whitish, and at one place presented two small papillomatous masses. The author asks if this dilatation may not have been due to compression caused by the ovarian cysts.

This case even more definitely establishes the neoplastic nature of certain mucoceles.

Baillet²⁴ exhibited a cyst of the appendix found during an operation performed to remove a papillomatous ovarian cyst. The orifice of communication with the cecum was occluded. The cyst was found to be made up of mucoid tissue, embedded in which were many large cavities lined with non-ciliated cylindrical epithelium.

PERSONAL CASE.

The case which came under my own observation occurred in a woman aged 50, who had had glycosuria for a number of years and finally died of rapidly developed acidosis. I had not seen her during life, but made the postmortem examination, at which the enlarged appendix was discovered.

Autopsy.—Performed March 18, 1899.

The body was that of a very large white woman, probably 5 ft. 10 in. in height, and quite fat. Rigor mortis was well developed. On section into the body, the subcutaneous fat was found excessive, being in places as much as an inch and one-half in thickness. The fat itself was of a light yellow color, and the muscles were rather lighter colored than normal. The pleural cavities were free of exudation, and there were no adhesions. The anterior mediastinum was occupied with a considerable excess of fat, so that the heart and pericardium were completely covered. On section into the pericardial sac, a little clear fluid was discovered. The epicardium contained an excessive deposit of fat at the auriculo-ventricular furrow on the right side and over the root of the great vessels, but not much anywhere else. The coronary vessels were distinctly seen as white streaks, the whiteness being due to the fibrous character of the walls of the vessels. The root and larger branches were so sclerotic that when cut transversely and compressed they immediately sprang open from the stiffness of their walls. The heart muscle was flabby on both sides, and there were some fatty patches here and there. The endocardium was white in places, and there were marked patches of atheroma in the anterior mitral leaflet, as well as in the root of the aorta around the mouths of the coronary vessels and at the edges and floor of the sinuses of Valsalva. The valvular orifices were normal, as were also the leaflets of the aortic and pulmonary valves. The tricuspid leaflets showed slight thickness at the edges.

The lungs were congested at the bases and somewhat edematous, but there was no marked change.

The abdomen contained an immense quantity of fat in the mesentery and omentum. The gastro-intestinal tract presented no abnormalities, except in the region of the appendix. The latter was found greatly distended in a somewhat cystic fashion, and at one point a small hernious protrusion of the mucosa, through a defect in the serous and muscular coats, causing a clear vesicular projection of about the size of a grape seed, was discovered. The mouth of the appendix seemed to be occluded, but the bowel was kept unopened for subsequent dissection. The coils of intestines were everywhere more or less matted together by thin, veil-like adhesions, but there were no gross adhesions of any sort.

The liver was much enlarged and compressed by the ribs (*Schnurr Leber*). On section, its substance was rather flabby, and spots and streaks of light color were discovered. There was evidently some slight sclerosis and also a considerable degree of fatty change.

21. Med. Jahrb., Wien., 1867, vol. xiii, p. 179.

22. Boston Med. and Surg. Jour., 1884, vol. cx, p. 131.

23. Bull. Soc. de la Soc. Anat., Paris, 1887, vol. lxii, p. 608.

24. Bull. de la Société Anat. de Paris, 1891; 55, vol. v, lxvi, p. 67.

The pancreas was soft and presented no macroscopic change.

The spleen was enlarged, fully twice its normal size, and the capsule was rather hard. The substance was soft and rather light in color, with spots of dark, almost hemorrhagic, appearance. The splenic artery was atheromatous in a high degree, and the splenic vein greatly increased in size. The kidneys were a little enlarged. The cortex was rather wider than normal, and the substance deeply congested. There was no definite visible disease. The suprarenal capsules were normal.

The abdominal aorta and its branches, as far as they could be felt, were sclerotic, though much less so than the splenic artery.

The uterus and its appendages were normal.

Description of Cyst.—The appendiceal cyst was found to be about 2½ inches in length and a little over 1 inch in diameter at its thickest part, which was near the middle of the organ. The cyst extended from the cecum to the tip of the appendix. At the cecal attachment the lumen was occluded and the expansion to the cyst was sudden, as if a much enlarged appendix had been constricted at its origin by a tightly drawn encircling ligature. The unopened cyst was quite firm and elastic, giving the impression of a sac tensely distended with gelatinous matter (Fig. 1). The wall was white and glistening on the outside, and the small blood vessels showed plainly in the serous covering. On transverse section, the cyst was found to be of semi-solid consistence; its walls tough and fibrous, 1 to 2 mm. thick, and in most places cleanly separated from the whitish or opalescent gelatinous contents (Fig. 2). The latter were easily shelled out when the thickness and firmness of the walls prevented their collapse, leaving a firm shell.

Microscopic Examination.—Microscopic examination showed the wall or capsule of the cyst to consist of a thickened and fibrous muscularis and serosa (Fig. 3). The submucosa was of loose areolar character, containing considerable leucocytic infiltration and new-formed spindle cells. The mucosa was nowhere intact. In its place was found a layer of varying thickness, composed of polymorphous epithelial cells without definite arrangement. Generally this epithelial lining was thin and in some places wanting; in a few places it was more massive, and at such places there could be traced into the gelatinous cyst contents, irregular extension of columns or strata of epithelial cells.

Cyst Contents.—The contents of the cyst were made up of a lamellated gelatinous substance of compact character. Between the lamellæ were found cells (leucocytes) and many minute spherical bodies staining deeply with hematoxylin (calcareous spheres, Fig. 4), and here and there masses of irregular epithelial cells or less frequently ill-defined glandular acini. While the epithelial cells were more numerous near the periphery, that is, just within the capsule, they could be found at the very center and scattered everywhere through the gelatinous substance. There was no evidence of a well-developed reticulum, but in places the irregular course of the gelatinous lamellæ and the presence of short strands of spindle cells lying end to end suggested an old reticular arrangement. Nearly all the cells within the gelatinous material were degenerate, the groups of epithelial cells in particular.

The appearances strongly indicated some active cellular process with mucoid degeneration rather than an exudation from a diseased, but more or less intact, mucous membrane. The presence of acini deeply embedded in the gelatinous material and of irregular heaps of polymorphous epithelial cells further confirmed the suspicion that the process was akin to that of gelatinous carcinoma elsewhere.

There was no evidence in any of the sections of invasion of the walls of the appendix by a heterotopic epithelial process, though here and there in the submucosa small collections of epithelium-like cells could be identified. It is possible that a more extensive sectioning of the appendix might have discovered definite invasion, but at the time when the sections were made no suspicion was entertained of a possible malignant nature of the condition.

It may be recalled that in some of the reported cases, such as that of C. C. Norris,²⁵ but little invasion of the walls was

evident, the growth appearing, in the main, as a mass within the lumen of the appendix.²⁶

Chemical Examination.—Chemical examination of the contents of the cyst showed the presence of considerable quantities of calcium which the microscopic examination had indicated. The fact that the mucoid material was a mucoprotein of the same general character as those met in gelatinous tumors of the ovary and the like seems to me of some importance, though not a certain proof of the nature of the condition.

Dr. Edsall reported his chemical study of the cyst contents as follows:

"The substance from the interior of the appendix was (after preservation in alcohol) of somewhat the same consistency as a dense jelly. It was of grayish-white color, partially opaque. It was soluble in water, forming a slightly glistening solution. On boiling, the solution became opalescent, but did not coagulate. It did not coagulate on the addition of acetic acid. Absolute alcohol produced a precipitate which was still readily soluble in water after standing under alcohol for more than 24 hours. After boiling with 2 per cent. sulphuric acid, the substance reduced Fehling's solution readily. The lack of coagulation after adding acetic acid is sufficient to indicate that it was not mucin; the solubility in water distinguished it from colloid. The reactions are those characteristic of pseudomucin. After allowing the preserving alcohol to evaporate from two small portions of the substance, these portions were weighed, incinerated, and the calcium of the ash determined. In the two instances the weight of the calcium (as Ca O) was, respectively, 1.1 per cent. and 0.85 per cent. of the total."

It must be admitted that the nature of the process in the appendix I have described is somewhat uncertain. At first I was disposed to think it simply a dilated appendix filled with mucoid material unusual in amount and character. Closer study, however, and in particular the discovery of irregular acini, epithelial tubular structures and collections of epithelial cells scattered everywhere through the gelatinous substance made me suspicious that the condition might be allied to colloid carcinoma. A review of the scant literature of the subject and especially the features of the cases reported by Vimont and Baillet seem to lend some support to my suspicion.

DISCUSSION.

DR. WALTER L. BIERRING, Iowa City, Iowa, said that he had seen only one case that bore any resemblance to the interesting condition described by Dr. Stengel and this was evidently not identical. An enlarged appendix, obtained at operation, measured 3½x1½ inches, and resembled a retention cyst, the entire mass having an elliptical form. The content was composed entirely of gelatinous mucoid material; at first it was thought to be purely an evidence of a perverted secretion of the lining epithelium due to chronic inflammation, following which the excessive mucoid substance had become retained in the dilated appendix, but the fact that portions of the mucosa and the other coats of the appendix were replaced and invaded by this same material made it appear, in part at least, like a retrogressive tissue change. Dr. Bierring said that the question

26. Since the above was written, an interesting specimen has been shown me by Dr. A. J. Smith, to which I am permitted to refer by his and Dr. B. C. Hirst's courtesy. Dr. Smith writes as follows:

"Appendix was removed in Howard Hospital in June, 1905, by Dr. B. C. Hirst, from a negro woman; character of case was not appreciated prior to operation. The woman was supposed to have fibroid pelvic growths. At operation both ovaries were found much enlarged and the seat of widely infiltrating papillary adenocarcinoma, with surface papillary extensions, with both tubes and peritoneal surface of uterus similarly involved. Appendix was free, short, with smooth and pale and glistening exterior, but with distal extremity distended into a small rounded cyst-like dilatation. On section the latter was found filled with clear mucus, the walls grossly showing no thickening save at one or two points where there were tiny white subserous nodules, which may possibly be foci of starting invasion from the cancerous growth of the genitals."

The appearance of the gross specimen was strikingly similar to that of my own.

Fig. 1.—Slightly reduced view of the cecum (a) and appendix (b). The mesentery of the appendix was heavily loaded with fat (c).

naturally arises as to whether the original process in the case Dr. Stengel reported was not neoplasm, carcinomatous in character, in which the epithelial elements had become completely replaced by the mucoid material. He thought the condition certainly unique, and regarded the name proposed, mucocoele, as distinctive as any that could be suggested.

Fig. 4.—Calcarous spheres imbedded in the gelatinous substance within the appendix. Zeiss. oc. 1; obj. 1/12. Specimen stained with hematoxylin.

Fig. 3.—Section of the wall of the appendix and part of the substance within. Toward the center of the appendix is seen (a) an acinous or section of a glandular tubule; at (b) an irregular mass of epithelial cells suggesting by its position and the relations of adjacent structures that it had been separated from the mucosa. No remains of normal mucous membrane are observable.

Fig. 2.—Transverse section of the appendix showing the stiff walls and opalescent mucoid contents. Life size.

The Value of Brief, Pointed Practical Articles.—Dr. George J. Monroe said in the *Cincinnati Lancet-Clinic* about a paper which he wrote: I am not gifted with a large vocabulary of words, yet on reading the first paper I prepared I found that it took me fifty-four minutes to read it. I had not then said all I wanted to. I began to eliminate and condense. I rewrote the paper. I found that it took thirty-two minutes to read this paper. Again I rewrote the paper, and on reading it this time I found that it only required nineteen minutes. I compared this paper with the first and second, and I was surprised to find that I had really said more than I had in either of them. This will apply to the majority of papers read before medical associations. The greater number of them

could be materially condensed and yet contain all that is expressed in the long papers. How many papers we find that are filled with useless matter that has no relation to the subject. What cares the busy doctor about a lot of statistics and data (that have been published elsewhere and that can be looked up by anyone interested)? We want to learn the best way to diagnose and treat these diseased conditions.