1875	Wash. m. t.	p	Nr. comp.	wt.	s	Nr. comp.	wt.	Obs.
Mar. 25		56.4	4	3	1.122 = 11"16	3	2	N
April 6	7 ^h	56.6	5	3	1.105 10.99	4	3	H
8	7	56.2	6	3	1.142 11.36	6	3	"
14		56.1	5	2	1.107 11.01	5	2	"
21		56.6	5	2	1.101 10.95	4	2	"

In the preceding observations *p* and *s* denote the angle of position and the distance. The first numbers in the column *s* give the distance in parts of a revolution of the micrometer screw, and the second the equivalent of these numbers in arc; one revolution being equal to 9"948. This value of a revolution has been determined by Professor Holden from a large number of transits of stars and is probably very nearly correct. The numbers in the column *wt.* refer to the condition of the images, 1 denoting a very poor condition, 3 an

average and 5 an excellent condition. In the column of observer the letter N, denotes S. Newcomb; H, A. Hall, and Hn, E. S. Holden. Two achromatic eyepieces, made by E. Kahler of Washington, and giving magnifying powers of 610 and 890 have been generally used in the observations of satellites; and on a few occasions a power of 1290. In observing the companion of Sirius the power was generally 400. All the observations were made with bright wires in a dark field.

1875, Aug. 3.

A. Hall.

On the Determination of the Mass of Mars.

By A. Hall.

Since the employment of large telescopes in the search for minor planets their number has increased so rapidly that the calculation of their orbits and their perturbations has become a serious labor for the astronomers of the present time. Should this number increase for another decade as it has during the last, this calculation will become so oppressive that doubtless it will be given up: at least, the laborious method of special perturbations now in use. Perhaps some general method may be devised for tabulating the principal perturbations of Jupiter on this ring of small planets, so that without great labor ephemerides may be computed which shall be accurate enough for following and observing these planets. Without some improvement of this kind many of the planets will be lost.

In order to show the rapidly increasing number of the small planets it is sufficient to state the numbers found by three of the principal searchers; and to notice that those found by the American astronomers have all been discovered since 1861, and also, that with the extension of their elaborate star charts the yearly number of their discoveries is not diminishing.

Discoverer	Nr. discovered	Diameter of Objective	Mean Magnit.
C. H. F. Peters	22	13 inches	11.4
R. Luther	20	4.5 "	11.0
J. C. Watson	16	13	11.3

In view of the great labor attending the observa-

tions and computations of such a multitude of planets it is well to seek for any practical result which can be deduced from their theory, and to learn which of these planets may possess any special interest. That the motions of some of these small planets will give an excellent determination of the mass of Jupiter has been shown by Mr. G. W. Hill. *Memoirs of the Boston Academy*, Vol. IX, pag. 417. I have made a similar calculation in order to find the coefficients produced by the perturbations of Mars in the longitudes of the minor planets, and which have for their argument twice the longitude of the planet minus the longitude of Mars. These coefficients are those which depend on the first powers of the excentricities. The formulae necessary for this calculation are given in their general form by Laplace in his *Mecanique Celeste*, tome I, pag. 274 to 280. Since in this case we have to do with a superior planet disturbed by an inferior one, or since *a* is greater than *a'*, we have to use the formulae of pag. 274 and 275 to get the proper values of

$$a A; a^2 \frac{dA}{da}; \text{ and } a^3 \frac{d^2A}{da^2},$$

If we denote by *n* and *n'* the mean motions of the minor planet and of Mars, and put

$$\gamma = \frac{2n - n'}{n},$$

$$\gamma G^{(-1)} = J; \text{ and } \gamma F^{(-1)} = -H;$$

Prospectus.

In Kurzem erscheint in meinem Verlage:

ABHANDLUNGEN VON FRIEDRICH WILHELM BESSEL.

HERAUSGEGEBEN

VON

RUDOLF ENGELMANN,
DR. PHIL., VORMALS OBSERVATOR DER LEIPZIGER STERNWART.

In drei Bänden.

Erster Band: I. Bewegungen der Körper im Sonnensystem. II. Sphärische Astronomie.

Mit dem Bildniss BESSEL's und einer lithogr. Tafel.

gr. 4. Preis ca. 15 Mark.

Der Herausgeber beabsichtigt in dieser Ausgabe der wichtigsten Abhandlungen und Aufsätze des grossen Königsberger Astronomen den Schwierigkeiten einigermassen abzuhelpen, die sich dem Studium der Originalarbeiten Bessel's immer mehr entgegenstellen. Durch die Seltenheit und Kostbarkeit der Werke und Zeitschriften, in denen sich die meisten und werthvollsten Arbeiten zerstreut finden (vor allen der Astronomischen Nachrichten), wird selbst ihre Kenntniss immer umständlicher und die Einsicht in den organischen Zusammenhang, in die allmälige Entwicklung und Vervollkommnung in der Lösung der Probleme unbedingt erschwert. Der Herausgeber hat versucht durch Auswahl aus dem überreichen Material das unbestritten Werthvolle und Dauernde einem grösseren Kreis zugänglich zu machen und durch systematische Anordnung des Gewählten auch ein möglichst vollständiges Bild der geistigen Fortbildung Bessel's in der Behandlung der einzelnen Aufgaben zu geben. Dementsprechend enthält die Ausgabe nicht nur die als besondere Abhandlungen und Aufsätze in den verschiedenen Zeitschriften erschienenen Arbeiten, sondern auch bestimmte Theile und Capitel aus grösseren Werken (z. B. den Fundam. astron., den Tabulis Regiomont., den Werken über Gradmessung etc.), sowie selbst kürzere Bemerkungen (z. B. Auszüge aus Briefen), sobald dieselben nur ein Ganzes bilden und für das Verständniss des Früheren oder Späteren von Werth erscheinen. — Aus den die Zahl 390 fast erreichenden Druck-

sachen Bessel's wurden über 130 ausgewählt, die sich in etwa 150 Stücken auf die folgenden 8 Hauptabtheilungen vertheilen: Erster Band. I. Bewegungen der Körper im Sonnensystem (23 Stücke); II. Sphärische Astronomie (28). Zweiter Band. III. Theorie der Instrumente (22); IV. Stellarastronomie (21); V. Mathematik (19). Dritter Band. VI. Geodäsie (10); VII. Physik (12); VIII. Verschiedenes (ca. 15 Stücke).

Um dem Werke den Charakter eines möglichst einheitlichen, bis zu gewissem Grade vollständigen und auch in der Form würdigen und harmonischen zu verleihen, ist dem 1. Band neben einer kurzen Lebensskizze das Bildniss Bessel's nach dem bekannten Stich von Prof. Mandel beigegeben; und soll der 3. Band ausser dem vollständigen chronologischen Verzeichniss sämtlicher Schriften von und über Bessel das Facsimile eines Briefes von allgemeinerem Interesse bringen.

Der 1. Band wird Anfang November d. J., der 2. im Frühjahr, der 3. im Sommer 1876 erscheinen; jeder in der Stärke von etwa 45 Bogen Gross-Quart. — Mittheilungen über Fehler, besonders in den Formeln der Originalabhandlungen, deren vollständige Controle nicht möglich war und die nicht schon durch den Druck (z. B. in den Astron. Nachrichten) bekannt sind, wird der Herausgeber mit Dank annehmen und im 3. Band veröffentlichen.

Leipzig, im September 1875.

Wilhelm Engelmann.

we have by reduction of the general formulae for the values of $D^{(i)}$, $E^{(i)}$, $G^{(i)}$, and $F^{(i)}$, for the present case, in which $i = -1$,

$$J = \frac{1}{1-\gamma^2} \left[4(3-\gamma+\gamma^2) \cdot b_{1/2}^{(2)} + (3-b\gamma+\gamma^2) \cdot \alpha \frac{db_{1/2}^{(2)}}{d\alpha} - \gamma \cdot \alpha^2 \frac{d^2b_{1/2}^{(2)}}{d\alpha^2} \right],$$

$$-H = \frac{1}{1-\gamma^2} \left[\frac{3(\gamma^4+\gamma^3+3\gamma^2-5\gamma+4)}{2(\gamma-1)(\gamma-2)} \cdot \frac{1}{\alpha^2} + \frac{(-5\gamma^4+27\gamma^3-67\gamma^2+69\gamma-36)}{2(\gamma-1)(\gamma-2)} \cdot b_{1/2}^{(1)} \right. \\ \left. + \frac{(-\gamma^4+21\gamma^3-47\gamma^2+39\gamma-12)}{2(\gamma-1)(\gamma-2)} \cdot \alpha \frac{db_{1/2}^{(1)}}{d\alpha} + \gamma \cdot \alpha^2 \frac{d^2b_{1/2}^{(1)}}{d\alpha^2} \right].$$

Since we have $\alpha = \frac{a^1}{a} = \frac{1}{(2-\gamma)^{2/3}}$; J and H are functions of α , and may be tabulated with α as the argument. If now we put

$$-H \sin \varphi \cos \pi + J \sin \varphi' \cos \pi' = k \cos \beta,$$

$$H \sin \varphi \sin \pi - J \sin \varphi' \sin \pi' = k \sin \beta,$$

the inequality in longitude takes the form

$$\frac{km'}{\gamma^2} \sin (2\lambda - \lambda' + \beta).$$

We may also compute the values of k and β from the following equations;

$$-H \sin \varphi + J \sin \varphi' \cos (\pi - \pi') = k \cos (\pi + \beta),$$

$$J \sin \varphi' \sin (\pi - \pi') = k \sin (\pi + \beta).$$

I have derived the values of $b_{1/2}^{(1)}$, $\alpha \frac{db_{1/2}^{(1)}}{d\alpha}$, etc. from

Runkle's Tables of the coefficients of the Perturbative Function; Smithsonian Contributions to Knowledge, Vol. IX.

The following table gives the logarithms of J and H with the argument α .

α	log J	log H	α	log J	log H
0.615	0.86609	0.42522	0.655	0.84045	0.39281
0.616	0.86516	0.42401	0.656	0.84004	0.39227
0.617	0.86423	0.42283	0.657	0.83964	0.39174
0.618	0.86332	0.42168	0.658	0.83923	0.39121
0.619	0.86243	0.42056	0.659	0.83883	0.39068
0.620	0.86155	0.41946	0.660	0.83843	0.39016
0.621	0.86069	0.41841	0.661	0.83804	0.38965
0.622	0.85985	0.41738	0.662	0.83765	0.38915
0.623	0.85903	0.41637	0.663	0.83727	0.38865
0.624	0.85824	0.41539	0.664	0.83690	0.38815
0.625	0.85749	0.41442	0.665	0.83654	0.38766
0.626	0.85679	0.41346	0.666	0.83617	0.38718
0.627	0.85613	0.41252	0.667	0.83581	0.38671
0.628	0.85550	0.41160	0.668	0.83545	0.38624
0.629	0.85490	0.41070	0.669	0.83509	0.38578
0.630	0.85428	0.40981	0.670	0.83473	0.38532

α	log J	log H	α	log J	log H
0.631	0.85361	0.40896	0.671	0.83437	0.38486
0.632	0.85288	0.40814	0.672	0.83402	0.38440
0.633	0.85210	0.40736	0.673	0.83366	0.38394
0.634	0.85134	0.40660	0.674	0.83330	0.38347
0.635	0.85059	0.40585	0.675	0.83295	0.38301
0.636	0.84990	0.40510	0.676	0.83260	0.38255
0.637	0.84926	0.40436	0.677	0.83224	0.38210
0.638	0.84869	0.40362	0.678	0.83188	0.38164
0.639	0.84817	0.40288	0.679	0.83152	0.38118
0.640	0.84766	0.40214	0.680	0.83116	0.38073
0.641	0.84714	0.40141	0.681	0.83079	0.38027
0.642	0.84661	0.40071	0.682	0.83043	0.37981
0.643	0.84608	0.40003	0.683	0.83006	0.37934
0.644	0.84555	0.39937	0.684	0.82968	0.37887
0.645	0.84502	0.39874	0.685	0.82930	0.37840
0.646	0.84452	0.39811	0.686	0.82891	0.37793
0.647	0.84403	0.39749	0.687	0.82852	0.37745
0.648	0.84356	0.39687	0.688	0.82812	0.37697
0.649	0.84309	0.39625	0.689	0.82773	0.37649
0.650	0.84263	0.39564	0.690	0.82733	0.37600
0.651	0.84218	0.39505	0.691	0.82692	0.37550
0.652	0.84173	0.39448	0.692	0.82650	0.37499
0.653	0.84129	0.39391	0.693	0.82607	0.37447
0.654	0.84086	0.39336	0.694	0.82563	0.37395
0.655	0.84045	0.39281	0.695	0.82517	0.37343

The elements of Mars which I have used are the following:

$$m' = \frac{1}{3,000,000}$$

$$n' = 1886''6558$$

$$\log a' = 0.1828760$$

$$\varphi' = 5^\circ 21' 4''5$$

$$\pi' = 333 \ 17 \ 53.7 \text{ M. Eq. 1850.0.}$$

The following are the elements of the minor planets, the longitudes of the perihelia being referred to the epoch 1850.0:

	n	log a	φ	π
Vesta	977.6698	0.3732096	5° 4' 22".0	250° 35' 9"
Hebe	939.5953	0.3847105	11 44 17.0	14 55.6
Iris	962.5806	0.3777130	13 20 50.0	41 23.4
Flora	1086.3310	0.3426963	9 0 56.0	32 56.2
Metis	962.3390	0.3777857	7 5 2.0	70 56.8
Parthenope	923.6220	0.3896749	5 42 11.0	317 41.8
Victoria	994.8347	0.3681389	12 38 45.0	301 38.6
Thetis	912.5902	0.3931539	7 25 46.0	261 12.2
Melpomene	1020.1198	0.3609032	12 34 20.0	15 2.2
Fortuna	930.0764	0.3876587	9 10 27.0	30 38.3
Massalia	948.8831	0.3818626	8 13 2.0	98 41.7
Lutetia	933.5544	0.3865780	9 19 45.0	327 1.5
Phocaea	954.0216	0.3802989	14 47 32.0	302 23.2
Enterpe	986.6944	0.3705493	10 0 56.0	87 42.7
Urania	974.5001	0.3741498	7 16 35.0	31 21.2
Isis	930.9057	0.3874010	13 2 21.0	317 52.8
Nysa	941.1804	0.3842225	8 40 7.0	111 31.6
Nemausa	975.4748	0.3738603	3 51 18.0	174 18.0
Echo	958.2732	0.3790115	10 35 26.0	98 10.8
Ausonia	955.5447	0.3798371	7 7 1.0	269 59.7
Asia	942.2820	0.3838838	10 45 17.0	306 9.5
Feronia	1040.1468	0.3552747	6 52 46.0	307 33.1
Eurynome	928.8736	0.3880333	11 12 50.0	43 57.4
Sappho	1019.7815	0.3609997	11 32 35.0	354 53.3
Beatrix	936.6616	0.3856159	4 45 49.0	191 29.7
Clio	976.8636	0.3734484	13 39 10.0	338 55.3
Artemis	969.7655	0.3755599	10 4 28.0	242 12.6
Iphigenia	934.6788	0.3862296	7 23 9.0	337 44.2
Amalthea	968.4048	0.3759665	5 2 10.0	198 33.0
Thyra	966.6444	0.3764932	11 11 19.0	42 41.9
Peitho	933.5089	0.3865920	9 25 32.0	76 45.2
Velleda	930.9792	0.3873777	6 5 31.0	347 29.7

In most cases the preceding elements are osculating, but I hope the values of the mean motions are so nearly correct that they will give sufficiently approximate values of the coefficients, at least in all cases except when $2n - n'$ is a very small quantity.

The following table gives the value of each coefficient and its period; and it will be seen that Massalia, Echo, Beatrix and Peitho are the planets best adapted for determining the mass of Mars. It should be noticed that in opposition, when these planets are observed, the coefficients will be increased in the ratio approximately of a to $a - 1$. Thus the apparent coefficients of Massalia, Echo, Beatrix and Peitho, when in opposition are nearly, 793".3 : 121".7 : 563".3 and 230".3.

Vesta $9''.2 \sin(2\lambda - \lambda' + 7^\circ 19' 4'')$ Period = 51.7 years
Hebe 484.4 ,, (,, + 78 17.3) 475.3 ,,
Iris 30.0 ,, (,, + 77 38.5) 92.1 ,,

Flora	$0''.5 \sin(2\lambda - \lambda' + 62^\circ 55' 8'')$	Period = 12.4 years
Metis	33.7 ,, (,, + 50 33.6)	93.3 ,,
Parthenope	16.3 ,, (,, + 17 21.8)	90.0 ,,
Victoria	2.2 ,, (,, + 329 48.0)	34.4 ,,
Thetis	10.1 ,, (,, + 200 21.2)	57.7 ,,
Melpomene	0.1 ,, (,, + 82 11.3)	23.1 ,,
Fortuna	48.2 ,, (,, + 64 35.2)	133.9 ,,
Massalia	464.0 ,, (,, + 45 24.9)	319.4 ,,
Lutetia	40.4 ,, (,, + 16 25.1)	181.5 ,,
Phocaea	47.9 ,, (,, + 313 55.1)	165.9 ,,
Enterpe	8.2 ,, (,, + 52 8.9)	40.9 ,,
Urania	9.3 ,, (,, + 55 40.4)	56.9 ,,
Isis	18.2 ,, (,, + 331 8.1)	142.8 ,,
Nysa	3273.3 ,, (,, + 41 48.9)	826.1 ,,
Nemausa	12.9 ,, (,, + 22 26.8)	55.2 ,,
Echo	70.8 ,, (,, + 49 9.2)	118.7 ,,
Ausonia	62.1 ,, (,, + 358 13.5)	145.2 ,,

Asia 4533'9sin(2λ—λ'+344°21'0) Period=1696.3 years	Iphigenia 68'2sin(2λ—λ'+31° 3'3) Period=205.1 years
Feronia 0.8 „ („ + 8 6.9) 18.3 „	Amalthea 21.2 „ („ +15 44.8) 70.7 „
Euryome 49.4 „ („ + 70 5.7) 122.7 „	Thyra 19.7 „ („ +70 7.5) 76.1 „
Sappho 0.8 „ („ + 70 47.7) 23.2 „	Peitho 135.7 „ („ +54 54.8) 180.7 „
Beatrice 331.5 „ („ + 13 55.2) 266.1 „	Velleda 57.0 „ („ +33 22.0) 143.7 „
Clio 1.3 „ („ + 68 41.3) 52.9 „	U. S. Naval Observatory. 1875, July 23.
Artemis 18.4 „ („ +353 8.6) 67.1 „	

On the Washington Observations of Flora in 1873.

I was absent when Dr. Galle wrote here inquiring about some of the Washington observations of Flora in 1873. Lately I have received his memoir on the solar parallax, and have examined the differences he finds between our observations and those made at some of the other observatories. The reductions by Mr. Skinner are correct; and the only note I find is one on Oct. 25, when the preceding star is noted as too faint and the illumination was changed from dark to bright wires.

Professor Galle assumes that the larger differences are due to a want of firmness in clamping our instrument in declination. With this assumption I cannot agree. As it is important that we should know the source of the discordances that are found between the micrometrical measurements made at various observatories, and as they are interesting, tending as they do to give us some idea of the real accuracy of such measurements, and of the results deduced from them, both in the case of solar and of stellar parallax, I give below in detail the Washington observation of Oct. 16, 1873. The method of observing with this instrument followed by Mr. Ferguson was to clamp the instrument in declination and to leave it free in right ascension, except in the case of very great hour angles. In 1869 this instrument was handled by many persons and its counterpoise was disturbed so much, probably in reversing the telescope, that afterwards I found it necessary always to clamp the instrument firmly both in right ascension and declination. Before beginning the observations of Flora the instrument was examined, and had there been a looseness in the declination clamp it could hardly have escaped notice. Again, a change in the declination of the instrument would affect the observed difference of declination only when this change occurred between the bisection of the star and the bisection of the planet, and as we cannot suppose such a change to be the same in each case the instability of the instrument would be shown by reducing each comparison separately. But nothing of this kind appears.

The cause of the discordances is to be found, I think, in the micrometer itself and in the manner of using it. (1) After some preliminary trials and as the observations were purely differential I decided not to reverse the micrometer, and the head of the screw was always kept downward, in the most convenient position. This was a mistake that may have given use to small errors when the objects differed much in magnitude. (2) The wires of this micrometer are too thick for delicate work. (3) In all cases of the larger discordances at Washington the difference of declination was considerable, so that the star and the planet were observed on different wires and the distance between these wires was measured by coincidences.

On Oct. 16th Professor Galle finds the difference Lund-Washington = -1"103. On this day the observations at Washington were made with dark wires in a bright field, and the images are estimated as of average quality. I take from my observing book the observed comparisons, which are reduced to 1^h39^m sidereal time. The value of one revolution of the screw is 15"3107, and the constant 13"064 has been added to the difference of the micrometer readings.

	Sid. Time	Micr.	Red.	Δδ
Oct. 16	0 ^h 35 ^m 35 ^s	24.655	+0.835	25.490
	42 32	24.842	.743	.585
	0 49 55	24.826	.646	.472
	1 6 32	25.051	.428	.479
	12 39	25.138	.374	.512
	18 41	25.277	.268	.545
	24 55	25.273	.185	.458
	31 40	25.416	.096	.512
	37 43	25.455	+0.017	.472
	44 8	25.648	-0.068	.580
	50 26	25.620	.150	.470
	1 56 28	25.650	.230	.420
	2 2 49	25.872	.314	.558
	8 55	25.984	.394	.590
	15 19	26.014	.478	.536