Mrs. S., colored, aged 40, gave a history of suppuration of the left ear for fifteen years, with occasional attacks of acute pain in the ear and side of the head. For three weeks before coming to the clinic she had suffered continuously from a dull pain in the ear and mastoid, which sometimes extended over the side of the head and as far forward as the eye. Examination showed that the membrana tensa had disappeared. The ossicles could not be seen. A cholesteatomatous mass occupied the attic and upper portion of the cavity of the middle ear. When portions of this mass were removed the surface beneath was covered with granulations. The temperature during her attendance at the clinic varied from normal to 101 F. Deep pressure over the mastoid caused pain. In testing the comparative resonance of the two mastoids I found that when the bell of the stethoscope was placed on the tip of the mastoid with the handle of the tuning-fork over the antrum the resonance of the left side was much greater than that of the right. The time which I could hear the fork on the affected side was thirty seconds, and on the normal side was sixteen seconds.

A mastoid operation was done December 20. The usual post-auricular incision was made, and the mastoid cortex found to be of ordinary density. The air-cells were small and filled with granulations. Hemorrhage from the mastoid wound was unusually profuse. The cavity was cleared of diseased tissue, the posterior wall of the auditory canal and the outer wall of the attic chiseled away, and the granulations and cholesteatoma carefully removed from the attic. The facial nerve was found exposed just above the oval window, and sponging of the cavity caused twitching of the left eye-lid and side of the face. A slit was made throughout the entire length of the posterior part of the membranous auditory canal; and the mastoid cavity, the middle ear and the auditory canal were lightly packed, through the meatus, with iodoform gauze. The posterior incision was completely closed with sutures.

The only thing of special interest in the case is the condition of the mastoid in relation to the test with the stethoscope and tuning-fork. The fork I have used in these tests is a C 512, so constructed that it should be heard by the normal ear about thirty-five seconds. The stethoscope has a metal bell five-eighths of an inch in diameter.

100 State Street.

OCULAR MANIFESTATIONS OF DIABETES MELLITUS.*

L. A. W. ALLEMAN, A.M., M.D.
BROOKLYN.

To the practitioner of general medicine, the eye symptoms of diabetes have only a general interest, save in their diagnostic and prognostic significance. I shall, therefore, in this discussion, avoid a detailed consideration of the ocular findings, from the standpoint of one interested in diseases of the eye, and briefly examine the relation of the more important symptoms to the primary disease. It is unnecessary to caution you against that unwarranted faith in ophthalmoscopic findings which one sometimes finds among the laity. Many patients feel perfectly assured of their physical soundness when told that no evidence of disease can be found in the eye, but I am not so sure that it is unnecessary to warn you against a

too broad skepticism. It is true that no systemic disorder writes its name in large type on the background of the eye. Syphilis does not have the sole prerogative of producing iritis, nor diabetes of causing hemorrhagic retinitis, but it is true that in certain general diseases ocular symptoms are found with sufficient frequency, and a clinical picture of sufficient definiteness is often enough obtainable to make us reasonably certain, from the eye symptoms alone, of the underlying disorder.

Further, there is information to be obtained by the ophthalmoscope, of the condition of the circulation, of the changes in the vessel walls, as well as of disease of the nerve, which is in many instances of the very highest importance and which I believe is obtainable in no other way. This is a broad field, one as yet but little cultivated; we have studied the findings discovered in more advanced conditions with much care, but it is the early conditions which we should investigate. When the student with the ophthalmoscope shall work with the student of general medicine and when their tentative findings shall have been shifted by the pathologist; when such results have been patiently collected and digested, I am convinced that the opportunity for inspection of living tissues, which the eye alone affords, will take rank as one of the most important of diagnostic methods.

While ocular lesions may be present in all forms of diabetes, they are more frequently found in the chronic cases without marked general disturbances. It is, possibly, the supposed need of change in glasses which brings the majority of these cases under our observation, for refractive changes are incidental to many of the eye symptoms of diabetes. A few cases are on record in which hypermetropia has developed as a result of diabetes, and in some instances¹ it has varied with the amount of sugar in the urine. This peculiar phenomenon is due, according to Landolt, to a change in the index of refraction of the vitreous.

The development or the increase of a pre-existing myopia is a much more common occurrence. This is frequently associated with changes in the lens, but is also seen when no evidence of cataract is present. These refractive changes are often early symptoms and are important in suggesting the possibility of some serious general disease. Langford² formulates the proposition that, "The onset of myopia in patients of 50 years of age or upward without discoverable abnormality in the lenses should always excite suspicion of diabetes;" and it is well on general principles to question any too frequent necessity for change of glass.

Paralyses of the extraocular muscles occur in mild as well as severe cases of diabetes, and may be present in patients unaware of any serious constitutional disorder. These paralyses may be transient and improve and recur with the progress of the general condition. This variability is particularly observed in the lesions occurring early in the disease; the later paralyses are much more permanent. Any of the extraocular muscles may be involved. The cause of these paralytic conditions is believed to be nuclear or peripheral hemorrhages, or peripheral neuritis. Paralysis of accommodation is probably the most frequent of the ocular complications of diabetes. It was first observed by von Graefe, and as in most other instances his opinion has been confirmed by later observers. Paralysis, or more properly paresis, of accommodation, for the paralysis is often incomplete,

*Read at the Sixteenth Annual Meeting of the Fifth District Branch of the New York State Medical Association, held in Brooklyn, May 22, 1900.

¹ Landolt: *The Refraction and Accommodation of the Eye*, p. 417.

² Norris and Oliver: *System*, Vol. iv, p. 667.

is in many cases among the very earliest symptoms of diabetes.

An incomplete paralysis is often unrecognized by the patient as such. They only observe increased difficulty in reading, which is relieved by a stronger glass, and belong to the class, to which I have previously referred, who first learn of the diabetic condition when examined for glasses. The relation of diabetes to cataract is a question which has occasioned no little discussion; some authors claim that there is a true diabetic cataract while others contend that lenticular changes are no more common among diabetics than in patients of the same age suffering from some other debilitating disease. Among patients past middle life I am inclined to believe that the influence of diabetes on cataract formation is through the general nutritive disturbances and arterial degeneration by which it is accompanied; but in the young there seems to be some direct causal relation, and the unexplained development of juvenile cataract requires exclusion of diabetes, as a possible cause, by a careful physical examination.

The most interesting, if not the most frequent, ocular symptoms of diabetes are found in the retina. It has long been known that pathologic changes are frequently found in the retina of diabetics, but owing to the similarity of these lesions to those found in other diseases attended with vascular degeneration and particularly to those of renal retinitis, many observers have denied that a true diabetic type of retinal disease could be distinguished. Hirschberg,³ who has given much careful study to this subject, is convinced that there is a distinctive diabetic retinitis, presenting in most cases either a degenerative or a hemorrhagic type, and he points out the features which distinguish it from the other forms of retinal disease. He admits that his contention is not universally accepted, and explains this fact as largely due to the confusion of other observers, who have not taken into account the frequency with which renal lesions develop in diabetics. When the two diseases co-exist we may have a mixed type of retinal disease, or a diabetic retinitis may take on a renal character when the disease of the kidney is developed in the diabetic, or a patient suffering from diabetes and renal disease may not develop a diabetic but an albuminuric retinitis. Again, there are periods in the progress of the case, during which either the sugar or albumin may disappear from the urine of a patient, who, nevertheless, is suffering from both diseases. In such an event the ophthalmoscopic findings would not be sustained by a single urinary examination made at this time. Making due allowance for these facts there is no difficulty in recognizing a diabetic retinitis which can always be confirmed by physical examination. These contentions of Hirschberg seem to be well grounded and I am inclined to accept them. The typical diabetic and renal cases are surely distinct enough, but I have never chanced to see the mixed type. Retinitis is often a late symptom of diabetes and develops after the general symptoms have been long present and are well marked; but this is by no means always the case and the diagnosis of diabetes may be first made when patients are examined for glasses or consult us for some blurring of vision, flickering before the eyes or other ocular symptoms.

Retinal hemorrhages, with or without other retinal changes, are always suggestive of diabetes, particularly the small punctate hemorrhages. They are often associated with conjunctival hemorrhages; they are absorbed and recur. They are sometimes very numerous, but fre-

quently cause little disturbance of vision. Retinal changes are always present in a case of diabetes which has existed for more than ten or twelve years, and are usually considered terminal symptoms of the disease; this is not however entirely unquestioned. I believe, while always an unfavorable symptom, retinal complications do not by any means indicate an immediate termination of the disease. Hirschberg makes a distinction between the degenerative and hemorrhagic forms, and considers the cases with profuse hemorrhages far more serious. In my own experience I have known diabetics with recurrent retinal hemorrhages to live many years in comparative comfort, and I can not view the diabetic lesions with the same alarm as those of renal disease.

There is another ocular symptom which is not uncommon and which has occasioned much discussion. This is the central scotoma.

Diabetics sometimes present a clinical picture which is almost identical with that found in the toxic amblyopias. So similar are the conditions that they have been called tobacco amblyopias in diabetes, and the frequent occurrence of these symptoms was supposed to be due to the lessened resistance to the tobacco poison among diabetic subjects. As the symptoms are found in patients who use neither tobacco nor alcohol, and who are suffering from no known toxic influence, we are forced to conclude that the poison must be in some way due to the disease. Recent studies of toxic amblyopias have largely changed our ideas of the pathology of these conditions, and the peripheral neuron, which has of late been upsetting most of the established theories of the pathology of the nervous system, seems to be the offending member in this condition as well. The altered state of the blood and the profound nutritive disturbances seem to furnish us with sufficient explanation of these symptoms, which are probably similar pathologically to the other toxic amblyopias.

There are many other diseases of the eye which are caused or modified by diabetes. Iritis of a severe plastic type is met with and is always dreaded after operation on diabetic subjects.

Keratitis, which is pre-eminently an attendant of faulty nutrition, is a natural consequence of the disease, and when occurring is often a serious condition. Optic-nerve atrophy and amblyopia, without discoverable lesions, are sometimes seen. The latter symptom is probably similar to uremic amblyopia in origin, and is sometimes a forerunner of diabetic coma.

The eye symptoms are on the whole of far more diagnostic than prognostic significance. They are nearly all seen as both early and late symptoms, and are indications rather of the present nutritive condition than of the extent of the irreparable damage to the organs. The most valuable prognostic indications are to be discovered by the ophthalmoscope in the retina, and particularly in the condition of the retinal vessels.

A patient may rally from a profound toxemia, but "he is as old as his arteries."

Remote Prognosis of Serofibrinous Pleurisy.—A French author reports in the current issue of *Arch. de Méd. Navale*, all the cases of serofibrinous pleurisy observed in the navy between 1877 and 1899, a total of 352 cases: 320 patients were discharged cured, and 32 died in the course of the disease. He finds that 131 have since died, and that in 118 of these death was due to tubercular infection. The remainder, 189, are still living. He therefore concludes that in a third of all the cases serofibrinous pleurisy, apparently primary, is in reality of tubercular origin. He found the left side more frequently affected than the right.