Figs. 11 and 12 are exact drawings of a dissection of the cells and turbinates of a child at birth. In this case there are three ethmoidal turbinates, but in 14 of the 18 babies' heads which I dissected there were only two, as in most of the adults. Note the relation of the cribriform plate to the palate bone almost parallel now.

Fig. 11

Fig. 12

FIG. 11. Outer wall of nasal cavity of a baby at birth. Natural size.

FIG. 12. Dissection of cells of same from orbital side. A. Sphenoidal sinus(?). B. Antrum. C. Frontal sinus. D. Lachrymal duct.

There are four cells opening into the superior meatus, one into the highest meatus and six into the middle meatus, two of which open into the groove of the unciform. The antrum is well developed but not more so than the other ethmoidal cells which contain about three times as much air space at birth as the antrum.

A CASE OF COINCIDENT TUBAL AND INTRA-UTERINE PREGNANCY.

BY CHARLES M. GREEN, M.D., BOSTON.

(From the Gynecological Clinic of the Boston City Hospital.)

ON June 6, 1907, Dr. Ernest B. Young, first assistant visiting surgeon of the Department for Diseases of Women, reported a case of coincident extra- and intra-uterine pregnancy, and gave an interesting statement of the present knowledge of this rather rare condition, with an inclusive, recent bibliography.¹ This was the first case of this character that had occurred in the clinic. In October, 1907, a similar case came under my observation; and a brief report of the experience, and the interesting pathological findings seem worthy of record.

¹ BOSTON MED. AND SURG. JOUR., vol. clvi, No. 23.

H. E., aged thirty-three, born in Canada, married eight years, had had two children, in 1903 and 1905 with normal labors and puerperia: she had never miscarried. The menstrual history was not noteworthy, and the last period the patient said had occurred three weeks previously. The present illness began two weeks before entrance; during this time there had been sharp intermittent pain in the rectum near the anus, not aggravated by defecation, and a continuous dull ache over the entire abdomen. The woman had had one chill, and had vomited two or three times. Two weeks before there had been a similar attack, on which the patient remained in bed for two hours, and found relief with hot water bottles. Two months before entrance, and again after the last supposed menstruation three weeks before admission, the patient flowed intermittently for a week after the regular period had ceased.

On admission the pulse was 128, and the temperature 100.2° F. General physical examination showed no abnormality, except a cardiac systolic murmur. The abdomen was lax, tympanitic, and not tender, but bi-manual examination, besides the traumata of parturition, revealed a soft, cyst-like, median mass rising from the pelvis to within 6 cm. of the umbilicus. In the pelvis, to the right and posterior, was a soft, sausage-shaped, sensitive mass connected with what felt like a cyst. The uterus was not differentiated, and was thought to be concealed by a superimposed cyst. The white blood count was 13,000: the haemoglobin test showed 70%. Neither the history nor the physical examination seemed to me and to several colleagues to warrant a diagnosis of pregnancy, either tubal or uterine; and the probable diagnosis made was ovarian cyst, with twisted pedicle, with a supervening inflammatory process. In the face of this false diagnosis, however, the satisfaction remains that the treatment pursued was the most appropriate for the condition found.

On opening the abdomen the cyst-like mass was found to be the uterus, four months pregnant; and the supposed thickened and twisted pedicle was seen to be a pregnant tube surrounded with from four to six ounces of blood clot, forming a mass in which the vermiform appendix was involved. Taking especial care to disturb the uterus as little as possible, I removed the appendix and the tubal mass, and closed the incision. On the same evening the foetal membranes ruptured. Four days later a slightly macerated, four months' foetus was cast off spontaneously, and the uterus was cured. The subsequent convalescence was uneventful.

The intra-uterine foetus had apparently been dead two or three days, and if so, was alive at the time of the removal of the partly aborted tubal pregnancy. Young states² that "in a majority of the published records, the disturbance of one of the twin pregnancies usually the extra-uterine, has ended the other"; and such was the result in this case, in spite of every effort to ensure the continuance of the intra-uterine pregnancy.

PATHOLOGICAL REPORT.

BY ARTHUR B. EMMONS, 2d, M.D.,
Senior Pathological Intern.

GROSS DESCRIPTION.

The specimens are four in number: A fallopian tube, a small piece of soft, grayish, gelatinous tissue, a fetus, and the vermiform appendix.

1. The fallopian tube measures 8.5 cm. in length. It has been opened by a longitudinal incision throughout its superior surface and has been stripped back from its contents.

² Loc. cit.

A CASE OF PREGNANCY - GREEN

FIG. No. I. Fetus expelled from uterus four days after operation for extra-uterine pregnancy.

FIG. No. II. Cephalic extremity of fetus removed from Fallopian tube in which extra-uterine pregnancy had taken place.

FIG. No. III. Fallopian tube in which extra-uterine gestation took place. The tube wall has been pulled away from the products of conception, showing the point of attachment and pedunculated appearance of the latter. At A, the free extremity of the products of conception, the fetal membranes can be seen.

There is considerable difference in the size and general character of the proximal 2 cm. and the remaining distal portion of the tube. The proximal portion shows little change. It averages 8 mm. in diameter, the mucous surface is smooth and glistening and its lumen is completely shut off from that of the remainder of the tube by a sharp constriction.

The lumen of the distal 7 cm. of the tube is considerably enlarged and contains a circular, elongated, rather firm and elastic mass, which is firmly attached at the extreme proximal end of the greatly enlarged portion of the lumen of the tube, into which it hangs as a pedunculated mass. This mass measures 1 cm. in diameter at its point of attachment. Beyond this point it rapidly increases in size, measuring 3.5 cm. at its middle and distal portions. Its external surface is lightly granular, light gray to dark red, and attached to it are a few dark red, fragile tags. The distal extremity of the mass is somewhat depressed (funnel-shaped) and is lobulated.

The glistening appearance of this lobulated extremity is due to a thin, delicate membrane, which lines it and hangs down from its free edge.

On section the pedunculated mass described above shows the following picture: The proximal 6 cm. is composed of greyish, elastic tissue throughout which are small, dark red, rather soft areas. The distal 2 cm. consists of a dark red, elastic substance which is sharply demarcated from the proximal portion. The proximal portion resembles placental tissue; the distal, blood clot, which has formed about the placental tissue and which has separated the veil-like membranes from their placental attachment.

2. In the region of the distal extremity of the fallopian tube there is a small piece of grayish, gelatinous tissue, which measures 0.9 mm. x 5 mm. x 9 mm. Its surfaces are smooth and glistening except at one point where it is torn. The general shape of this specimen is hemispherical. At one point there is a slight depression in the surface. Just below and to each side of this there is a semi-opaque area 1 mm. in diameter. Its surface is semi-transparent and shining through it is a dark bluish area. This specimen is the cephalic extremity of a fetus which has developed to the point where the eyes are well differentiated.

3. The fetus was received at the pathological laboratory four days after the other specimens. It measures 18 cm. in length, is dark red, somewhat softer than normal and shows evidence of having been dead at least two or three days before being expelled from the uterus. The head, body and extremities are well formed. The sex is easily determinable as that of a female.

4. The vermiform appendix measures 5 cm. in length. Its external surface is smooth and glistening. The blood vessels are not injected. The mucous membrane is normal.

GROSS DIAGNOSES.

Extra-uterine pregnancy (tubal).
Cephalic extremity of small fetus. (Age questionable.)
Fetus 18 cm. in length (age about four months).
Normal vermiform appendix.

MISROSCOPICAL FINDINGS.

Sections from the greyish tissue within the lumen of the fallopian tube show chorionic villi, large pale staining cells with rather small vesicular nuclei (decidual cells) and an occasional multinuclear cell (giant cell of pregnancy) and polymorphonuclear leucocytes (acute inflammation).

The cephalic extremity on section shows embryonic

muscle cells and cartilage, the nuclei of which, for the most part, fail to take the stain.

The vermiform appendix shows small groups of lymphocytes and eosinophiles in the submucosa (slight chronic appendicitis).

SUMMARY AND CONCLUSIONS.

This is, unquestionably a coincidental intra- and extra-uterine pregnancy.

The interesting question arises whether we have in this case an instance of superfecundation, the fertilization of the two ova taking place at or about the same time; or of superfetation, in which several months may intervene between the two impregnations. The uterine cavity does not become completely obliterated until the reflexa and the vera fuse at the end of the third month of pregnancy; after that time superfetation is impossible, but prior to that period its theoretical possibility must be admitted (Williams). On the other hand, superfetation is not considered possible in this connection by most investigators (Young). Simpson concludes that there is no positive proof that the ectopic ovum may become fecundated after the uterine, as the difference of size is probably due to insufficient nourishment of the former. In view of the difficulty of classification in the case reported, the following discussion of the subject by Dr. Emmons and associates in the Pathological Department is of interest:

DISCUSSION.

Coincidental intra- and extra-uterine pregnancy might take place at or about the same time; the intra-uterine pregnancy might precede the extra-uterine or the extra-uterine pregnancy might precede the intra-uterine.

In coincidental extra- and intra-uterine pregnancy it is generally conceded that the coincidental intra-uterine and extra-uterine gestation occur, as a rule, at or about the same time. The difference in the size of the two foetus is due to the adverse conditions offered for growth without the uterine cavity.

The following facts might make it possible that in this case the extra-uterine pregnancy took place some time after the intra-uterine pregnancy. 1. It is not an anatomical impossibility. 2. This patient was regularly exposed to the possibility of becoming pregnant. 3. The tissues of both foetus stained well, suggesting that neither had been long dead. 4. The foetus within the uterine cavity corresponded to one at about the fourth month of normal gestation; the foetus within the fallopian tube might be easily mistaken for one at about the sixth week of normal pregnancy. 5. The placental tissues of the ectopic gestation covered a considerable area and their blood supply was good, affording sufficient nourishment to have developed a foetus within four months considerably larger than this one. 6. The placental site within the uterus, as determined by curreting, was upon the side opposite the tube that contained the ectopic gestation, making access to this tube possible.