pair points to the externus; crossed diplopia to the internus; homonymous diplopia of the second pair points to the inferior oblique; crossed diplopia to the superior rectus; homonymous diplopia of the third pair points to the superior oblique, and crossed diplopia to the inferior rectus.

The affected muscle being found it only remains to determine whether it is the muscle of the right or left eye. This last question is answered by the well-known fact that the image of the affected eye is projected in the direction of the normal action of the paralyzed muscle; and that the distance between the double images increases in this direction and diminishes in the opposite. If, for instance, we have found a paralysis of the externus and the images separate when the eyes are rotated to the left, but come together when the eyes are rotated in the opposite direction, we know at once the paralyzed muscle must be the externus of the left eye.

I have taught this method a number of years and always found the quick and easy way of making a correct diagnosis was a great revelation to the students who are accustomed to look upon the paralyzes of ocular muscles as one of the most difficult and perplexing problems. Dr. Starkey has elaborated a similar method and I am pleased to learn that he too has been successful and well satisfied. It shows that students may readily understand the clinical pictures of ocular paralyzes if they are presented to them in a comprehensive manner.

DR. W. H. WILDER, Chicago—Dr. Starkey's table will be a valuable addition to the class room. It is certainly difficult to get students to comprehend this matter of ocular paralysis and any way of simplifying it is of value. The only suggestion I would make would be that this table might be supplemented by a diagram giving the evidences of torsion or tilting of images that we see when certain straight muscles or the obliques are paralyzed. I do not agree with Dr. Hotz on this point. If you have an affection of one of the straight muscles it is easily recognized without taking the patient into the dark room, but the difficulties come with the partial paralyzes or pareses, where the degree of diplopia is not great and the difficulties are worse when a group of muscles are affected. It is not an easy matter to work these cases out to a satisfactory conclusion.

DR. M. F. WEYMANN, St. Joseph, Mo.—The subject of the diagnosis of paralysis affecting a single muscle of the eye may be reduced to three mechanical rules whose application requires little thought and analysis: 1. Diplopia appears in the direction of the action of the affected muscle. 2. The faulty image is always in advance; this is reversed only when the paretic eye is used for fixation. The first two rules determine easily any paresis in the horizontal plane. 3. Of the muscles acting in the vertical sense, the obliques produce homonymous, the recti heteronymous diplopia.

DR. EDWARD JACKSON, Denver—I think it is extremely important for practical reasons not to attempt to burden the mind with any such table, but to follow a plan similar to that indicated by the last speaker. It is well enough to have such a table for reference, but I think for practical work we have the whole theory of the subject in the rules Dr. Weymann has given. The diagnosis of partial paralyzes in a large majority of cases is not rendered simpler by such a scheme, and I find that they can not be settled in a few minutes or even in an hour by repeated tests. You must remember that no ocular movement is dependent upon any one muscle and no movement is lost by the paralysis of a single muscle, unless it be direct lateral movements of a certain nature. I think the measure having the greatest value as a final test is the accurate measurement of the fields of binocular fixation.

DR. HORACE M. STARKEY, in reply—If I understand Dr. Hotz, this paper illustrates his method of teaching to-day as well as that of ten years ago. The object of my study has been to make the teaching of this rather complex subject as simple as possible, and certainly it has been much more easily understood by physicians and students, when explained in connection with this table, than by any other means that I have been able to devise. It is not expected that the table

will be memorized, but rather to be at hand for ready reference. The reason that no supplementary table was added, as suggested by Dr. Wilder, is because it seemed to me that the simpler the table could be made the more valuable it would be.

MULES' OPERATION.*

FRANK C. TODD, M.D.

Clinical Professor of Ophthalmology and Otology, University of Minnesota; Attending Eye and Ear Surgeon to City and Asbury Hospitals.
MINNEAPOLIS, MINN.

The operation of evisceration was devised by Graefe in 1884, his object being to prevent death from meningitis following removal of suppurating globes; and by Mules of Manchester, England, at about the same time, his object being to prevent sympathetic ophthalmitis which occasionally follows enucleation. Since which time this operation has been urged by some surgeons as furnishing a more suitable stump for prosthesis.

In 1885 Mr. Mules devised and executed the operation which bears his name. This consists in ordinary evisceration, including ablation of the cornea and the insertion of an artificial vitreous.

In 1886 Mr. Adams Frost of London practiced the insertion of a glass ball immediately after enucleation; his method was as follows: After incising the conjunctiva all around the cornea the tendons of the recti were secured by sutures, then enucleation followed with the insertion of the globe, after which the sutured tendons were fastened and the conjunctival flaps brought together.

Mr. Lang of Moorfields shortly afterwards modified this operation by inserting the artificial globe into Tenon's capsule.

In 1895 Dr. Fox of Philadelphia devised the operation of inserting an artificial globe into an orbit where the eyeball had previously been enucleated. His operation consists in opening the conjunctiva horizontally, dissecting a hole in the orbital tissues into which the globe is inserted, the tissues then being sutured over the ball. Dr. Fox records a success of 85 per cent. in 48 operations. Simple evisceration in the opinion of most surgeons presents no advantages over enucleation. It does not, at any rate, after a few months, make a better stump than results from a careful enucleation. In those cases where Mules' operation can not be performed, the Frost-Lang operation may be substituted. The Fox operation is a very ingenious one and certainly causes a great improvement in the appearance of appropriate cases, but it is probable that this operation will not be very generally adopted, since the reform Snellen eye has been invented and is giving such satisfaction. Many of the members of this Section have used the reform eyes, no doubt, and will give us the benefit of their experience. There are two forms made, of which I show some samples. In both of them the back is filled in, but in one it is filled in more than in the other. (Figs. 2 and 3.) They are certainly a great improvement over the old form. The large cavity which results from enucleation is partially filled in and does away, in part at least, with the collection of mucus, besides giving far better results from a cosmetic standpoint. Recently I received a letter from Dr. de Schweinitz, in which he says that by the use of the reform eye in cases where he has recently performed enucleation according to the improved

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Ophthalmology, and approved for publication by the Executive Committee of the Section: Drs. J. A. Lippincott, Casey A. Wood and H. V. Würdemann

technic, i. e., suturing the tendons, he has had as good results as after Mules' operation, though he states that he still performs the latter operation in selected cases. We certainly have much to thank Dr. Snellen for in giving us this great improvement, and it only seems strange that some one did not devise such an artificial eye before.* There are many cases in which the Snellen eye will be of great benefit, but I do not believe that it can ever take the place of Mules' operation where the latter can be performed, because there are at least the following advantages of the latter operation over improved enucleation with the insertion of the reform eye.

FIG. 1.

FIG. 2.

FIG. 3.

Fig. 1.—The original hollow artificial eye, or eye-shell, for cases of atrophic eye ball, and after Mules' operation. Fig. 2.—A double-walled shell for cases where a smaller stump remains, as after a simple evisceration. Fig. 3.—An artificial eye-globe, where the conjunctival sac is emptied and spacious.

ENUCLEATION WITH SUTURING TENDONS AND INSERTING REFORM EYE.

MULES' OPERATION.

1. Growth of orbits uniform.

2. Mobility.

3. Orbit entirely filled in.

1. Growth of orbits not uniform.

2. Mobility less and will decrease in time as is the case following simple evisceration.

3. Orbit only partially filled in.

As yet the use of the reform eye has not been sufficient to enable us to make further comparison.

I shall mainly give you my own experience with Mules' operation and conclusions derived therefrom. If you desire a more full treatise upon this subject and also regarding other substitutes for enucleation, I refer you to the papers read before the Section of Ophthalmology of the Thirteenth International Congress of Medicine at

Mules' Glass Spheres.

Paris, and particularly to the excellent paper read at that meeting by De Schweinitz, summarizing the opinions of 67 American surgeons who had given their experience with substitutes for enucleation, 32 of whom had performed Mules' operation. It is worthy of note that those who have performed a number of these operations have had a large percentage of successes and are enthusiastic in their commendation, while many who have performed but few and had a larger percentage of

failures do not consider the operation advisable. The great discrepancy in the results of the various operators surely indicates a difference in technic and I believe that statistics gathered a few years hence will show a greater percentage of successes and a larger number of operators who will agree with those of larger experience, notably Fox, Buller and Allport, who are strongly in favor of the operation. In England we find enthusiasts in Carter, Bickerton and others, who have performed the operation for many years. Swanzy says in his last edition of "Diseases of the Eye," regarding Mules' operation:

"This proceeding—a modification of the foregoing—was also proposed by Mr. Mules for cases of threatened sympathetic ophthalmitis, and, like simple evisceration, has not yet met with universal acceptance in those cases, because many fear that it does not afford sufficient protection against sympathetic ophthalmitis. I do not participate in this feeling. In cases of staphyloma, however, and in some other conditions where the questions of sympathetic ophthalmitis in the other eye, or of a new growth in the eye to be operated on, do not enter into consideration, no proceeding is, in my opinion, more satisfactory, at least in young persons, than this beautiful one of Mules'. The prosthesis it gives is almost perfect

Author's Introducer.

"The danger that the glass sphere may get broken by a blow upon the eye has been put forward as an objection to this method. No doubt it is an accident which may occur, and would then necessitate the enucleation of the eye, but no case of the kind has as yet been recorded, although the operation has been in use for fifteen years. Silver spheres, instead of those of glass, have been sometimes employed to obviate the danger referred to.

"I can heartily recommend this procedure. I use it frequently, and I am much pleased with it, for, with a well-fitting glass eye, the cosmetic result it gives is infinitely better than that produced either by complete enucleation or by evisceration of the eyeball."

De Schweinitz speaks very highly of Mules' operation and states in conclusion that "the best cosmetic results among the substitutes for enucleation, if successful abscission be excluded, are secured by Mules' operation which is only positively contraindicated by malignant disease, sympathetic ophthalmitis, extensive laceration of the sclera, and extreme phthisis bulbi." Yet he states finally: "Mules' operation, when successful, certainly furnishes admirable results, but I feel sure that although at the present time, from the cosmetic standpoint, it seems to be one of the best of the substitutes for enucleation it is not likely to endure as an operative measure, unless the percentage of failures is greatly reduced." There can be no doubt that this percentage of failures can and will be reduced.

The number of cases upon which I have performed Mules' operation are 31; these include two cases of Dr.

* Since reading this paper I have discovered an article published in the Ophthalmic Record, May, 1901, which states that the closed artificial eye was devised by J. L. Borsch, in 1894.

Allport (whom I assisted) which I have had an opportunity to follow out, also another case of Dr. Means of Columbus, whom I assisted in his first Mules' operation in 1899. All of these cases are either under my personal observation or I have recently heard from them by correspondence. Two were operated five years ago, and the others during the past four years and up to the present time. All of these patients are still retaining the globe excepting two in which the ball escaped, and the case of Dr. Means, in which the glass ball was removed eighteen months after insertion by another surgeon. These three cases are given in detail later.

Age at Operation.—The youngest person operated was one year old, and as the child had an extra large scleral cavity, having lost the eye from ophthalmia neonatorum followed by staphyloma, a good-sized glass ball was inserted in order to better assist in maintaining uniform growth of the two orbits. This child is here and I show the case to demonstrate the uniformity of orbital development. The oldest was 54 years. The only reason for not inserting a globe in old people has been the fact that

the cosmetic improvement has not been as important as in young people. The other cases ranged in age from 4 to 45 years.

Time in Hospital.—Regarding the time in the hospital, while at first I kept them in two weeks I now keep them only one week. I do not consider it essential to keep the patient in bed but a few days, and simply advise him not to use his good eye for reading and not to exercise to any extent for two weeks, though I keep the patient under observation for two weeks and prefer to see him one month. The prosthesis was inserted in some cases two weeks after the operation, but in many cases it

was thought best to wait a month, and in one case two months, before allowing the patient to continually wear the glass eye. Regarding the material used as an artificial vitreous I have used glass in all my cases. Fox sums up this point as follows: "I prefer first glass balls, then gold balls. Silver balls produce argyria and undergo oxidation. Aluminum globes become disintegrated, as was shown in one case where I used a perforated aluminum sphere as suggested by Dr. Bryant. This caused great pain, because the tissues surrounding it grew through the opening."

Contra-indications.—As absolute contra-indications I

would consider malignant intra-ocular growths and diseased or injured eyes which have already excited sympathetic ophthalmitis; extreme phthisis bulbi and sclera so lacerated that it could not be made to retain an artificial vitreous. (Frost-Lang operation may be substituted in the two latter classes.)

I have had no experience in performing Mules' operation where sympathetic ophthalmitis was present and have not cared to be a pioneer in the experiment, though I could not have had worse results than I have had fol-

lowing enucleation in these cases, for in the few cases I have had, enucleation failed to permanently arrest the disease, and in two cases at least total blindness ultimately resulted. Old diseased eyes, traumatic or non-traumatic, in which iridocyclitis is present and which are said to be most apt sooner or later to cause sympathetic ophthalmitis, even when sympathetic irritation is present, I do not consider contra-indications and have on my records 7 such cases; in all these cases there has been a complete and immediate subsidence of the sympathetic irritation and no return. Three years ago, however, I enucleated such an eye because the patient considered

the time element more important than cosmetics and he has since had attacks of sympathetic ophthalmitis. Had I inserted an artificial vitreous in this case I probably would have considered the operation at fault. It was simply a case in which enucleation failed to arrest a developing sympathetic ophthalmitis.

Phthisis bulbi if not extreme need not prevent the insertion of a glass globe within the sclera. Case 28 was one of phthisis bulbi. A small globe was inserted into the scleral cavity. Glaucoma is not a contra-indication if we may judge from the success attained in Case 2.

Neither do I consider suppuration of the globe a con-

tra-indication, for in all such cases the results have been highly satisfactory and suppuration has ceased with the operation. I have had four such cases, of which the worst was that of J. T. M., male, age 28, machinist. A piece of brass entered the vitreous. Mules' operation was performed ten days after the injury, the vitreous chamber was full of pus, but the tissues healed as readily as any case. The fact that the presence of tuberculosis of the iris is not necessarily a contra-indication if limited to the iris is evident from the record of Case 1. In this case Dr. Allport had performed an iridectomy for tuberculosis of the iris, and upon recurrence Mules'

operation was performed. Gunshot wounds are not contra-indications unless the sclera is too much lacerated or shot is present in the orbit and causing irritation. I have had only one such case, and in this the shot had passed completely through the ball and become embedded in the bone. Mules' operation was performed with success.

Failures would include escape of artificial globe, the development of sympathetic inflammation and a stump over which an artificial eye can not be worn because of its irritability.

Sympathetic Inflammation.—No opinion is more valu-

able on this point than that of De Schweinitz, who concludes after his thorough investigation of 317 operations by 32 operators, as follows: "Eyes so wounded that they are likely to excite sympathetic ophthalmitis if seen before two weeks have elapsed need not be enucleated, that is, evisceration or Mules' operation may be performed, because, *with perhaps the exception of a single case, there is no positive proof that those operations have of themselves excited sympathetic disease. They may fail to arrest the development of sympathetic ophthalmitis, just as enucleation may meet with a similar failure.*" (Italics mine.)

Dr. de Schweinitz reports painful stump requiring removal three times. The only case in which I have had

this experience is the one of Dr. Means, whom I assisted two years ago while in Columbus.

J. R., male, 22, left eye blind and at times some pain. Right eye in good condition. Mules' operation performed as usual, no reaction, recovery uneventful. From time to time Dr. Means wrote that the patient, who was away at college, complained of inability to wear an artificial eye, and at my suggestion the Doctor removed some of the stitches which he thought might be causing the irritation. May 17, Dr. Means writes in reply to my letter as follows:

"The patient you refer to has had the glass ball removed. He came to me last December and I removed another stitch that was deeply embedded in the upper fornix. I thought

probably this was causing the irritation. He returned to college, and claimed that his eye did not improve. I received a letter from him later stating that the other eye was becoming irritated. I wrote him to come to me at once, but received a letter from an oculist in his college town, saying that he had removed the glass ball to relieve irritation. He simply removed the ball and allowed the sclera to remain. The Doctor intimated that the ball was too large, and therefore made undue pressure. I have operated on only one case since, and it has proven an entire success."

I am not prepared to explain this case, but do remember that we used a ball which was quite tight and

think the explanation of the surgeon who removed the globe quite probable.

Escapes are due to the following causes: reaction, hemorrhage, sepsis and consequent sloughing of edges of the sclera, imperfect ball, too large ball, too small ball, improper coaptation of sclera, use of absorbable sutures, insufficient number of sutures, too early wearing of prosthesis, and later sepsis introduced by means of the prosthesis or otherwise, resulting in abscess.

Reaction.—Those of greatest experience at first had a great deal of inflammatory reaction following their operation, and Fox attributes many of his early "escapes" to this cause. In the 317 cases tabulated De Schweinitz reports excessive reaction 77 times. The only case in which I have experienced any greater reaction than following enucleation are in Cases 1 and 2; only one of these could be called excessive. These I attribute to the fact that just before inserting the globe the cavity was dried out with iodoform gauze, enough of the powder remaining to cause the inflammatory reaction, and I believe a common cause of reaction is the use of irritating antiseptics. I have found very hot sterile water of the most service, because it acts as an antiseptic, is non-irritating and effectually stops hemorrhage, which may in itself be a cause of expulsion.

Moist dressings are undesirable because they promote suppuration, and a dry, sterile or non-irritating antiseptic dressing with a pressure bandage will aid in preventing reaction. This bandage should not be removed before two or four days have elapsed, unless complications arise, and the good eye need not be bandaged at all though the patient should be cautioned to keep it quiet.

While an irritating antiseptic should be avoided as much as possible, and if used it should be followed by irrigation with hot sterile water, yet a condition of asepsis is important and the balls should be sterilized in boiling water, and before inserting should be examined to discover any small hole through which the water may have entered. Occasionally such imperfect glass globes will be discovered.

The ball inserted should be as large as the scleral cavity will admit and yet small enough so that the edges of the sclera will not be pulled apart during the healing process. Case 15, already spoken of as being the only case in which there was an escape, I believe is an example of the use of a too small ball.

Laura D., aged 4, consulted Oct. 31, 1899. Dirty wire cut left eye yesterday while at play. Find large wound across cornea with escape of contents of the globe and evidence of infection. Operation delayed till November 2, awaiting consent of mother. November 2, performed Mules' operation as usual, the only difference being that a globe was inserted smaller than the cavity would have admitted comfortably, because I had on hand no globes of medium size and could not obtain one in the city. There was no reaction and result was successful. An eye was adjusted (date not recorded) and patient returned to her home one hundred miles away. A letter was received Feb. 21, 1900, stating that on February 11 the operated eye began to "look sore," and for the next two days "she had a good deal of pain." After which there was relief, but the glass ball became visible, and February 20 it came out (three months and eighteen days after operation). My opinion of this case is that the ball being too small to become firmly encysted was movable in its cavity and acted as an irritant.

The sclerotic coat is so non-vascular that union does not quickly take place, and hence the importance of careful coaptation of the cut edges and the necessity of using many strong, non-absorbable sutures. These be-

come encysted and do not often cause irritation, but if they should the conjunctiva can be opened and the irritating sutures removed after firm union has taken place. Occasionally it has been found necessary to suture the conjunctiva during convalescence because it had separated.

As a rule, it is best to wait a month before inserting a prosthesis for permanent use, and when first used, the patient should be seen daily to ascertain the effects. The irritation produced might be sufficient to cause the recently united scleral edges to part. If this seems imminent, it may be necessary to wait a week longer, when the artificial eye may be tried again. If the sclera separates, it should be reunited with sutures as before. In my experience this has never happened.

The second case in which there was an escape of the glass ball was that of W. M., whose eye was operated upon May 28, 1898. It was an old case of recurrent iridocyclitis with sympathetic irritation. The original trouble had been gonorrheal ophthalmitis, which had occurred two and one-half years previous to the time Mules' operation was performed. The results of the operation were successful and the patient had no trouble of any consequence until May 18, 1901. While doing the ordinary work on a farm and wearing the prosthesis, not taking pains to keep the prosthesis clean, the tissues became infected and the result was that when he reached the city four days later I found the glass ball protruding and an abscess of the orbit present. Pus was found in the orbital tissues and in the sclera back of the glass ball. The glass ball was removed and the cavity irrigated daily. The stump is now quite healed. The escape in this case was clearly due to sepsis introduced on account of the carelessness on the part of the patient, who was constitutionally unclean.

I have always found great difficulty in securing a proper fitting glass eye to cover the stump after Mules' operation. After a successful operation the only thing required to secure perfect appearance is a well-fitting eye of the correct shade, and hence one feels like being very particular in the selection. Most of the eyes now on the market are not fitted to be used for these cases. To fit well and give the best results the iris should be placed well up and not far toward the internal side, leaving a broad white margin below while the temporal side of the eye should not be as large as the usual prosthesis: an extreme convexity is undesirable as such eyes appear too prominent when in place. The customary notch should be very slight, or may be absent.

In conclusion, I wish to state that I believe Mules' operation has come to stay and that as time goes on the percentage of failures will decrease and it will be more universally performed. The following are the advantages:

1. Absence of tears and secretion.
2. Development of orbit maintained.
3. Absence of enophthalmus.
4. Moisture of prosthesis maintained.
5. Mobility.

6. In general the satisfaction to the patient, because of natural appearance and consequent absence of self-consciousness, which is bound to make a vast difference in the success and happiness of the individual.

602 Nicollet Avenue.

BIBLIOGRAPHY.

1. Allport, Frank: *Northwestern Lancet*, Feb. 1, 1897.
2. Fox, L. Webster: *International Clinics*, vol. ii, 7th Series.
3. Fox: *International Clinics*, vol. ii, 8th Series.
4. Fox: *Transactions of the Section on Ophthalmology*, Amer. Med. Assoc., 1898.

5. Fox: The Medical Bulletin.
6. De Schweinitz, G. E.: The Comparative Value of Enucleation and the Operations which have been Substituted for It.
7. Todd, Frank C.: New York Medical Journal, Sept. 17, 1899.
8. Transactions of the Ophthalmological Society of the United Kingdom, vol. xviii, 1898.

DISCUSSION.

DR. FRANK ALLPORT, Chicago—Failures in this operation are more or less frequent from the ambitious desire to fill in the scleral sac to its utmost capacity, in order to provide a large stump upon which to place the artificial shell. Considerable reaction and swelling usually follow the operation, and if the ball is too large, stitches are liable to pull out and expulsion of the ball will follow. An important point in this operation is the use of plenty of scleral sutures. Some operators make a rule to insert three, four or five sutures. I use as many as the incised sclera will hold, and place them well back in the scleral tissue. I found it useful to follow the suggestion of Dr. Prince, of Springfield, in swabbing out the interior of the scleral sac with a 95 per cent. solution of carbolic acid just before inserting the globe, and neutralizing the acid immediately with alcohol. It certainly adds to the safety of the operation by rendering the interior of the sclera as aseptic as possible. I insert my conjunctival sutures very far back in the conjunctival tissue, as I was very much annoyed by the conjunctival sutures pulling out and leaving a central opening in the conjunctiva, thus exposing the scleral sutures. In my last few cases I have placed the sutures as far back in the conjunctiva and underlying tissue as was practically possible, and I have not since been annoyed by the unpleasant and unfortunate gaping of the edges. I always wait until hemorrhage has practically ceased before inserting the glass globe. Some operators pay little attention to this point, but the hemorrhage following the curetting of the sclera is usually excessive and while the wait is tedious, I believe it better to endure it than to be in too great haste to insert the globe. The hemorrhage can be controlled to a certain extent by the use of very hot bichlorid solutions and packings with iodoform gauze. I do not like aluminum globes, because I saw one case in the practice of a colleague in Chicago where the globe greatly disintegrated and had to be cut away with considerable disfigurement to the orbit. Up to the present time I have had unvarying success in my Mules' operations, but I am sorry to say that I have to report my first failure. It was a case in which I found a calcareous lens and orbit. When I inserted the scleral sutures in this case, I noticed that the tissue was fragile, and I should have had better judgment than to have gone ahead with the operation, but never having had an experience with an eye of this kind, I completed the operation. I subsequently lost this ball, and had to treat the case as an ordinary eversion. I think this case should teach us the lesson never to make a Mules operation in a calcareous eye, for I believe that in such cases we will always find the sclera incapable of standing the necessary pressure.

DR. MELVILLE BLACK, Denver—Concerning the use of Snellen's eyes I believe that with their use this operation in the future will not be so frequently performed. In my experience with the Snellen eye, covering a period of two years, it has been so satisfactory that I have abandoned the Mules operation entirely and have had just as good cosmetic effects as with the implantation of the glass ball. We secure by the use of this shell all that the Mules operation can give us without the long delayed healing process, together with the fact that a large percentage of these people are unable to spend the time necessary to secure a good cosmetic result.

DR. H. FRIEDENWALD, Baltimore—I should like to make the statement that insertion of the glass globe after enucleation does not prolong the period of healing; it may after the Mules operation. I have not had experience with that. I have had only one case in which the ball came out and in that case I had used a gold one. In my short experience with the Snellen eye I am inclined to think we shall probably do these operations more rarely than we used to.

DR. H. MOULTON, Fort Smith, Ark.—One of the arguments advanced for the Mules operation is that the orbit, in the case

of children, develops symmetrically with the orbit of the other eye. I have observed a number of cases in which the globe had atrophied in childhood and in every one of them the development of the orbit was very insufficient and in some the growth seemed to have absolutely ceased at the time the atrophy took place. I have wondered why operators should expect that the orbit would develop more after the Mules operation than in these atrophy cases, and I would like to ask if operators are satisfied that they have observed in practice such a development of the orbit as they expected after the Mules operation.

DR. F. C. TODD, in reply—In regard to the development of the orbit my experience has not been long enough to observe it in many cases. One of the cases just shown demonstrates the uniform development of the orbits. In that one, however, there was a large anterior staphyloma and I was able to insert a large ball, which must certainly assist in the development of the orbit. This was the one operated when one year old.

THE TREATMENT OF THE ACUTE PSYCHOSES IN PRIVATE PRACTICE.*

C. EUGENE RIGGS, A.M., M.D.

ST. PAUL, MINN.

The various forms of mental perversion usually first come under the observation of the general practitioner. As for the acute insanities, their prognosis varies in proportion to their early recognition and treatment. Skilful attention at the outset will frequently avert an attack, or, at least, cause to run a benign and favorable course that which would otherwise have developed into a prolonged and serious illness. "In regarding the causation of insanity," says Dr. Mercier, "we should not exclusively concentrate attention upon the nervous system, but should remember that this is a question of general medicine. We must not wrap ourselves up too closely in our specialty; we should take wide views, and regard insanity as a malady, not of the brain alone, but of the whole man; it is a disease of the whole organism and it can only be properly understood and properly dealt with when so regarded."

Since the representative medical schools have made a course in psychiatry an essential part of their curriculum, the attitude of the profession has radically changed with reference to insanity. It is a comparatively short time since it was thought that only a state hospital or private institution for the treatment of the insane could care for these patients. It is now believed by some of the most experienced in our profession that no person should be sent to an institution who is a suitable subject for home treatment. I am convinced, as the result of a large experience, that home care in properly selected cases (and by "home care" I mean that either in a private family or a private hospital) will shorten the length of an attack by a period varying from a few weeks to several months. There can be no question, not only that the association of the insane with each other adds greatly to their suffering, as the result of personal contact, but that auto-suggestion intensifies and prolongs indefinitely the morbid mental state. It is the absence of these conditions, together with the opportunity for individualization of treatment, which makes home care so desirable and successful.

I am far from being antagonistic to our state hospitals; on the contrary, I am most appreciative of the great work they are constantly doing. They have a defin-

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Nervous and Mental Diseases, and approved for publication by the Executive Committee: Drs. Frederick Peterson, Hugh T. Patrick and H. A. Tomlinson.