Great Northern Hospital (recorded at page 576). Although this opening led into the joint of which we have spoken, there were still no projecting fragments issuing through it.

For the subjoined notes we are indebted to the kindness of Mr. W. P. Swaine, house-surgeon to the hospital:—

Benjamin C. P.—, aged thirteen, a native of London, of a strumous habit of body, was admitted Nov. 5th, 1857, with extensive disease of the lower portion of the femur and knee-joint. On admission, the knee was permanently flexed at an angle of about 45°, swollen, and much inflamed; the patella was displaced outwards and completely ankylosed. Just on the inner border of the external condyle there was an opening through which a probe could be passed both inwards and outwards, but only in the former direction could bone be felt. The femur, on its anterior aspect, from the condyles to about the middle of the shaft, was considerably thickened by deposit of new bone, causing the femur to appear curved from before backwards. On the outer aspect, at about the middle of the thigh, was a small opening, through which bare bone could be felt by a probe.

The patient states that about last Christmas he first experienced difficulty in extending his leg. This gradually increased; the knee became very painful; and about a month after, he first noticed the opening above the knee. At this period the joint had become perfectly immovable. He had no advice, however, until the end of September, when he attended a London hospital; but receiving no benefit, he was recommended to see Mr. Fergusson, who advised him to come into King's College Hospital.

On Nov. 14th, the inflammation about the parts having in the meanwhile subsided, the patient was brought into the theatre, and, being placed under the influence of chloroform, Mr. Fergusson proceeded to remove the necrosed bone from the lower portion of the femur. In order to effect this, the opening in the middle of the thigh was enlarged; and after a great deal of new bone had been removed with the bone forceps, Mr. Fergusson was enabled to extract, as he considered, the whole of the necrosed bone. The opening near the external condyle was then enlarged, and necrosed bone removed from that side also. After the operation the boy progressed favourably; the discharge decreased; and the cavity in the femur appeared to be covered with healthy granulations.

Under these circumstances, Mr. Fergusson thought the patient was in a favourable condition for excision of the knee-joint, in order to straighten the limb, and give the patient a useful leg.

Dec. 5th.—The patient was again brought into the theatre, and whilst under the influence of chloroform, Mr. Fergusson proceeded to excise the joint. Having made the usual **H** incision over the knee, he dissected back the integuments, exposing the condyles of the femur, the patella, and head of the tibia; he then sawed through the femur just above the condyles, and finding that the bones did not come into apposition after a slice of the tibia had been removed, he took away a second portion from the end of the femur, and then divided the tendons of the hamstring muscles subcutaneously, and was enabled to put the limb in a good position; the flaps were then brought together by sutures, and the limb placed in the usual excision splint. After the operation, the boy suffered very little from shock, passed a quiet night, and on the following morning was surprisingly well, considering the severity of the operation which he had undergone.

7th.—This morning the lint was removed from the knee, and water-dressing applied; healthy suppuration has already set in, the limb is in perfect position, and the general condition of the patient leads us to hope for a favourable result.

ST. GEORGE'S HOSPITAL.

SUCCESSFUL RESULT OF THE TREATMENT OF A CASE OF ANEURISM BY ANASTOMOSES OF THE LEFT EAR OF A YOUNG MAN.

(Under the care of Mr. PRESCOTT HEWETT.)

In a previous "Mirror," (Nov. 14th, p. 496,) we gave a report of a case of aneurism by anastomoses of the left ear of a young man, which was treated by the electric cautery, styptics, and ligature of the tumour itself. The subsequent results were of the most satisfactory character; the tumour sloughed off, healthy granulations sprang up, and the ulcer healed. The swelling of the ear subsided, and it resumed somewhat of its natural shape; the pulsation and constant beating about the ear and head disappeared; all the vessels became obliterated, excepting a small part of the tumour situated over the edge of the pinna, which was not touched at

the first operation, but which was now quite soft and flaccid. The posterior auricular artery and other vessels resumed their natural size, leaving the deep groove quite prominent over the mastoid process of the temporal, (itself hypertrophied,) and which would with ease admit the radial artery itself. On the 10th of December, Mr. Hewett, without giving chloroform to the patient, strangulated the small tumour on the edge of the pinna, and he is now quite cured.

This is a result which is not often met with in these dangerous cases in adult life. The case referred to in our previous report as at the present moment exciting the attention of the surgeons at the Hôpital Beaujon, in Paris, was submitted to ligature of the opposite carotid artery to arrest the bleeding, and with success, the other carotid, on the same side as the disease itself, having been tied about forty years ago. The hæmorrhage has ceased since this latter operation, and the man is alive, and apparently rational, with both of his carotids obliterated.

CLINICAL RECORDS.

WEBBED FINGERS IN A YOUNG MAN.

It is the almost invariable custom at the present day to remove any congenital deformities in childhood, as they are more easily dealt with at that period of life, and with less danger to the little patient, than when left till adult age is reached. There is less risk in operating upon the child than upon the adult. Now, if nævi, tumours, deformities, and other vices of development are not eradicated in early life, they may give much trouble at a later period. In our "Clinical Records" of the 25th of April last, we said a few words upon a case of webbed fingers in a child at King's College Hospital, treated by Mr. Fergusson. We did not meet with another instance till the 10th November, at Guy's, when a young man, with the middle and ring fingers of both hands in a congenitally webbed condition, was brought into the operating theatre. Both fingers in each hand were of the same length, and, under the influence of chloroform, Mr. Hilton divided the integument of the united fingers of the left hand first, leaving the right to be dealt with on another occasion, should that on the left heal up without the supervision of erysipelas or other bad consequence. The most curious part of the history of this young man is, that he thought this condition of his fingers was a natural one until pointed out by some one to be otherwise. He had not suffered any inconvenience from them.

MUCOUS TUBERCLES FROM THE VULVA.

No better proof can be offered of the truth of the observation, that these growths arise from dirt and filth on the part of the patients themselves, than that of their occurring amongst the lowest orders, and at the same time the most uncleanly. Plenty of soap and water does an immense deal in preventing their appearance, together with dryness of the cutaneous surface. In the upper walks of life, when blenorrhagic affections may have been contracted, it is very seldom that we see mucous tubercles about the anus or vulva, because cleanliness is scrupulously attended to. On November 17th we saw Mr. Brooke remove a large number of these growths from the vulva of a young girl at the Westminster Hospital with a pair of scissors; they had resisted other treatment, and he considered this the best mode of getting rid of them. The bleeding was somewhat troublesome at first. Some months back we recorded the case of a girl under Mr. Curling's care, in whom these growths were very extensive indeed; they were removed with a pair of scissors, and she did well. The result of the treatment proves, together with what we know of their structure, that they are confined solely to the epithelial or epidermal layer of the skin, the true dermis beneath not being involved.

RECURRING TUBERO-CYSTIC TUMOUR OVER THE STERNUM.

WHEN we saw the patient, who was the possessor of this adventitious growth, laid upon the table in the operating theatre of St. George's Hospital, on the 26th of November, we at first sight imagined the diseased mass to be an aneurism by anastomosis, with discoloration of the integument; but in the progress of events which followed, it turned out to be otherwise. The subject of it was a very stout, elderly woman, with what would be considered a very great development of the adipose tissues, which showed itself especially in the mamæ,