ment himself. Dr. Todd performed the operation in my clinic very successfully, even with the very inferior instrument in my possession.

DR. OSCAR WILKINSON, Washington—I have been with Dr. Worth in a great many of his operations and have watched the results, and in my own practice no other advancement operation has given such good results. The operation depends largely, of course, on the object we wish to obtain and the amount of defect to be corrected. Where there is a large defect I should be skeptical as to the results of the tucking operation. It may be very valuable where the defect is slight, and it is certainly an ingenious idea.

DR. MARK D. STEVENSON, Akron, Ohio—I make a long, vertically curved incision through the conjunctiva, the center nearly touching the limbus, the ends curving away from the cornea. Tenon's capsule is next opened and the conjunctiva and capsule being pushed well back the muscle is seized by my advancement forceps and then cut free from the sclera. A sharp needle on a silk suture is next passed from without inward through the junction of the upper and middle third of the tendon and is then passed outward through the upper margin of tendon, and again passed through the tendon just behind the first point of entrance. It is then carried near the edge of the cornea and passed deep through episcleral tissue into the sclera. The needle on the other end of the suture is likewise used on the lower portion of the tendon and the two being first tied in a single knot, are tightened until effect required is secured, when a surgeon's knot may be made. All superfluous tissue should be removed. By this suture the upper and lower portions of the tendon are securely held, while the middle portion also affords support by the cross piece of the suture. The circulation in the middle portion is not interrupted, however, and only one knot is necessary.

DR. JOHN E. WEEKS, New York City—Any operation that necessitates doubling the tendon or muscle into a cord is certainly theoretically imperfect. The operation of Dr. Savage is such an operation. That difficulty has been avoided to some extent by the last speaker. The ideal operation is one that will maintain the normal width of the tendon, in addition to accomplishing the result desired. In order to maintain the width of the tendon, sutures must be entered at the upper and lower borders of the tendon. If only two sutures are used the middle portion can not be fastened as it should be, but with three sutures the width of the tendon can be controlled perfectly. My experience has led me to the conclusion that deviation of the muscle in any way imperils the result of the operation. For some years I used the advancement by tucking as devised by de Wecker, with very good results. One can not tell by half a millimeter the amount of the effect that will be produced by any operation. Another thing that is detrimental is the strangling of a portion of the tendon. That is done when the middle or the sides of the tendon are tied. It can be avoided by quilting the sutures. It is important to go well into the scleral tissue in passing the bulbar sutures. Another thing, do not delude yourselves in believing that a sufficient plastic process will be set up in four or five days to retain the advanced muscle. It ordinarily requires a longer time. Leave the sutures in from seven to ten days in order to be assured of the permanent retention of the tendon in the position in which it is placed at the time of operation.

DR. W. E. LAMBERT, New York City—I should like to substantiate what Dr. Weeks has just said. I am not an advocate of the tucking operation. I find advancements, or resections and tenotomies perfectly satisfactory.

DR. J. L. BORSCH, Philadelphia—Dr. Todd in his paper says that he can overcome a secondary strabismus by tucking the muscle that allowed the eye to deviate. The tendon, in a secondary strabismus, is so retracted that in most cases it can hardly be found; and what is advanced is the capsule, and the eye is thus straightened. It has been my privilege to assist de Wecker in many operations for capsular advancement, and have performed many myself, and fail to understand why an operation so simple and so effective should be modified. The manner of placing the sutures, it has been said, prevents their

slipping; if the sutures are of the proper kind this will not occur. Dr. Todd assures, you can regulate the effect exactly with his method. This can also be done by grasping the capsule in the broad forceps, but in the majority of operations one must overcorrect a few degrees for a day or two. As for judging the amount of deviation corrected, I wish to say, that by using the de Wecker strabometer, which is extremely simple and portable, you can measure the exact deviation of the eyes and regulate the operation accordingly. I venture to say that the fraternity who do many operations will bear me out in the statement that we must simplify and not complicate operations in ocular surgery. For myself I am not ready to abandon the de Wecker operation.

DR. F. C. TODD—The great number of operations that have been described is evidence that we are not satisfied with what has been done so far in this work. As to Dr. Savage's criticism that torsion could not be effected by my operation, I would say that that is one of the advantages of the use of the instrument; the torsion can be easily regulated. In the second case I operated on I was obliged to produce torsion. By changing the axis of the instrument and inserting the scleral sutures above or below, as the case may be, the desired effect may be readily produced. Of course, the idea of the ligatures above and below is to prevent the cutting of the suture along the fibers of the tendon, which takes place in many of these operations and, as it seems to me, would take place in Dr. Savage's operation. I would remind Dr. Weeks that while he cuts off that portion of the tendon I ligate it, and I can not see that it makes any difference whether it is removed by atrophy or cutting; the results are the same; we do not care to preserve that portion of the tendon. Another advantage of this method is that the flap lies down and makes a smaller lump, which later disappears. In regard to the administration of chloroform, in these cases I rarely use chloroform; I prefer not to. As to the matter of pain, of course, many of these operations are somewhat painful, but that depends largely on the pulling of the muscle. With this instrument the pull is gradual and steady, while the pain is caused mostly by jerking the muscle. Now someone has spoken of doing a tenotomy. Each case must be treated on its merits. I would suggest that where one eye is involved it might be wise to do a tenotomy and an advancement, but where you have an alternating strabismus or where the movement of the two eyes is about equal we might desire to do two advancement operations.

THE PASSAGE OF DIFFERENT FOODSTUFFS FROM THE STOMACH.*

WALTER B. CANNON.

Assistant Professor of Physiology, Harvard Medical School.
BOSTON.

In the transfer of food from the stomach to the intestine two factors are concerned, the gastric waves of peristalsis and the pyloric sphincter. It is now known that under normal conditions the peristaltic waves course from the middle of the stomach to the pylorus without interruption during the whole of a digestive period. How the waves will affect the food is determined by the action of the pylorus. If the pylorus remains closed, the repeated waves over and over again churn the food with the gastric secretions; if the pylorus opens, these same waves act as the motive power to force the food onward.

The determinative function of the pylorus was recognized by the fathers of medicine, for they gave the ancient terms *pylorus* (keeper of the gate), *rector* and *janitor* to the sphincter, and assumed that it possessed a "will" which fixed the fate of ingested food. Van Helmont, in the days when the learned confused mysticism with science, presented his reasons for believing that the pylorus was the seat of the soul. It is certain that this

*Read at the Fifty-fifth Annual Session of the American Medical Association, in the Section on Pathology and Physiology.

wonderful sphincter holds the chances for good and evil in many of the most essential processes of digestion. The retention of food in the stomach to undergo gastric digestion, the gradual delivery of food from the gastric reservoir into the region of intestinal digestion, so gradual that the important digestive processes near the pancreas shall proceed regularly and without disturbance; the checking of irritants in their onward passage through the alimentary canal—all these functions are in the control of the pyloric sphincter.

Differing views of the way in which the pylorus opens and permits the food to pass have been set forth. Oppenheimer,¹ in 1889, and Rossbach,² in the next year, declared the food was all thoroughly mixed with the gastric juice, and that only four hours or more after eating did the sphincter relax and allow the stomach to empty. These observations, however, have not been substantiated; the errors in them probably were due to the nature of the method employed, for both clinical findings with the stomach tube, as Ewald and Boas³ have demonstrated, and studies with the x-rays on the undisturbed subject,⁴ prove that the stomach is not emptied all at once at the end of gastric digestion, but progressively during the period. In an investigation of the passage of food from the stomach we have, therefore, to consider agencies maintaining an intermittent closure of the pylorus, agencies which usually keep the channel closed, yet occasionally open it to permit portions of the digested food to advance.

Both mechanical and chemical agents have been supposed to determine the emptying of the stomach. Rossbach,² for example, thought that the stomach emptied only at the completion of gastric digestion, because the muscular contractions at that time were (under the conditions of his observations) more intense than before. Penzoldt,⁵ after an extensive study of the periods during which various foods remain in the stomach, noted that such foods as delayed the appearance of free hydrochloric acid remained longest in the stomach. Verhaegen,⁶ on the other hand, declared that it mattered little in the passage through the pylorus whether the food was acid or neutral. In criticism of these statements, it may be said that Rossbach's observations were made on morphinized animals, with opened abdomen, conditions known to disturb the normal activities of the digestive tract; and in Penzoldt's studies, although the amounts of food are stated, no attempt was made to have these amounts uniform, or to have the food of the same consistency. Furthermore, Penzoldt was dependent on securing remnants of the food by means of the stomach tube, a procedure which furnished no data concerning the rate at which the stomach was emptied, and was also objectionable in interfering with the digestive processes. This method, moreover, leaves the investigator in doubt as to which of the factors, the gastric peristalsis or the pylorus, caused the retardation of the food. Verhaegen's⁶ conclusions are based on observations with the stomach tube on only four individuals, two of whom were admittedly pathologic. In the presence of strong evidence against his conclusions, the small number of cases observed by him must furnish him inadequate support.

Because of the differences of opinion which the above discussion has shown to exist, and because of the impor-

tance which the passage of food from the stomach bears both to physiology and to practical medicine, further investigation of the factors concerned in the process is desirable.

THE METHOD OF INVESTIGATION.

In studying the agencies controlling the pyloric passage, it is well, for the sake of simplicity of conditions, to employ foods predominantly proteid, fat or carbohydrate. Such foods were used in this investigation. Boiled beef, free from fat, boiled whitefish and the white meat of fowl are examples of proteids which were fed; beef suet, mutton and pork fat are representatives of the fats; starch paste, rice, and potatoes, of the carbohydrates. In all cases full-grown cats, deprived of food for the twenty-four or thirty hours previous to the experiment, served for the observations. The food was invariably given in uniform amount, 25 cubic centimeters, and with uniform consistency, the consistency of thick mush. Before the food was given to the animals it was mixed with 5 grams of subnitrate of bismuth. Comparison of subnitrate of bismuth with other insoluble heavy salts has shown that it has no peculiar effects on the movements of the alimentary canal; and clinical studies by Schüle⁷ have proved that the addition of subnitrate of bismuth to food does not interfere with gastric motility. The animal thus fed with food mixed with bismuth subnitrate was exposed to the x-rays, and the movements of the food in the stomach and small intestine observed by means of the shadows cast on a fluorescent screen. The observations were recorded at regular intervals, one-half hour, one hour, and every hour thereafter for seven hours following the feeding. The records consisted of outlines of the shadows of the food, traced on transparent tissue paper laid over the fluorescent surface. Inasmuch as the diameter of the intestines varies only slightly, the area of cross-section of the contents may be disregarded, and the aggregate length of the shadows of the food in the small intestine may be taken to indicate the amount present. By comparing the aggregate lengths of the shadows cast by the intestinal contents it is possible to know the relative amounts of food in the intestine at different times as well as the relative amounts of different foods present in the intestines of different animals at the same time after eating.

It is obvious that the method has defects. For example, loops of intestine are not always parallel to the screen, and the loops not parallel may not always form the same angles with the screen, and they would, therefore, be variously foreshortened. It may be said in answer to this objection that the animals were stretched on their backs and the abdominal wall flattened both by the stretching and by the pressure of the fluorescent screen, so that the loops must lie nearly parallel to the screen, except, possibly, at short dorsoventral turns from one loop to another. That these turns were not a serious source of error was proved by massage of the abdomen, which rearranged the coils; tracings made before and after such treatment showed only slight variations in the aggregate length of the shadows. Another possible defect of the method arises from such overlapping of the loops that two loops, or parts of two loops, cast a single shadow. Care was always taken to obviate this error by separating by pressure with the finger loops lying close together. Still further faults of the method are found in the variations in the thickness of the food masses in the intestine, and the changes due to absorption, which proceeds at different rates with the different foods. But

1. Oppenheimer: *Deutsche med. Wochenschrift*, 1889, xv, p. 125.

2. Rossbach: *Deutsches Archiv für klin. Medizin*, 1890, xvi, pp. 296, 317.

3. Ewald and Boas: *Virchow's Archiv*, 1885, cl, p. 364.

4. Cannon: *American Journal of Physiology*, 1898, i, pp. 368, 369, 378.

5. Penzoldt: *Deutsches Archiv für klin. Medizin*, 1893, li, p. 545.

6. Verhaegen: *La Cellule*, 1897, xiv, p. 33.

7. Schüle: *Zeitschrift für klin. Medizin*, 1895, xxviii, 6, 500.

these defects must be regarded as slight, especially in the early stages of intestinal digestion, when compared with the great and characteristic differences in the amounts of food present in the intestine when carbohydrates, fats and proteids are separately fed.

What these differences in the treatment of the three kinds of foodstuffs by the stomach are, will now be stated. The results here set forth are based on between 600 and 700 observations on 80 to 90 cases.

RESULTS.

The characteristic differences in the aggregate length (in centimeters) of the intestinal content with the various foodstuffs at definite times after eating are illustrated by the following average figures:

Hours after feeding.	0-½	1	2	3	4	5	6	7
Fats (15 cases).....	5	9	14	17	16	13	13	9
Proteids (10 cases).....	1	5	18	21	23	19	17	14
Carbohydrates (14 cases).....	8	30	39	33	25	19	14	6

Note that the carbohydrates and fats both begin leaving the stomach soon after they are ingested; but the carbohydrates leave rapidly, while the fats leave slowly. The curve representing the carbohydrate rapidity of passage would have a quick rise to a high maximum at the end of two hours, then a slow drop; the curve for fats would be a long, low curve, with no prominent maximum point. The figures for proteids are of especial interest, for they show that the proteids begin to pass out much later than either carbohydrates or fats (in many cases of proteid-feeding nothing left the stomach during the first half hour, and frequently nothing was out at the end of an hour, though peristaltic waves were running regularly from the beginning), and, moreover, after the proteids begin to leave the stomach, they leave slowly. The curve for proteids would show a slow rise to a maximum at the end of four hours, and then a gradual fall.

Many confirmations of these figures have been secured. For example, when proteids are fed first and carbohydrates later, the proteids are pressed more into the pyloric while the carbohydrates lie more in the cardiac end of the stomach. Under these circumstances, the presence of the proteids near the pylorus delays the discharge of the carbohydrates, as the following comparison of average figures in two such cases with the average proteid figures will clearly show:

Hours after feeding.	0-½	1	2	3	4	5	6	7
Proteids (10 cases).....	1	5	18	21	23	19	17	14
Proteid first, carbohydrate second.....	2	6	15	21	29	34	28	22

On the other hand, when carbohydrates are fed first and proteids second, the presence of the proteids in the cardiac end of the stomach does not materially check the exit of the carbohydrates. A comparison of average figures in two such cases with the average carbohydrate figures will make this relation clear:

Hours after feeding.	0-½	1	2	3	4	5	6	7
Carbohydrates (14 cases).....	8	30	39	33	25	19	14	6
Carbohydrates first, proteids second.....	4	23	35	28	25	18	14	7

The most significant feature of the above tables is undoubtedly the remarkable difference between the figures for carbohydrates and for proteids. Why are the carbohydrates, which are not digested by the gastric juices, permitted to go quickly onward into the intestines, where they are digested, while the proteids, digested in the stomach, are retained there to undergo digestion? What fine mechanism of the digestive economy could see to it that the gastric part performs its function well when food it acts on is present, and lets it waste but little time or energy with food it can not change? Such apparently purposive functioning recalls the ancient attribution of a "will" to the pylorus, supposed to control the movements of material through the channel with def-

erence to the body's welfare. 1. there, however, a more simple and direct explanation of these activities?

If the presence of free acid in the stomach and near the pylorus is a stimulus to the relaxation of the pyloric sphincter, the interesting mechanism is largely explained. Both carbohydrate food and proteid stimulate the gastric secretion in abundance, as researches on dogs by Pawlow and his pupils,⁸ and as clinical studies on man, have shown.

Now, inasmuch as carbohydrates provoke an acid secretion and do not unite chemically with it, free acid is at once present in the stomach; whereas with proteids, since they join with the acid secretion to form acid proteids, free acid does not make its appearance for some time. Thus the carbohydrates would begin almost immediately to pass through the pylorus. Thus proteids would be delayed.

The hypothesis that the presence of free acid in the stomach opens the pylorus can be tested. Feeding a carbohydrate mixed with an alkaline fluid would delay the appearance of free acid in the stomach, and, according to hypothesis, should also delay the passage of the food. The similarity between the average figures from two cases fed on crackers moistened with 1 per cent. sodium bicarbonate, and the average figures for proteid, shows how closely the addition of the alkaline fluid makes the figures approximate.

Hours after feeding.	0-½	1	2	3	4	5	6	7
Proteids (10 cases).....	1	5	18	21	23	19	17	14
Alkaline carbohydrate	0.45	19	28	31	29	25	21	

The converse of this process also was tried. Proteids changed to acid proteids, and freed from all excess of acid were fed and the rate of discharge recorded. The following average figures of three cases of acid proteid (lean beef acid with hydrochloric acid), compared with the average figures for carbohydrates, show a remarkably close relation throughout:

Hours after feeding.	0-½	1	2	3	4	5	6	7
Carbohydrates (14 cases).....	8	30	39	33	25	19	14	6
Acid proteid	6	30	43	39	28	21	14	10

These two tables show that alkaline carbohydrate, like ordinary proteid, leaves the stomach slowly, while acid proteid leaves the stomach quite as rapidly as carbohydrate food. All cases which have been tested harmonize with the idea that conditions delaying the appearance of free acid in the stomach delay the discharge of the food, and conditions favoring its appearance hasten the discharge.

At first thought it might seem that acid added to carbohydrate food would still further increase the rapidity of the discharge. Such, however, is not the fact. When 0.4 per cent. hydrochloric acid is added to crackers in sufficient quantity to make a thick mush, the figures secured do not vary much from those secured when crackers and water are fed:

Hours after feeding.	0-½	1	2	3	4	5	6	7
Crackers and water.....	10	23	43	46	43	30	19	5
Crackers and 0.4 per cent. HCl.....	8	18	41	51	45	25	21	17

But if free acid is the stimulus to the opening of the pylorus, why does not the addition of free acid to the food open the pylorus and hasten the emptying of the stomach? How is the paradox to be explained? It is here that the reflexes from the duodenum must be considered.

In 1893 Hirsch⁹ observed that the introduction of acid into the duodenum through a duodenal fistula inhibited for a time the passage of food by the opening. Serdju-

8. Pawlow: The Work of the Digestive Glands, English translation, London, 1902, p. 35.

9. Hirsch: Centralblatt für klin. Medizin, 1893, xiv, p. 73.

kow,¹⁰ one of Pawlow's students, confirmed the work of Hirsch. It was impossible to tell by the methods these investigators employed whether the acid in the duodenum checked the gastric peristalsis, or compelled a prolonged contraction of the pyloric sphincter. Observations with the x-rays show that, although there is no cessation of gastric peristalsis after the advance of acid food into the duodenum, the pylorus does not for some moments relax. The duodenal reflex of Hirsch and Serdjukow must, therefore, affect the pyloric sphincter.

It is now possible to explain the slow passage of fats from the stomach. Fats are unlike either proteids or carbohydrates in that they are seldom eaten as a pure, or even predominant, foodstuff. When fats in almost pure form are fed to animals the gastric peristalsis is often observed to be slight, and, according to Pawlow, the flow of gastric juice is not stimulated. When fats enter the duodenum, however, they cause the same reflex that acid causes.¹¹ Thus, by a diminution of the influence opening the pylorus, and by a continuance of the influence closing it, the fats are long retained in the stomach, and only slowly passed along.

THEORY OF THE ACTION OF THE PYLORUS.

Evidence has been presented in the foregoing paragraphs which would explain the remarkable activities of the pylorus in the presence of food as largely due to chemical stimulation. The degree of contraction of the pyloric sphincter when at rest varies,¹² but the effect of particles of food in keeping the sphincter contracted⁴ indicates that the mere mechanical presence of food in the stomach will produce a closure of the pylorus. After food has reached the stomach all evidence points to the probability that the signal for a movement onward into the intestine is the appearance of free acid. Free acid on the stomach side of the pylorus would thus be the stimulus to open the pylorus and initiate the chemical control of the sphincter. The opening of the pylorus permits the exit of a portion of the acid chyme. There is proof that acid in the duodenum closes the pylorus and keeps it closed. Thus the acid chyme would go forth from the stomach, and, with its acid, close the gate behind it. But acid in the duodenum is known to be the stimulus for the flow of the alkaline secretions of the pancreas.¹³ No inorganic acid is normally present beyond the first few inches of the small intestine.¹⁴ The acid, therefore, is here neutralized. As the neutralizing of the acid proceeds the stimulus to the closure of the pylorus is weakened, until the acid in the stomach can again open the pylorus. Again the food would pass out and the gate close till the duodenal changes finish their slower course.

Thus, automatically, carbohydrates, on which the stomach works no change, would go early and quickly into the intestine, where they meet their proper ferment; thus, automatically, proteids would be retained in the stomach to suffer alteration by the gastric juice, and only after such alteration be permitted to go on; thus, automatically, the intestine would be saved from oppression by overwhelming discharges from the gastric reservoir; and, by this same automatic mechanism, the gastric secretion, harmful to the action of the intestinal fer-

ments, is rendered innocuous because of its admission, little by little, into the duodenum.

DISCUSSION.

DR. WINFIELD S. HALL, Chicago—I will note in a few words the relation between Dr. Cannon's findings and something that I have discovered in my experiments in dietetics in my department, namely: Patients with acute gastritis can take with comfort egg lemonade earlier than they can take any other form of food, and it is our custom to give that at the beginning of feeding after they have had the period of rest that Nature is calling for in an acute case of gastritis. The earliest food that we can give is very small doses of egg lemonade every two hours. Dr. Cannon's discoveries regarding the influence of acid make it evident that this egg lemonade, being in the condition of acid albumin and already half digested, passes readily right on through the stomach, without exciting or irritating it, and down into the duodenum, where its digestion is completed.

DR. CHARLES HARRINGTON, Boston—Did the gluten curve take that form on account of the presence of carbohydrates, which are commonly found in large amounts in commercial gluters?

DR. D. H. BERGEY, Philadelphia—How does Dr. Cannon account for the prolonged time of absorption from the stomach? Our usual idea has been, I think, that food passes from the stomach in a much shorter time than his curves indicate. Does he believe that the food really remains in the stomach for that length of time, or is it due to the shadows cast by the substrate of bismuth remaining on the mucous membrane?

DR. FENTON B. TURCK, Chicago—One interesting point that Dr. Cannon has brought out here is that the presence of acid or alkali does not materially influence the peristaltic motion and that the chemical composition of the food does not influence the motions of the stomach. The peristaltic waves he found extending over to the pylorus were independent of the character of the stomach contents. In this difference between starch and proteid, the more rapid passage of the starch was not due to increased peristaltic waves caused by carbohydrates, nor to lessened peristaltic waves influenced by diminished stimulatory action of the proteid on the muscle fibers of the gastric wall, but to difference in the opening or closure of the pylorus. This somewhat corresponds with my own observations, published last year, on the influence of various chemical bodies on the gastric mucosa. To the entire mucosa were applied various chemical bodies—acids, alkalies, varieties of food, such as bouillon, peptones, starch—and they did not seem to influence the peristaltic motions a particle; but as soon as tension was brought to bear on the muscle wall of the stomach by distending the stomach with air or water, introduced directly, or indirectly in a rubber bag inserted in the stomach, it promptly caused rhythmical contractions—peristaltic waves. The degree of tension or intermittent tension, within limits, increased these peristaltic waves proportionally. By prolonged and excessive tension, lasting several hours, the gastric muscle of the dog became fatigued and resulted in dilatation of the stomach. I also observed, as Dr. Cannon has done, the influence of acid on the mucosa of the stomach and duodenum, causing the opening and closing of the pyloric door; and this work that Dr. Cannon has brought out in another direction not only confirms my observation of the negative effect of chemical stimulation or excitation of stomach movements, but goes further and gives an explanation of the manner in which the pyloric door is opened. Does this, then, conflict with the popular theoretical idea that excessive acid in the stomach causes spasm and a closure of the pylorus, as is held by clinicians? I am impressed more with the physical and mechanical effect of the stomach contents in causing rhythmical contractions, which necessarily includes relaxation as well as contraction. The acidity may be an incident indirectly influencing the passage of food through the pylorus.

DR. CLAUDE A. SMITH, Atlanta, Ga.—Has Dr. Cannon made any observations on the effect of exercise on the passage of these different foods from the stomach?

10. Serdjukow: Reviewed in Hermann's *Jahrsbericht über die Fortschritte der Physiologie*, 1899, viii, p. 214.

11. Lintwæw: *Biochemisches Centralblatt*, 1903, I, p. 96.

12. Oser: *Zeitschrift für klin. Medicin*, 1892, xx, p. 285.

13. Dolinsky: *Archives des sciences biologiques de St. Pétersbourg*, 1895, III, p. 399.

14. Moore and Rockwood: *Journal of Physiology*, 1897, xxi, p. 373.

DR. W. B. CANNON—In order to find the reason for gluten leaving the stomach early and in large amounts, just as carbohydrates do, I took equal amounts of gluten paste and of the proteid (meat proteid) which I had been using, and put into each the same amount of 0.4 per cent. hydrochloric acid. In about thirty minutes the free hydrochloric acid had entirely disappeared from the meat, but the free hydrochloric acid was still present in the gluten paste twenty hours after being introduced. Evidently this gluten does not readily unite with the acid; so that, if acid is poured out on the gluten in the stomach, free acid must be present immediately, just as it is in the case of carbohydrates. I used Dr. Johnson's gluten, and I understand that a large amount of starch is present in it. The figures I have presented do not show the amount of food in the stomach. The figures show the amount of food in the small intestine. The stomach may be empty while food is still present in the small intestine. It is true, as Dr. Turek has suggested, that I have seen no considerable variation in the peristalsis. Gastric peristalsis runs regularly from the beginning to the end of digestion unless stopped by emotional disturbances. I have not examined the effect of exercise, but one of my students is at the present time at work on the effect of exercise and other factors.

THE DIAGNOSIS OF ENLARGED BRONCHIAL LYMPH NODES.*

ALFRED FRIEDLANDER, M.D.,
CINCINNATI.

Among the phenomena of the developmental period of childhood, there is none more striking perhaps, than the activity for the lymphatic system. The lymph glands constitute an apparatus for filtration. Given the factors of rapid growth, on the one hand, and of poorly developed powers of resistance, on the other, and it can readily be seen that the period of childhood must necessarily be accompanied by marked reactive irritation and inflammation of the entire glandular system. During this period the mucous membranes are prone to irritation, with resulting inflammation. Added to this, there is the disposition, often hereditary, not infrequently aggravated by faulty conditions of development and hygiene, to chronic inflammation of mucous membrane, skin and bone. Under such conditions it is not surprising that the filter system should be taxed to its utmost capacity—hardly less astonishing that we should find very often, in the inability of this system to hold the various poisons localized—an avenue for general systemic infection.

From the nature of things we should expect to find inflammation of the tracheobronchial glands to be enormously common. The three groups of these glands—the tracheal, the tracheobronchial and the peribronchial intercommunicate freely. They receive the lymphatics of the trachea, the bronchi and the lungs and the pleuræ. But, more than this, through the deep lymphatics of the neck, lying both in front of and behind the carotid sheath, they are in direct communication with the cervical glands. Again, these glands are directly connected with the mesenteric lymph nodes.

It will thus be apparent that very many different factors and very diverse conditions may induce inflammation of these glands. Of these factors there are many that are particularly frequent in childhood. Diseases of the nasopharynx and irritation of the bronchial mucosa are excessively common. Indeed, if we consider the morbidity of but two diseases, which in-

variably cause a good deal of irritation of the bronchial mucosa and thus of the bronchial glands, namely, measles and pertussis, the frequency of the condition can not be doubted.

SIMPLE ADENITIS.

Simple adenitis of these glands is very frequently only the precursor of the far more serious tubercular involvement. There is no need to quote many figures as to the frequency of such tubercular infection. As long ago as 1865 Steiner and Neuritter showed that in 302 autopsies in tuberculous children, the bronchial lymph nodes were affected 275 times (91 per cent.). And the oft-quoted figures of Northrup, who found the nodes tubercular in every case, in a series of 125 autopsies, irrespective of the cause of death, hardly need comment. It is not only true that tubercular bronchial adenitis is present in at least four-fifths of all cases of tuberculosis in children, but—and this is the more important—it is certain that in the majority of these cases, it is the primary lesion.

There can be no doubt that many cases of enlarged bronchial glands tend to recovery, without subsequent tubercular infection. The danger of such infection, however, is always present, and it is, after all, this factor which adds so greatly to the gravity of the condition.

If we consider that the morbidity of measles is probably greater than that of any other infectious disease; if we add to this the enormous frequency of pertussis, influenza and of such forms of irritation as adenoid vegetations, the statement that almost every child, at some time or other, exhibits a bronchial adenitis, does not seem exaggerated. Of course, in many cases, such a simple adenitis recedes with the disappearance of its cause. With the subsidence of the special infection, the strong and otherwise healthy child recovers, and is to all intents perfectly well again. Yet every adenitis of this sort is but a preparation of the soil whereon, under proper condition of diminished vitality and lowered bodily resistance, the ubiquitous tubercle bacillus may grow and flourish. And other things being equal, the longer such adenitis persists, the greater the danger of tubercular infiltration.

TUBERCULAR ADENITIS.

Yet, even though it be admitted that tubercular disease of these glands is enormously common, it is not to be forgotten that this form of tuberculosis—so long as it remains localized—is still decidedly amenable to treatment. Indeed, tuberculosis, in infancy and childhood, is essentially a disease of lymphatic structures. It may remain so localized for months and years, gradually progressing to other and more incurable forms. There is, it is true, the ever-present danger of general infection; and owing to the anatomic peculiarities, to the peculiarly intimate relation of these glands to other lymphatic systems, such infection is particularly apt to occur in the form under consideration, namely, enlargement of the bronchial glands. A recent writer insists on "the predominating influence of the tracheobronchial glands, as compared with any other lymph glands, in the origination of tubercular processes elsewhere in the body."

DIAGNOSIS OF ENLARGED BRONCHIAL GLANDS.

From the viewpoint of prophylaxis, as well as of direct therapy, it is thus very desirable that we should be able to make a definite diagnosis of enlargement

* Read in the Section on Diseases of Children of the American Medical Association, at the Fifty-fifth Annual Session, June, 1904.