superficially situated, the patient was never dangerously ill, and, indeed, made a good recovery without any permanent pitting.

Several photographs of patients treated with red light, who recovered with no scarring or only insignificant scars, published in Finsen's book, represent what we in this country would regard as mild and favorable cases, the eruption being on the skin rather than in it.

It must be remembered that Denmark, Sweden and Norway are, with the exception of Germany, the best vaccinated countries of Europe. It is illogical and misleading to definitely attribute favorable results to certain measures employed in the treatment of smallpox in persons once vaccinated. The vast majority of such patients recover without scarring under any treatment and even without any treatment. Again, in the absence of severe epidemics, and there have been no severe epidemics in the past decade, young children with variola (who are, of course, unvaccinated) usually escape without permanent scarring.

The character of the prevailing form of smallpox, whether mild or severe, is a matter of great importance in determining both mortality and disfigurement. Dur-

Fig. 3.—Example of a case of smallpox with a semi-confluent but superficial eruption in an unvaccinated mulatto, which was not followed by any permanent scarring.

ing the year ending June 30, 1902, there were in the United States 55,857 cases of smallpox with 1,852 deaths (a mortality rate of 3.31 per cent.), and in the year previous 38,506 cases and 689 deaths (a mortality rate of 1.79 per cent.). What would the verdict have been if red light had been used in these cases?

In the February (1903) number of a popular monthly magazine there appeared several articles on "Niels Finsen and His Healing Rays." Among other unwarranted statements concerning the red-light treatment of smallpox appeared the following: "All the world might now have smallpox without fear of disfigurement."

No one would for a moment endeavor to detract from the hard-earned and well-deserved reputation of Finsen, but in the interests of truth such statements as that just quoted should be challenged.

AUTOMOBILE CONJUNCTIVITIS.—Ophthalmologists report many cases of conjunctivitis caused by rapid riding in automobiles without any protection to the eyes. Davis, in the *Postgraduate*, says it is traumatic, due to friction of the wind and chilling of the eyeball. The treatment is to remove the cause, apply 15 minute cold compresses thrice daily and mild astringent washes by an eye cup twice daily.

THREE ESSENTIAL POINTS IN THE OPERATION FOR CICATRICIAL ECTROPIUM.

F. C. HOTZ, M.D.

Professor of Ophthalmology and Otology in Rush Medical College, CHICAGO.

Some one once said that blepharoplastic operations look very pretty—on paper—but in reality the results are anything but pleasing and satisfactory. The operations for cicatricial ectropium are not excepted from this criticism. The restored lids often look hideous, and still oftener are again everted after a few weeks or months. But we can overcome these drawbacks if we pay strict attention to the following points:

1. The proper division and fixation of the skin flaps.
2. The selection of the most suitable material for covering the lids.
3. The shortening of the overstretched lid border.

THE PROPER DIVISION AND FIXATION OF THE SKIN FLAPS.

The greatest difficulty we have to contend with in the operations for cicatricial ectropium is the tendency of the shrinking skin flaps to evert the lids again. As all transplanted flaps will undergo more or less shrinkage and as the lid border is least capable of offering any resistance to the traction of the shrinking flap, a re-eversion of some degree is almost inevitable as long as the transplanted flap is attached to the non-resisting lid border on the one side and to the non-yielding skin of the forehead or cheek on the other side, because the lid border is then exposed to the full traction force of the shrinking flap. It is evidently much easier for this flap to pull the lower lid down than to draw the skin of the cheek up, and therefore re-eversion is an especially common occurrence after the operation for ectropium of the lower lid.

It is plain, then, that to prevent the reoccurrence of ectropium we must place the lid border beyond the reach of this traction force. And this can be accomplished if instead of covering the whole wound with one skin flap we make use of two flaps, a small one, which is to cover the lid surface only and which we will call the lid flap, and a large one, which is to be spread over the remaining wound area; and furthermore, if we make provision that the contraction of the larger flap can have no effect on the lid flap.

In ectropium of the upper lid this point is gained if the upper edge of the lid flap (Fig. 1 ba) is firmly attached to the upper border *a* of the tarsus, while its lower edge *b* is united with the free margin. The lid flap is thus anchored above and below to the tarsus and its contraction can not turn the lid over because to do so the traction force must have a fixed point of purchase outside of the lid. The shrinkage of the other skin flap *ac*, which has its point of purchase *c* outside of the lid, however, can not disturb the position of the lid margin, because its pulling force is expended entirely on the upper tarsal border *a*; for, thanks to its firm union with this border, the lid flap can not be stretched or drawn upward by the contraction of the other flap, and as long as the lid flap can not be drawn up the lid border is effectually guarded against re-eversion.

In ectropium of the lower lid, the same principle is adopted for the protection of the lid against the vicious traction of the shrinking skin flap. But we must bear in mind that the tarsus of the lower lid is very small and narrow and that normally the lid skin reaches farther down than the lower border of the tarsus, to a slight furrow a little above the infraorbital margin (Arlt's tarso-malar furrow) where the integument passes from the upright plane of the lid into the sloping surface of

the cheek. This normal boundary line between the lower lid and cheek must be re-established by our division and fixation of the skin flaps; and the lid flap *dg*, therefore, must not be united with the lower border of the tarsus, but fastened to the tarso-orbital fascia (at *g*) in a line a little above the infraorbital margin. As the fascia is, as it were, the anatomical continuation of the tarsus, the lid flap adherent to the surface of the tarsus and fastened to the fascia can not evert the lid. And the shrinkage of the large flap *ge* covering the wound of the cheek, is prevented from pulling on the lid margin and causing re-eversion of the lid, because its traction can not reach beyond the firm union of the lid flap with the fascia. So strong is the resistance of this union that you may draw the skin of the cheek down as much as you please and you will not succeed in everting the lid.

THE SELECTION OF THE MOST SUITABLE MATERIAL FOR SKIN FLAPS.

If it is our ambition to obtain the best possible results in ectropium operations—and nothing less ought to satisfy us—we can not be contented with the mere reposi-

Figure 1.

tion of the lids, but should strive also to restore as much as possible their natural appearance. The lids contribute more to the expression of the eye than the eyeball itself; and the slightest alteration of shape and outline, or the least impediment of motion seriously mars the expression of the face. This is a very important point and should always be taken into account when we choose the material for our flaps. The normal integument of the lids is a thin and delicate skin of such perfect flexibility and adaptability that it allows perfect freedom of motion and accommodates itself readily to the numerous slight changes in contour so characteristic for the various facial expressions. In order, therefore, to gain a perfect cosmetic result we must select as a substitute for the lost lid skin a material which possesses the same qualities; it must be thin, light and adaptable.

It goes without saying that the thick skin of the forehead, temples or cheek does not possess these qualities, and therefore flaps from these regions should not be transplanted on the lids. Besides the transplantation inflicts additional wounds on the face, and I hold we should avoid marring the face by unnecessary scars. The use

of Wolfe's flaps taken from the arm avoids this latter objection, but they, too, are usually so thick that, like the pedunculated flaps, they make a heavy, clumsy-looking lid. These two kinds of flaps are as well suited for a substitute of lid skin as shoe leather would be for kid gloves.

A good and suitable material for lid flaps which answers all the requirements is the cicatricial skin usually found in the immediate vicinity of extensive ectropium. I have shown in 1896¹ that this skin can be successfully transplanted and makes a perfect lid skin. In ectropium of the lower lid this cicatricial skin can nearly always be utilized for the lid flap. In ectropium of the upper lid it can sometimes be used when the eyebrows are absent. If on account of the eyebrows the cicatricial skin is not available, a Thiersch graft is the only suitable material. It accommodates itself to the surface of the lid, brings out its contours perfectly and does not in the least interfere with the free movements of the lid.

THE SHORTENING OF THE ELONGATED BORDER OF THE LOWER LID.

As the diameter is shorter than the half circle, so in the case of complete ectropium the free lid margin of the lower lid turned from its almost straight line between the canthi into a long downward curve is elongated a good deal; and after being held in this abnormal state for a

Figure 2.

long while it will not recover its normal length when the lid is replaced. The elongated margin of the lower lid will not closely fit the curvature of the eyeball and will drop away from it, as in senile ectropium. The reposition of the everted lower lid, therefore, can not be perfect and permanent, unless the overstretched lid margin is reduced to its proper length. In several instances I did not do that, because the lid margin appeared to lie in perfect apposition to the globe; but every time I had cause to regret this omission, and had to correct my mistake by a second operation. I therefore regard the shortening of the elongated border of the lower lid a very essential point in the operation for cicatricial ectropium.

And now, after having so strongly emphasized the importance of these several points, I wish to indicate briefly the plan of operation based on the views expressed in the foregoing remarks. As the operation is not alike in details for both lids, I believe it is better for a clear understanding to describe the method for each lid separately.

TECHNIC OF THE OPERATION ON THE UPPER LID.

If the eyebrows are partly absent and there is a good expanse of cicatricial skin above the everted lid, we cut from this cicatricial skin the lid flap in the following manner:²

1. THE JOURNAL A. M. A., Sept. 19, 1896; and Archives of Ophthalm., vol. xxv, No. 3.

2. Archives of Ophthalm., vol. xxv.

"From a point (*a* Fig. 2) about 5 millimeters above the inner canthus an incision is made obliquely upwards into the cicatricial skin and then continued in a curve downwards to a point *c* about 5 millimeters from the external canthus. This incision outlines a large flap *abc* which is carefully dissected up from the underlying scar tissue as far as the lid borders. The lid is then released by dissection from all cicatricial connections until it can be easily turned down into its normal position; and now the edge (Fig. 3 *ac*) of the lid flap is fastened by silk sutures to the upper border of the tarsus."

If, however, on account of the eyebrows the lid flap can not be taken from the cicatricial skin, we make an

Figure 3.

incision along the lid border and after the reposition of the lid cut from the arm a Thiersch graft of suitable size, which is transported on the razor directly to the lid, spread out over its surface and fastened by fine silk sutures to the upper tarsal border as well as to the wound edge of the free border. These sutures must be inserted with great care to make sure the edges of the flap do not roll in; their application therefore is very tedious work; but the time is well spent, because the graft thus fixed

Figure 4.

is positively secured against being shifted from its place by any movement of the lid or by any accidental displacement of the dressings.

After the lid flap is fastened (as in Fig. 3) the lid is drawn down as far as possible and held in this position by two ligatures passed through the free border and fixed on the cheek by plaster strips. This is done for the purpose of immobilizing the lid during the healing process and also for enlarging the wound (*abc*) above the lid to its fullest capacity. Over this wound a Thiersch graft is spread out so that its edges lap over the surrounding skin; no sutures are used.

TECHNIC OF THE OPERATION ON THE LOWER LID.

Here the lid flap can always be procured from the cicatricial skin; but as this skin shrinks considerably as soon as it is dissected up we must take good care to cut the flap of very liberal dimensions. We begin the incision one centimeter below the inner canthus (Fig. 4 *g*), carry it obliquely down into the cheek to a point *m* 2 to 3 centimeters below the center of the everted lid margin; then we continue it in an oblique direction upwards and outwards to a point *b* even with and one centimeter from the outer canthus. This large flap *gmb* is then dissected up from the underlying scar tissue and all cicatricial strands and bands are cut until the lid is freed and can be turned up.

The next step is to reduce the overstretched lid margin to its proper length by removing a suitable piece of the lid (except the conjunctiva) near the outer canthus by the following incisions (Fig. 4): from the canthus *a* transversely to *b*; from *a* to *c* along the lid margin, and from *c* to *d* through the flap; the edges *cd* and *ab* are then united by two silk sutures. Now the lid is drawn up as far as possible

Figure 5.

and held in this position by two silk ligatures passed through the free margin and fastened on the forehead by adhesive plaster or collodion. This done the edge of the lid flap is anchored to the tarso-orbital fascia by silk sutures (Fig. 5). The lid flap should be evenly spread out, so as to be in perfect contact with the wound surface; but we must carefully avoid any undue stretching; and should we find that the flap is a trifle short and would be stretched if the sutures are placed very near the infra-orbital margin, it is better to put them through the fascia 1 or 2 millimeters higher up. Finally, the wound surface *gmb* below is covered with a Thiersch graft, the edges of which are made to lap over the surrounding skin. No sutures.

AFTER TREATMENT.

The after treatment is the same for the upper and lower lid. Strips of gutta percha protective are laid over the flaps, and on these strips a gauze compress wrung out of warm boric acid solution; this is covered by a large

square piece of protective to prevent evaporation and over this is put a layer of cotton to maintain uniform warmth. This whole dressing is secured in place by a roller bandage and best left undisturbed for three days. Then it is carefully removed; the flaps are thoroughly cleansed, the overlapping edges of the large flap are trimmed off and a new dressing is put on which is changed every day or two, as circumstances may require. At the end of the first week the ligatures and the sutures of the lid flap may be removed; during the second week some simple ointment (like borated vaseline) is applied daily on the flaps, and after this period no further treatment is required.

THE OBJECTIONS TO PRESCRIBING MEDICINES OF UNKNOWN COMPOSITION.

AUGUSTUS A. ESHNER, M.D.

PHILADELPHIA.

If the proposition contained in the foregoing terms were always presented so clearly and so directly there would be little room for discussion, as I think it will be generally conceded that the fact of the composition of a medicine being unknown would constitute in itself the most serious objection to its prescription by physicians. Nevertheless, I fear that many prescriptions are written daily for preparations concerning whose composition the prescriber is not in possession of such fulness of knowledge as is requisite for the most intelligent and most successful employment of the articles of the *materia medica*.

The modern manufacturing pharmacist has exhibited marked skill and ingenuity, worthy of a better cause, in concealing ignorance behind the mask of false knowledge. It has become the vogue for this enterprising individual, firm or corporation, to put forth products to which he gives fanciful names, for which he claims elaborately described marvelous effects, and of which he states the composition in accordance with his own views and purposes. He has a predilection for describing his product as a synthetic preparation with a formula comprising C, H, N, and O in startling proportions, and he gives it an almost interminable and unpronounceable chemical name, but is good enough to coin a descriptive and copyrighted substitute. Sometimes it is, at other times it is not, a coal-tar product, according to personal caprice. If the preparation be a mixture, only its constituents may be named, and sometimes not all of these, the proportions being omitted. In these and devious other ways a pretence of publicity is made, by which the careless and indifferent may be readily if not willingly deceived. Samples are distributed generously to physicians, who read with avidity the accompanying so-called literature skillfully devised by the manufacturer, and it is not long before testimonials indorsing the new preparation are forthcoming, and clinical reports of cases treated with the new preparation appear in certain medical journals. Sometimes the aid of the physician is further invoked by securing him as a stockholder or officer in the manufacturing concern, and his interests may be further looked after in the way of commissions or concessions in return for prescribing the remedy. The preparation is now launched on a career of success and prosperity, which may be of longer or shorter duration in accordance with a variety of conditions. Mayhap when the profession has done all in its power to contribute to this end, direct appeals to the laity, supported by the indorsement of "the medical faculty," are made

to the public through the advertising columns of the newspapers and the dead walls and spaces of the city and the country.

The practice of medicine, though art, is based on scientific principles, and it will be successful in proportion to the fulness and precision of the knowledge underlying it. If the identity of a medicine is not known through its composition the prescriber has no assurance that he will always receive the same article, but, on the contrary, there is good reason for believing that he will get something different, in quantity or in quality, at different times. In this way he is liable to do discredit to his art, and injury, positive or negative, to his patient. Additionally, injustice is done the patient in that he is compelled to pay a higher price for medicines of concealed or suppressed composition than he would if these were prescribed in the regular way. The practitioner who permits himself to fall into the habit of prescribing ready-made preparations surrenders thereby originality and independence in his therapeutics, and fails to make progress in his art. The manufacture and the use of secret preparations have also a blighting effect on legitimate pharmacy. Finally, after all has been said on both sides of the question, if there are two sides, there remains the practical argument that the prescription of medicines of unknown composition is unnecessary, as all that can be accomplished with them, barring the potent influence of secrecy, can be brought about as well if not better by means of medicines of known composition.

From the foregoing considerations it must be clear that the prescription of medicines of unknown or concealed composition is unscientific, unprofessional, unfair if not prejudicial to the sick, unprogressive and unstimulating and, most of all, unnecessary.

The remedy for the existing state of affairs is in the hands of the medical profession. The manufacturing pharmacists will not make preparations that they can not sell, and if physicians will not permit the creation of an artificial demand for illegitimate preparations, and will cease to prescribe them, the manufacturers will soon discontinue their production. To this end it is the duty of all medical men having the interests of their profession and their patients at heart to confine themselves to the employment only of such drugs or combinations of drugs whose composition is of public knowledge, and further, is vouched for by some recognized authority, official or otherwise, such as the Pharmacopeia, the Dispensatory and the Formulary; or, in lieu of such indorsement, in the intervals between successive revisions, is given authentic publicity by investigators of acknowledged standing and competency through legitimate and recognized channels, such as the professional journals and independent publications. Above all things, the so-called literature of the manufacturing pharmacist and the reprints of articles from publications of questionable character should be viewed with suspicion.

HABITS.—A cheerful habit of mind is like a fountain of youth to the body. Worry kills as surely as does arsenic, but steady, moderate work is a prophylactic against many of the ills to which flesh is heir. The greatest brain workers have generally been long-lived (as witness Gladstone and Bismarck), provided that they did not neglect the laws of hygiene and the natural demands of the body. The physician's irregular hours for eating and sleeping tend to disorder his faculties, depress his vitality and tempt to the use of drugs whose fatal allurements he knows best of all.—*Denver Medical Times*.