Under appropriate treatment, the colicky pains disappeared, and the patient began to improve in appearance; but thirteen days after the seizure which had attacked him the day after admission he began to be troubled with cough. This continued to increase, and the following note was made on Nov. 17th:—Lower lobes of both lungs rather dull; the right, bubbling crepitation; the left, tubular breathing; and at extreme base some crepitation; expectoration dark-coloured, with fetid stench; pulse 100. Temperature: Morning, 101·4° Fahr.; evening, 104·1°. On Nov. 20th, he had some decided hæmoptysis, and exhaustion rapidly set in. He died at 1.15 A.M. the next day.

Post-mortem examination.—Pericardium contained three ounces of clear serum. Heart weighed ten and a half ounces; no valvular disease, but a few small spots of atheroma at commencement of aorta. Pleuræ firmly adherent, especially the left. Right lung weighed twenty-five ounces; very pale; posterior part of lower lobe quite solid. Left lung weighed twenty-one ounces; greater part quite solid, except apex. At the posterior and outer part, a cavity about four inches long; the outer wall formed of thickened pleura, the other wall shreddy, brownish-green, softened, and gangrenous. The dead tissue formed an isolated, abruptly limited, gangrenous patch, about the size of a walnut, separated from the surrounding pneumonia by a yellowish membranous deposit of lymph. Brain not examined.

Remarks.—Dr. Ralfe observed that the gangrene of the lung in this case was, without doubt, proximately due to embolism, and that the mischief might be dated from the day after the admission of the patient to the hospital, when he became insensible; that attack being due, no doubt, to the simultaneous plugging of one of the arteries of the brain with an embolus. The inflammation excited in the lungs by the embolus was, at first, apparently of a very mild character, for it was not till eleven days after the first appearance of lung trouble that any suspicion of gangrene existing was entertained. From this slowness of onset, and also from the fact that the man had suffered from the effects of syphilitic, malarial, and metallic poisoning, it would appear as if the gangrene in this case resulted rather from a depraved textural condition of the lung itself, which rendered it unable to resist an attack of inflammation of moderate intensity, than from the actual cutting off of the blood-supply by the embolus.

ABERDEEN ROYAL INFIRMARY.

CASE OF PSOAS ABSCESS; ANTISEPTIC TREATMENT; RECOVERY.

(Under the care of Dr. OGILVIE WILL.)

J. N—, aged fourteen, suffering from angular curvature of the spine, involving the seven lower dorsal and first lumbar vertebræ, of long duration, was admitted to the infirmary on account of a psoas abscess pointing in the left groin, of which the following is a condensed history. He stated that about six months before admission, when taking violent exercise, he “felt something give way.” About a month afterwards he observed a small swelling in his groin, which gradually increased, but was unattended by pain. He applied to Dr. Garden, who aspired the swelling on three occasions, drawing off a small quantity of fluid at each tapping, but failing to perceptibly reduce its size. As aspiration was evidently insufficient, he was sent into Jacob’s ward for the purpose of having the abscess opened antiseptically. At that time a large, tense, fluctuating swelling was observed on the inner side of his left thigh directly under Poupart’s ligament, presenting the usual features of a psoas abscess.

On August 17th, 1877, the skin having been thoroughly purified by washing with carbolic lotion, the abscess was opened under spray, and a large quantity of pus, containing numerous flakes of curdy material, was allowed exit. Digital pressure was employed to squeeze out the remaining fluid, and several pieces of bone, evidently derived from the bodies of the affected vertebræ, were removed by passing a forefinger into the cavity. A long drainage-tube, of large calibre, was introduced, and the usual antiseptic dressings were applied, an elastic-web spica bandage holding all in position. It may be here mentioned that, had it not been for this elastic bandage, it would have been perfectly impossible to prevent the access of putrefactive elements to the wound, for the boy felt so well from beginning to end,

and he was, as Mr. Glass (one of the dressers of the case) truly remarked, so “full of life,” that he could not be kept still; he was so active in his movements that fifty times each day he ran the risk of displacing the dressings, but by the aid of the bandage, and a Liston splint, which it was subsequently found necessary to apply to restrain him, the danger was averted.

August 18th.—Dressings removed under spray, a small quantity of red, serous discharge staining the deep dressing. Temperature 99·4°; pulse 92. Dressing as before.

19th.—No sign of inflammation about wound. Tube filled with curdy material. A very little serous fluid obtained by pressure. Temperature 99·2°; pulse 96.

21st.—Very slight serous discharge. Curdy substance in tube. No inflammation. Temperature 98·8°; pulse 84.

23rd.—Discharge similar in character, but less in quantity. Temperature 98·6°; pulse 72.

Sept. 5th.—A smaller tube was substituted for that first inserted. From this date progress towards recovery was steady, the cavity gradually contracting, necessitating the lessening of the length and calibre of the drainage-tube from time to time; but as the notations, which were carefully taken on each occasion when the dressings were changed by Mr. Aitken, are similar to those given, repetition is uncalled for.

On Sept. 30th there was a sudden increase of discharge, the fluid being light-yellow in colour, mixed with curdy matter, but there were no signs of inflammation about the opening, no tenderness on pressure, and no putrefactive odour. At this date the temperature, which had been slightly higher than usual for some days previous, ran up to 102·4°; pulse 100. On the following day it fell to 100·9°, and the dressings were not removed. On Oct. 4th it was again normal. During the next four weeks the discharge varied in quantity, but the abscess cavity gradually became obliterated. On Nov. 5th, on the wound being exposed, after the dressings had remained unchanged for three days, only a slight stain on the gauze directly in contact with the opening was found. A very narrow drainage-tube, one inch long, was inserted with difficulty. On Nov. 12th it was shortened to half an inch; and on Nov. 17th it was removed altogether, after which the opening speedily healed, and the patient was dismissed from the hospital. He presented himself a short time afterwards in perfect health, and since his dismissal he has been disporting himself in a most active manner without showing the slightest inconvenience.

CARDIFF UNION HOSPITAL.

INTESTINAL OBSTRUCTION.

(Under the care of Dr. SHEEN.)

A. B—, aged eighteen, a pawnbroker’s assistant, was admitted at noon on January 21st. He was a Jew, had a strumous look, a red-and-white complexion, and dark hair. He stated that he had not passed wind or had a motion since the 18th, that his bowels had been regular up to that time, that he had a similar attack twelve months ago which lasted two weeks, cause unknown, and that he had been subject to such seizures since childhood. Up to the 18th he was quite well, and had had his bowels moved daily and naturally. In the morning he was roused from sleep by a pain round the umbilicus, which went away in half an hour, but soon returned, and got worse towards evening, when he went to bed and took two pills, sent him by the doctor, after which he slept, and the pain was easier. A medical man saw him on the morning of the 19th, and from that time to his admission he was treated chiefly by enemata and poultices.

On admission there was uniform enlargement of the abdomen, with tympanites and pain, occasional hiccough, and an anxious expression. A turpentine stupe was applied to abdomen, and an enema of turpentine and castor oil was administered. Fifteen drops of tincture of belladonna in half an ounce of water were given every two hours.

On the 22nd the abdomen was immensely distended and tympanitic throughout. There was no relief, and the anxious look continued. He had been sick once or twice during the night, but not afterwards. The pulse was fairly good; pupils not dilated, and the throat not dry. Treatment continued. A warm-water injection to one pint was given. After 4 P.M. the bowels acted four or five times, and several