plantation, the older methods are usually successful; but the cases of the central variety, where the cervix does not easily dilate, and in which the hemorrhage is profuse and uncontrollable then Cesarean section, in my opinion, is not only a justifiable procedure but is imperatively demanded.

THE CHEMICAL AND MICROSCOPIC EXAMINATION OF BLOOD.*

W. D. KELLY, M.D.
ST. PAUL, MINN.

Until recently little has been known of the blood as a factor in the cause of disease. The microscopic morphologic changes which take place in the blood, and their cell identity, have quite definitely been fixed or determined; however, there has not been much progress made from a chemical standpoint in blood examination. Recently, Schmidt, Hoppe-Seyler, Judell and others have contributed liberally to the chemistry of the blood, and among other things they have given us chemical blood analysis, and a working foundation whereby we are enabled to forward chemical analyses of the blood with more precision. The blood in the body is enclosed, so to speak, in a sealed tube, and is unable to gain exit, except by osmosis, or through solution of continuity in the blood-vessel. The depletion of the red blood normally takes place, principally, in the spleen, and it receives its supply from the lymphatics and red bone marrow.

The blood is composed of plasma, serum and blood-plates. Intermixed in the blood, gases are found—carbonic, oxid and oxygen. Salts are also found—magnesium, calcium, potassium and sodium. To these salts are accredited, by many writers, the alkalinescence of the blood. In the blood are found red and white blood-corpuscles. The red corpuscles are composed of oxy-hemoglobin and neucloproteids, lecithin and chloesterin.

In the leucocytes have been demonstrated fatty acids, lecithin, glycogen and salts, similar to those found in red cells. Blood-plates are regarded by Lilienfeld as belonging to the neuclo-albumins, and are supposed by him to be the remnant of the leucocytic neuclei.

The blood serum, containing serum-albumin and serum-globulin, is subject to chemical change in puerperal septicemia, and in febrile disease; serum-globulin is less subject to change than serum-albumin.

The inorganic salts of the serum are phosphates, chlorids, carbonates and sulphates. The pathologic variations in the phosphates of the serum vary in slight degree and importance. The variations in the chlorids of the serum, investigated by Schmitt, Barnicke, and Limbeck, have not been found extreme nor of notable pathologic import, although this principle is chiefly responsible for the isotonic relations of cells in serum. A high percentage of chlorid is usual in anemia. The larger the proportion of plasma, the greater is the percentage of chlorids in the blood. The alkalies, sodium and potassium, are principally combined with chlorin in the blood. Large proportions of phosphates and carbonates of sodium and potassium are found. Sodium salts are principally found in the plasma, being usually increased in the watery blood. Potassium, found in red cells, is diminished in hydremic conditions.

Sodium carbonate in the plasma is responsible for the alkalinity and the power of plasma to produce CO_2 . The iron in the blood is principally found in hemo-

globin; traces are found in plasma and neuclein. Seegen, Chaureau, Cavazanni, von Mering, Limbeck, found in normal blood traces of glucose, which is increased by a diet of carbohydrates and diminished by muscular exertions. The diastase ferment of blood is found in the red cells and serum, and is found, by Castellino, Tiegel, Plosz, and Precka, to have the power of coagulating blood. It is inhibited by the neuclein and increased by sodium sulphate and chlorid.

After a heavy meal, fat has been demonstrated in the blood, by Gunprecht. The occurrence of free fat, palmitin, stearin and olein in the blood in health and disease has repeatedly been observed. It has been found very much increased in healthy individuals after a hearty meal. Acetone is found in fevers, Jksch has demonstrated fatty acids in the blood of leukemia, acute atrophy of the liver and infectious diseases.

CHOLEMIA.

The poisonous symptoms developed in cholemia have been referred, by most authorities, to the presence of biliary acids. To the naked eye, icteric blood may appear yellowish-red in color in the serum or its foam. Small quantities of bile pigment have been readily detected by a peculiar yellowish tint. On rapid heating, the yellowish-red bilirubin changes to green biliverdin. Diminished isotonic tension and increased resistance of the red cells are peculiar characteristics of the red blood in jaundice. Limbeck found tension of the cells reduced .40, .38, and .32 per cent. sodium chlorid. The well-attested fact that in intense jaundice red cells are frequently dissolved by biliary principles, can with some difficulty be reconciled with that markedly hyper-isotonic quality of the serum. Bile acids affect the union of hemoglobin with the stroma of the red cells, rendering hemoglobin more easily soluble; this accounts for the solution of red cells in jaundice, as well as in other conditions. Other characteristics of icteric blood are increase of nitrogenous bodies, diminution of chlorids of both blood and serum, which is displaced by biliary acids, and well-marked increase in the volume of red cells. Well-marked cholemia may be detected by inspection of serum or foam on heating to 50 C. Bilirubin may be changed to biliverdin by this process.

By the intravenous injection of glacial acetic acid in the rabbit, I have been able to get an acid reaction several hours afterward by the phenol-phthalein test. It simulates that found in diabetic cases. Singular to say, I have also found that a rabbit weighing 1550 kilos can, apparently without any inconvenience, tolerate as high as a 2-gram intravenous injection of glacial acetic acid into the vein. Examination of the blood shows decrease in red cells and hemoglobin, also decreased leucocytes.

The specific gravity of the blood varies by the amount of water present and amount of salts contained. The conclusion reached by many authors has placed the specific gravity from 1.058 to 1.062 in man; 1.054 to 1.060 including both sexes. The specific gravity of the blood may be increased by sweating, lack of food, muscular exertion, and may be decreased by freely imbibing in water or fluids.

Osmosis is the force which leads to the interchange of blood through the animal membrane, until the salt in each is equal. When two fluids have equal content of salt, they are said to be isotonic. The isotonic tension of red cells is .46 per cent. sodium chlorid. Homberger finds that albumins, phosphates and chlorids behave differently after changing osmotic conditions. When a lit-

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Practice of Medicine, and approved for publication by the Executive Committee of the Section: Frank Billings, George Dock and J. M. Anders.

the acid is added to blood, albumin and phosphates pass from the red cells to the serum, while chlorids pass from the serum to the cells, but when an alkali is added the opposite transfer is induced.

Similar physical effects are produced by the passage of oxygen and carbonic acid through the blood. Homburger suggests that these factors take an important part in the metabolic exchanges in the capillaries.

It is an established physiologic principle that the absorbent power with relation to carbonic dioxid, entirely depends on its alkaliescence; venous blood contains more carbonic dioxid than arterial.

Febrile processes are generally accompanied by diminution of CO_2 in the blood. Often in proportion to the height of the fever this condition, according to Gepherd, and Minkowski, is referable for the abnormal production of acid metabolic principles in the blood. In the cachexia of carcinoma, Limbeck and F. Klemperer found marked diminution of the CO_2 in the blood, but did not estimate the acid principles.

The blood is frequently very deficient in CO_2 . There are certain unsaturated salts in the blood,—sodium (Na_2CO_3 ; NaH_2PO_4 ; Na_2HPO_4)—which, although alkaline to litmus are acid to phenol-phthalein and are capable of uniting with bases. When fresh blood is alkaline serum reacts as acid phenol-phthalein. The capacity of the blood salts to neutralize bases has been called its basic capacity, by Krous. In normal venous blood Krous found the capacity to be .162 to .232 per cent. NaOH , which increase in febrile conditions to .209 to .272 per cent.; this is a demonstration of the marked increase of basic capacity in diabetes and has been regarded as strong evidence of acid intoxication in this disease.

By increasing doses of phenylhydrazin it is possible to reduce the red blood count to one million corpuscles, or even to a greater degree.

EXAMINATION OF THE BLOOD.

Five things are essential in order to make satisfactory examination of the blood: 1. The apparatus must be absolutely clean. 2. The various stages in the process must be performed rapidly, because the cell coagulation of the blood will interfere with any of the tests. 3. The work must be done accurately. 4. Making large quantities of stain and keeping some in glass-stoppered bottle will standardize the solution, so one will receive minimum variations in intensity of stain. 5. Fixing of specimen, by continuous heat, with as slight a degree of variations in distribution of heat as possible; taking of blood specimens with reference to time of day.

In the healthy individual, with the above-mentioned methods, I have noted increase and decrease of the leucocytic count, varying from 1000 to 7000 per cubic millimeter in the course of a day.

In order to more accurately fix the blood count, I have found that specimens taken in the morning, before any undue excitement is indulged in and previous to taking of fluids, or solids, will give a more uniform blood count than those taken at any other period of the day.

A subcutaneous injection of quinin in the forearm in a case under treatment for malarial fever at the hospital at Nantes, France, resulted in paralysis of the thumb and forefinger, incapacitating the patient for work for several months. He sued the hospital, the physician in charge and the interne who made the injection, claiming that the latter had failed to obey instructions to make the injection in the thigh. The courts exonerated the defendants and dismissed the case with costs.

LEUCOCYTE COUNTS IN HEMORRHAGE.*

GEORGE DOUGLAS HEAD, M.D.

INSTRUCTOR IN PATHOLOGY, UNIVERSITY OF MINNESOTA.
MINNEAPOLIS, MINN.

It has been conclusively demonstrated that the loss of blood in man and the higher animals is followed by an increase in the number of leucocytes in the circulating blood. This increase in white cells following hemorrhage has been called post-hemorrhagic leucocytosis and ranges all the way from 11,000 to 30,000 white cells per cubic millimeter.

How soon after the hemorrhage the leucocytosis appears does not seem to have been determined with absolute certainty. Cabot, in his "Clinical Examination of the Blood" (page 111), makes this statement: "Immediately after a loss of blood, we can usually find a decided leucocytosis, despite the dilution of the blood." On page 90 of the same work he writes: "Within an hour after a large hemorrhage we commonly find a considerable increase, 16,000 to 18,000 cells."

The size of the hemorrhage required to produce a leucocytosis and the length of time during which the increase of white cells continues after the hemorrhage has stopped are questions upon which meager information is offered. In two cases of cerebral hemorrhage recorded by Cabot (page 111), the loss of blood was not great enough to affect the red cells, but the white cells were much increased. In one a count of 25,000, in the other a count of 15,000 white cells was made.

On page 91 of Cabot's work appear these words: "In hemorrhage from the stomach, the leucocytosis disappears in a day or two, while in traumatic hemorrhage it persists longer."

The statements above quoted sum up in a general way all the definite knowledge we possess on the leucocyte count in hemorrhage. Much, if not all, of the data upon which these statements are founded has been gathered from counts upon clinical cases in human beings, where other factors beside the hemorrhage per se may have played an important part in the rise or fall of the white cells and where the leucocyte counts must of necessity have been made sometime after the occurrence of the hemorrhage.

I wish to place on record some experimental work on the leucocytes in hemorrhage, which excludes these disturbing factors and gives us what appears to be definite results somewhat at variance with generally accepted views. The experiments were performed on dogs, and I am indebted to Dr. R. O. Beard for his kindness in permitting me the use of material upon which he was experimenting.

The normal leucocyte count in dogs is about that of man, 6500 to 9000. The same laws seem to govern the increase or decrease of leucocytes of the circulating blood in dogs and in men, so far as the writer has been able to follow them. There is a leucocytosis of digestion and a leucocytosis following septic infections in dogs as well as in human beings. The varieties of leucocytes in the circulating blood of dogs are essentially the same as those in man. In all probability the conclusions arrived at from these experiments in dogs would apply equally well to human beings.

EFFECT OF HEMORRHAGE ON THE LEUCOCYTES.

The first series of experiments deals with the effect of hemorrhage on the number of leucocytes in the cir-

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Practice of Medicine, and approved for publication by the Executive Committee of the Section: Frank Billings, George Dock and J. M. Anders.