mentre il signor Yendell (Astronomical Journal, No. 217) troverebbe un massimo di 6^m.4 soltanto, al 3 di Gennajo. Io credo che il notevole disaccordo fra le nostre osservazioni provenga dalla variabilità di una delle due stelle DM. +19°1110 e 1126, e dalla diversa scala fotometrica

Torino 1890 Maggio 24.

adottata per le stelle di comparazione; e per vero se io calcolo le mie osservazioni colle grandezze della DM., in media inferiori per queste stelle di 0^m.32 a quelle dell' Harvard Photometry, trovo numeri più discordanti fra loro, ma più concordanti con quelli del collega americano.

F. Porro.

The Supposed Variable in RA. = 12^h18^m.0 Decl. = -48°43' (1875.0).

The only star above the 10th magnitude lying between Stone 6875 and 6906 is a dull red one whose mean place for 1890.0 has this evening been determined by me as follows from five equatorial filar-micrometer comparisons with each of those stars:

Comparisons with Stone 6875	RA. = 12 ^h 18 ^m 40 ^s .03	N.P.D. = 138°49'50".0
» » » 6906	12 18 40.00	138 49 47.8
Mean:	RA. = 12 18 40.02	N.P.D. = 138 49 48.9

The mean places of the comparison stars have been deduced from the Cape Catalogue for 1880. I am unable to refer to the Argentine General Catalogue, but adopting 7.0 and 6.5 as the magnitudes of 6875 and 6906 respectively I should regard the red star as of the 7.2 magnitude. The star will be watched here in order to a determination of the limits of its variability.

Windsor, N. S. Wales, 1890 May 30.

John Tebbutt.

Measures of Double Stars.

A. G. C. 1. Sirius.

1890.252	359°6	4".17
269	361.6	4.20
304	358.6	4.19
1890.27	359.7	4.19

These measures are subject to more than the usual uncertainty from the fact that they were necessarily made several months after the usual time of measuring this pair, and when it was long past meridian. The companion was seen with great difficulty.

β 101. 9 Argus.

1890.249	88°3	0".28
252	84.8	0.29
255	83.0	0.31
269	82.4	0.34
1890.26	84.6	0.31

The motion since last year is 8". The angular change since the discovery of this pair in 1875 amounts to 155".

A. C. 5. 8 Sextantis.

1890.252	114°6	0".25
269	113.9	0.36
280	111.2	0.30
301	116.8	0.39
1890.27	114.1	0.32

The angle of this interesting binary is rapidly decreasing, the change since the measures of last year being 11". The first measures by Dawes in 1854-60 apparently require a correction of 180° in the angles, and in that case the angular movement has been something over 100".

Mt. Hamilton 1890 May 3.

There are but few measures. I found it sensibly single in 1879-81 with the 18½ inch at Chicago.

Regulus.

B and C (Winlock).

1890.255	86°0	3".01
269	86.8	3.36
282	84.2	2.78
1890.27	85.7	3.05

There is not much relative change in the double companion. They are evidently connected with the large star, since they have the same proper motion, and probably form one vast system.

β 1077. α Ursae Majoris.

1890.249	317°4	0".87
252	322.9	0.83
255	321.3	0.79
269	318.6	1.00
1890.26	320.1	0.87

The measures of last year gave 326".1 and 0".91.

β 1082. 78 Ursae Majoris.

1890.252	77°4	1".35
255	78.1	1.43
280	75.1	1.61
1890.26	76.9	1.46

The foregoing are substantially all the measures I have had an opportunity to make with the large telescope since the conclusion of my last series in September, 1889.

S. W. Burnham.