The case reported by Dr. Phillips is one of the severe types and presents several interesting features which were not present in the other cases described, yet in general it followed the same course.

It deserves mention, as a matter of no small interest, that the clinical course of this local disease coincides in all the essential points with systemic or general infection, and that the duration and departure of the clinical conditions are in harmony with all the vital phenomena manifested by the influenza organism. Thus, the life cycle of the bacillus is short, a culture thrives but a few days, the resistance to harmful agencies is feeble, and under the most favorable surroundings the organism soon dies. Herein, perhaps, we find an explanation for the rapid recovery of the disordered eye when treated antiseptically. In view of what has been observed it appears that we have to do with a pathologic condition which differs materially from other acute inflammations of the eye. And in view of its abrupt and somewhat violent onset, its rapidity of development, its shortness of duration, and in view of the repeated findings of the bacillus of influenza, it would seem that we are justified in making a diagnosis of influenza ophthalmia.

In the *New York Medical Journal* of Feb. 1, 1902, appears a translation of an extract of a paper published by Dr. F. Rymowitch in the August, 1901, number of *Roussky Archiv Pathologiyi, Klinicheskoy Meditsinyi Bacteriologiyi*. It appears that this observer has recently made a comparative study of the *B. influenza* and the bacillus of Koch-Weeks, and concludes they are one and the same organism. He finds that they are identical in morphologic, biologic and pathogenic characters. Both produced intoxication without a general infection. The most conscientious study did not reveal to him any differences between the bacteria, and he therefore believes that the bacillus of acute contagious conjunctivitis and that of influenza are the same. When attacking the mucous membranes of the respiratory tract the influenza bacillus produces influenza, and when affecting the mucus of the eyes produces acute contagious conjunctivitis, which disorder, according to Rymowitch, is the same as that produced by the Koch-Weeks bacillus. Not having had an opportunity to make a comparative study of these two organisms, I am not in a position to confirm or refute the observations made by Dr. Rymowitch. The subject is an interesting one, and because of its importance deserves further consideration, which I hope to give it in the near future.

THE FLY AS A CARRIER OF TYPHOID.

AN INQUIRY INTO THE PART PLAYED BY THE COMMON HOUSE FLY IN THE RECENT EPIDEMIC OF TYPHOID FEVER IN CHICAGO.

ALICE HAMILTON, M.D.

From the Laboratory of the Memorial Institute for Infectious Diseases.
CHICAGO.

During July, August and September of 1902 there was an unusually severe epidemic of typhoid fever in Chicago, which raised the death rate to 402 from this disease alone, as against 212 during the same three months of the previous year.

CHICAGO'S WATER SUPPLY.

The water supply of Chicago comes from the lake and is collected at five pumping stations: the Hyde Park station, which is two miles from the shore; the Fourteenth Street, four miles from shore; the Carter Harri-

son, two and a half miles; the Chicago Avenue, two miles, and the Lake View, two miles out. The Carter Harrison station is considered the best, the Fourteenth Street is almost as good. Specimens of water from each station are examined daily in the laboratory of the Chicago Board of Health and a bulletin as to the condition of the water at each is published in the daily papers. The state of the water varies greatly from day to day, and the practical value of these daily bulletins becomes somewhat problematical when one considers that twenty-four hours must always elapse between the collecting of the water and the published report, so that people who read on Tuesday morning that the water at a certain station is bad and take precautions accordingly are a full day late, for they were drinking the bad water on Monday. On the whole, such a statement as is made at the end of the week by the health department, "the water at this station averaged 60 per cent. good for the week," is really misleading, since nobody can ever know until too late which days made up the 40 per cent. bad. As a usual thing the water is worst when the rainfall is greatest, the sewage being carried farther out into the lake and contaminating the water at the intakes, except when there is the fortunate coincidence of light easterly winds, which, without stirring up the bottom, keep the sewage in toward the shore.

In discussing the causes of this greatly increased mortality from typhoid fever, Dr. A. R. Reynolds,¹ commissioner of health, speaks as follows:

"At the time" (the early summer) "there was nothing in the condition of the water to cause alarm. Notwithstanding the heavy rainfall of the previous months, the supply from all sources averaged 72 per cent. good (by 'good' in the laboratory characterization is meant a water free from pathogenic organisms, the colon group). But in the early part of August the typhoid deaths suddenly jumped up from a previous weekly average of 8.8 to 28 in the first week of August, to 44 in the second, and to 53 and 51 in the third and fourth weeks, with a total of 194 for the month, as against a total of 44 for July. As already stated, the water supply in July and previously was not of such a character as to account sufficiently for the increase of typhoid; but a study of the meteorologic conditions of the period beginning in October, 1901, reveals what are believed to be strong contributory causes. October, November and December, 1901, and January and February, 1902, were the driest five consecutive months on record in 53 years. During the period only 6 inches of rain fell, instead of the average 11.84 inches. In November only 0.85 of an inch instead of the average 2.68, and in January only 0.64 instead of the average 2.07. There was no sewer-flushing rainfall during the entire period, and the city sewer flushing, always inadequate, was wholly suspended in January on account of the lack of funds. The sewers were congested with filth, of which typhoid stools formed a component part, and the surface of the earth, in city and country alike, was covered with the five months' accumulations. The water supply was, however, uniformly good, since there was little sewage movement. From March to July inclusive was the wettest season on record—a total precipitation of 23.73 inches instead of the usual 7.87 inches. The sewers were repeatedly flushed out, and the accumulated surface filth was washed away into streams, ponds and the lake. Under ordinary circumstances the water supply would have been repeatedly and grossly polluted. Fortunately, the wind movement during this wet period was unusually light and its prevailing direction was from the east—thus keeping the sewage-laden shore waters away from the tunnel intakes. In August a succession of high variable winds set in, the strongest being from the west. The lake bottom was vigorously stirred up by high wave action, the sewage was drifted to the intakes, and the water supply from all sources became so contaminated that it averaged only 38 per cent. good for the month.

1. Chicago Medical Recorder, 1902, vol. xxiii, p. 219.

CAUSES OF TYPHOID INFECTION ASIDE FROM POLLUTED WATER.

This pollution of the water supply was undoubtedly the greatest causative factor in the epidemic of the past summer, but there are one or two subsidiary factors which are not brought out in the report of the board of health, and which yet may serve to explain the peculiar localization of this epidemic. The accompanying Map 1² shows that a comparatively small area on the West Side was the region most severely affected. Within the limits of the Nineteenth Ward, which contains only one thirty-sixth of the total population of the city, there were between one-sixth and one-seventh of all the deaths

Map 1.—Showing the central part of Chicago and the approximate locations of typhoid fever deaths in August, 1902. (For map showing the entire city with the source of water supply and the sanitary drainage subdivisions, see *THE JOURNAL*, Dec. 20, 1902, p. 1563.)

from this disease. This part of the city is inhabited largely by working people. It contains one of the largest Italian quarters, most of the Greek colony, a small Bohemian colony, the northern end of the Jewish quarter and the western part is chiefly American-Irish. So far as the general intelligence of the inhabitants is concerned, their knowledge of the laws of hygiene, their general housing conditions, cleanliness, overcrowding, etc., this part of the city does not differ from the other

semi-foreign quarters, yet it suffered much more than any in this epidemic. Evidently there must have been some local conditions which favored the spread of the infection. The drinking water alone could not be responsible, for this part of the city is supplied from the Chicago-avenue and Fourteenth-street tunnels, the same water supply as that for the whole region between 47th and Lake streets, Canal street and Western avenue. Nor could the milk be chargeable, for though in this neighborhood the milk is often badly diluted, yet it averaged quite as good as that supplied to a prosperous residence district to the west, as shown by analyses made of the milk of both districts by the University of Illinois in 1898.

FAULTY SEWAGE CONDITIONS.

To those who studied the distribution of the cases of typhoid fever it soon became evident that the number was greatest in those streets where removal of sewage is most imperfect. This is an old part of the city, the sewers in many of the streets were laid before the great fire, at a time when the neighborhood was more sparsely settled and when usually not more than one family occupied each house. Adequate at that time, they are far from adequate now, and it takes only a moderate increase in the rainfall to make the sewage back up into vaults and closets, while clogging is of common occurrence in dry weather. The yards and closets are often below the level of the street and are therefore easily overflowed. Last spring during the flooding rains it was no uncommon thing to see one of these yards, from six to fourteen feet below the level of the street, covered with several inches of foul water which in the neighborhood of the privy was distinctly sewage-contaminated. In this way the earth of the yards and that under the basement tenements became soaked with diluted excreta.

This condition of things is made possible by the primitive arrangements for the disposal of dejecta which prevail in this part of the city. Two of the residents of Hull House, which is situated almost in the center of the typhoid district, made a careful house-to-house investigation, noted the conditions as to drainage in each house and also the number of cases of typhoid fever which had appeared in each during the three months in question. The results are shown in the accompanying Maps 2 and 3. It may be remarked in passing that there was no other way of ascertaining these facts except by such a house-to-house canvass. There are no records in Chicago of sewer connections which antedate the great fire. The records were destroyed at that time and have never been replaced.

It will be seen from the maps that the methods of disposal of excrement in this neighborhood have been divided into five classes. The first class includes such equipment as complies with the law: plumbing with adequate water-flushing, traps and a periodically cleaned catch basin; in other words, modern sanitary plumbing. This form of plumbing was made obligatory for all buildings by an ordinance passed in 1896. The second class includes imperfect plumbing or legal plumbing so much out of repair that the excrement is not completely carried away. In several instances very bad leaks were found, the fluid overflowing on to the floor, and in two instances trickling through the ceiling and dripping into the tenement below.

The third class includes out-of-doors water closets, whose water supply is the rain water from the roof and the wash water from the kitchen sink. This is a fairly safe arrangement when the catch basin is kept clean and the sewer does not back up. The law requires catch

2. Prepared by the board of health and kindly given me by Dr. Reilly, Assistant Commissioner of Health.

basins to be of easy access and frequently cleaned, but many of them are buried deep in the earth and uncovered only when the accumulation of filth has become unbearable.

The fourth class includes the privy vaults with sewer connections but without water supply. These are cleaned either by flushing with a hose connected with the hydrant

month, but in the houses of absentee landlords the place may be flushed only once in three years or more. In some places it is impossible to ascertain whether or not there is any sewer connection, as no cleaning has been done in the memory of the oldest tenant.

The fifth class includes the old-fashioned country privy; no sewer connection, simply a hole in the ground.

Map 2.

Explanation of Maps 2 and 3.

or by a scavenger. If cleaned often enough by flushing the sewer connection can be kept open; otherwise the night soil must be removed by a scavenger. It can be seen that there is room for great variation in this class. When the landlord lives on the premises he is apt to flush his privy at least twice a year, sometimes even once a

It is often full and indescribably filthy, in spite of all efforts on the part of the tenants to keep it at least outwardly clean.

As can be seen from the maps, the cases of typhoid fever were not more numerous, on the whole, in houses where the disposal of fecal matter is most defective. Houses with good plumbing which are close to those with badly-drained cesspools are just as apt to have cases of typhoid as are the badly-drained ones, e. g., Morgan street on the east side. All that can be said is that in those areas which have many such places there was more typhoid than in those in which conditions are better. (Compare Bunker with Taylor street, for instance, Polk street east of Halsted with Polk street west of Halsted, Harrison with Sholto.) It must be remembered that the character of the plumbing is due largely to the newness of the building and to the conscience of the landlord, and is not a measure of the standard of living of the tenants.

The statistics which the Hull House residents were able to gather as to the number of cases of typhoid fever are as accurate as such statistics ever can be. No cases

were entered on the maps which had not had a physician's diagnosis, and inasmuch as the physicians of the neighborhood testified that there was an unusually large number of cases of "walking typhoid," many of which very likely never saw a physician, it is probable that the number of cases, as given in the maps, is below rather

Taylor street (Italian). Old-fashioned privy; no sewer connection. One of five privies in a yard between a 4-story front tenement and a 3-story rear tenement. While we were inspecting it, a woman came down with a vessel filled with discharge from a typhoid patient, which she emptied into the vault. No disinfectant was used.

Map 3.

than above the actual number. It was also impossible to show on the maps the actual condition of the various closets. Cesspools filled to overflowing and leaking are necessarily represented by the same shading as moderately full and periodically cleaned cesspools, to avoid overcomplicating the maps. A few extracts from the note-books of the Hull House residents will give an idea of some of the conditions found:

DeKoven street (Jewish). Vault, connected with the sewer, but full. Basement full of sewage-contaminated water from backing up of sewer.

Law avenue (Greek). Seventy-six persons using 3 small closets under the house; very filthy; apparently no sewer connection.

Bunker street (Bohemian and Polish). Unconnected vaults; very foul; 10 cases of typhoid fever, with 4 deaths in this tenement.

Law avenue (colored). Connected with the sewer, but out of order; full to the floor. Boards at back are broken away so that cesspool is quite exposed.

Ewing street (Italian). Cesspool, said to have sewer connection, but full and running over, so that stream of sewage runs down the yard.

Aberdeen street (Irish). One large vault, used by 16 families; very foul-smelling; unconnected with the sewer. This was cleaned by a scavenger during August, and the filth left standing in an open barrow in the alley between two houses for a week. It was so offensive that the tenants in these two houses were obliged to keep their windows on that side closed. Complaints to the health department and garbage inspector were fruitless, and finally the personal influence of a physician prevailed over the landlord and he removed it, but not until it had stood there during a week of warm weather when, naturally, the place swarmed with flies. There were five cases of typhoid fever in each of the two houses next to the alley.

INFECTION BY THE HOUSE FLY.

Now if there is any causative relation between the condition described above and the distribution of the cases of typhoid fever the agency of flies at once suggests itself as a possible factor. It is true that germ-laden dust blown by the wind may also be a mode of conveyance of the infection. The typhoid bacillus has been shown by Firth and Horrocks to retain its vitality in dry soil for twenty-five days, by Uffelmann for over sixty. However, it was only after the middle of August

that this agency could have come into play to any great extent, for up to that time there had been constant rains and there was practically no dust.

The importance of the common house fly in the spread of typhoid infection was emphasized by Majors Reed, Vaughan and Shakespeare in their report on the "Origin and Spread of Typhoid Fever in the U. S. Military Camps During the Spanish War in 1898." They state that in many of the camps:

Flies were undoubtedly the most active agents in the spread of typhoid fever. Flies alternately visited and fed on the infected fecal matter and the food in the mess tents. More than once it happened when lime had been scattered over the fecal matter in the pits, flies with their feet covered with lime were seen walking over the food. Typhoid fever was much less frequent among members of messes who had their mess tents screened than it was among those who took no such precaution. Typhoid fever gradually died out in the fall of 1898 in the camps at Knoxville and Meade with the disappearance

can not be accounted for by infected water or milk; also puzzling instances of individuals, who have taken all usual precautions during an epidemic, and yet fall prey to the disease. These cases may best be explained by assuming that the food became infected by flies which had previously fed on typhoid discharges.

In a recent number of the *British Medical Journal*⁴ there is a report by two British army surgeons, Majors Firth and Horrocks, of an investigation into the "Influence of Soil, Fabrics and Flies in the Dissemination of Enteric Fever." I quote simply the last part of the paper, that which deals with flies.

These investigators had adult and larval flies placed in a specially-constructed cage in which were also placed Petri dishes containing a mixture of syrup, a fresh typhoid stool and a fresh culture of typhoid bacilli, and sometimes the whole was lightly covered with earth. After feeding on this the flies would settle on other Petri

TABLE OF CULTURES BELONGING TO COLON-TYPHOID

	Morphology.	Gelatin Plate After Forty-eight Hours.	Agar Plate (Hiss) After Twenty-four Hours.	Potato.	Gel. Stab.	Agar Streak.
Typhoid control No. 15.	Short, slender rods, rounded ends, sporeless, very actively motile; decolorizing by Gram.	Deep colonies are small, yellowish, granular, round. Surface colorless, delicately veined, edge lobulated. Center turning yellowish.	Deep colonies are greenish, bristly or thready, small. Surface colorless, granular, greenish center.	Moist, slightly slimy, colorless, growth scanty.	Delicate, granular, growth along stab, thin, greenish on surface, not spreading much. No liquefaction.	Whitish, shining streak, even rounded borders, spreading very little. White precipitate in water at bottom of tube.
A 1 from vault	Same	Same	Same	Same	Same	Same
A 2 from yard	Same	Same	Same	Same	Same	Same
A 3-a from yard	Same	Same	Same	Same	Same	Same
P 1-a from yard	Same	Same	Same	Same	Same	Same
P 2 from sick room.	Same	Same	Same	Same	Same	Same
A 3-b from yard	Same	Same	Same	Moderately abundant growth, slightly yellowish. Potato brown in two days.	Same	Thin, grayish, spreading more than 15.
P 3-a f'm vault	Same	Same	Same	Same. Potato brown in 24 hours.	Same	Same
A 3-c from yard	Same	Same	Same	Like 15	Same	Same
P 3-b f'm vault	Not so actively motile as 15.	Same	Same	Dry, granular, brown growth, all over surface.	Same	Same
A 4 from yard.	Like 15.	Same	Same	Like 15	Same	Same
P 1-b from yard	Same	Same	Deep colonies brown, granular, large, smooth-bordered. Surface large, granular, brown.	Same	Same	Like 15
P 3-c f'm vault	Not so actively motile as 15.	Same	Same	Abundant, slimy, colorless growth.	Same	Rather more abundant than 15.

of the fly, and this occurred at a time of the year when in civil practice typhoid fever is generally on the increase. . . . The first pits at Knoxville contained, before the first twenty-four hours had passed after the arrival of the troops, fecal matter infected with the typhoid bacillus. Flies swarmed everywhere. Instead of abating, the disease increased. The soldiers were using the same water used exclusively by the inhabitants of West Knoxville, and among the latter there was not at that time a case of typhoid fever. Certainly the disease was not disseminated through the drinking water.

Dr. H. A. Veeder,³ writing on the same subject in the fall of 1898, said that in his opinion flies were more important agents in the spread of typhoid infection in the military camps during the late war than were drinking water or milk.

There have been several mysteriously spreading epidemics, in farming regions, for instance, where the houses are far apart and where the spread of the disease

dishes which contained sterile culture medium. The typhoid colonies were recovered in every instance.

EXAMINATION OF FLIES.

Interesting as these results are, they do not really prove the point in question, the agency of flies in the dissemination of typhoid infection. They show, to be sure, that flies can take up typhoid bacilli from fresh cultures and then replant them a few minutes after on sterile culture medium, but it remained to be proven that flies having fed on mixed discharges, which have lain for some time in privy vaults, can subsequently crawl over various surfaces and yet finally deposit the living typhoid bacilli on foodstuffs at a considerable distance from the original place of infection. In order to do this, it would be necessary to catch the flies on the spot and make the bacteriologic examination without having subjected them to any artificial conditions.

3. Medical Record, 1898.

4. Brit. Med. Jour., Sept. 27, 1902.

Two places were selected in the neighborhood of Hull House as specially favorable for such an examination. The first was an unconnected privy on Polk street into which the discharges from two cases of typhoid fever were being thrown without any attempt at disinfection. The vault was either very shallow or very full, for the dejecta lay within three feet of the opening and had caught on the projecting scantling within a foot of the opening. The flies⁵ caught within the vault, on the fence of the yard and inside the sick-room of one of the patients, which was also used as a kitchen, were dropped into test tubes containing culture medium and allowed to remain there for periods varying from fifteen minutes to twelve hours. Eight tubes were thus inoculated, three of which contained plain bouillon in which the flies were drowned, three the slightly acid gelatin-agar of Hiss and two the potato-water gelatin of Holz. In two of the bouillon tubes, the one from the sick-room and the one

agar. From three of these tubes the typhoid bacillus was recovered; that is, from one bouillon tube inoculated with flies from the privy and from one bouillon and one agar tube inoculated with flies from the walls of the house and from the fence. From another bouillon tube with flies from the fence a bacillus was isolated belonging to the intermediate group above referred to, and another also of this group was isolated from one of the bouillon tubes which contained the typhoid bacillus.

At the time the collection was made there were no fresh typhoid discharges being emptied into this vault, so that the presence of living typhoid bacilli on the legs of the flies apparently proves either that the bacilli already in the vaults were living and multiplying, or that the discharges of the recovered cases still contained typhoid bacilli. In the Polk-street house the flies caught in the sick-room are especially interesting. This room was also used as the kitchen and at the time the collec-

GROUP, OBTAINED FROM TUBES CONTAMINATED BY FLIES.

Litmus Milk.	Litmus Whey.	Lactose Bouillon.	Glucose Bouillon.	Saccharose Bouillon.	Glucose Agar.	Indol in Peptone Bouillon. 7 days at 37° C.	Agglutination.
Faintly acid at the end of twenty-four hours. No coagulation after eight weeks.	Faintly acid after twenty-four hours and after four weeks. Hardly at all cloudy.	Uniformly cloudy. No pelticle, no acid, no gas.	Acid, no gas.	No acid, no gas.	Uniformly cloudy. No gas.	—	1-200+ 1-400—
Same	Same	Same	Same	Same	Same	—	1-200+ 1-400—
Same	Faintly acid in twenty-four hours. Neutral in twelve to sixteen days, then alkaline. Like 15	Same	Same	Same	Same	—	1-150+ 1-250—
Same	Same	Same	Same	Same	Same	—	1-100+ 1-250—
Same	Same	Same	Same	Same	Same	—	1-250+ 1-400—
Alkaline in twenty-four hours. Dark blue in two weeks. Persistent, growing translucent.	Alkaline in twenty-four hours. Persistent. Slightly cloudy.	Same	No acid	Same	Same	—	1-100+ 1-250—
Same	Same	Same	Same	Same	Same	—	1-25—
Faintly acid in twenty-four hours. No coagulation. Alkaline on fifth day, persistent. Bleaches finally.	Faintly acid in twenty-four hours. Alkaline on ninth day. Cloudy.	No acid	Acid, gas	Same	Gas, not abundant.	—	1-25—
No acid. Alkaline on third day. Persistent. Bleaches finally.	Faintly acid in twenty-four hours. Alkaline on fourth day. Cloudy.	No acid	Acid, gas	Same	Same	—	1-70+ 1-150—
Acid in twenty-four hours. Coagulating on third day. Alkaline on fifth day. Persistent.	Acid in twenty-four hours. Alkaline by ninth day. Cloudy. Persistent.	Faintly acid in twenty-four hours. Alkaline by third day, persistent.	Acid, gas	Same	Same	Slight+ reaction.	1-25—
Acid in twenty-four hours. Coagulating on third day.	Acid in twenty-four hours. Cloudy.	Acid gas	Same	Acid, gas	Abundant gas.	+	1-25—
Acid in twenty-four hours. Coagulating on third day.	Same	Acid gas	Same	Acid, gas	Abundant gas.	+	1-25—

from the yard, a bacillus was isolated which corresponds culturally with the typhoid bacillus, and which agglutinates in dilutions of 100 to 200 with the serum of immunized animals. In one of the tubes inoculated by flies from the vault a bacillus was isolated closely related to but not identical with the typhoid bacillus, belonging apparently to the group intermediate between the typhoid and colon groups.

The second place chosen was a yard on Aberdeen street containing one large, full and filthy vault, not connected with the sewer. This is used by sixteen families. Flies from the three privies built over this cesspool were used to inoculate two tubes of bouillon and one each of agar-gelatin and of potato-gelatin; other flies from the fence of the yard and from the walls of the two houses bounding the yard, at varying distances from the vault, were dropped into two tubes each of bouillon, gelatin and

tion was made the table was covered with food on which could be seen flies both living and dead.

The conditions found at these two places are repeated over and over again in this neighborhood, and when one considers that, in addition to the privies and defective plumbing, there are the sewer-infected back yards, that food is kept more or less exposed, not only in the houses, but in the groceries and fruit stands, that the houses are not furnished with screens, it can easily be seen what an important part the house fly may play in the spread of typhoid infection in such a neighborhood. It can also be seen that this is one factor which does not come into play to such an extent in more well-to-do parts of the city, where screens and good plumbing are the rule, or in newer parts of the city where the excrement is properly disposed of. There are, as was stated above, no available statistics as to the number of privies still in use in Chicago, but Dr. Reilly, assistant commissioner

5. In all cases the common house fly, *Musca domestica*.

of health, is of the opinion that more than half of them are within the Nineteenth Ward. It seems, therefore, not unwarranted to assume that the concentration of the typhoid fever in this particular quarter of the city during the late epidemic was due to the spread of the bacilli by flies from the undrained or imperfectly drained cess-pools, the leaking closets and the infected earth of the yards.

BACTERIOLOGIC EXAMINATION OF THE FLY TUBES.

As already stated, the flies caught at the different places were dropped into six tubes of bouillon, six of potato-water gelatin and six of Hiss' agar-gelatin. They were allowed to walk over the solid, slanted media, sometimes for fifteen to thirty minutes only, sometimes from one hour to twelve hours; others were drowned in the bouillon tubes. The flies were then removed with sterile forceps and the tubes set aside for twenty-four hours, the gelatin tubes at room temperature, the others in the incubator. All of the gelatin tubes liquefied in forty-eight hours to such an extent that they were useless. The agar tubes, with the exception of two in which the flies had remained not more than half an hour, were covered with a thick slimy growth, loopfuls of which were transferred to bouillon and the plating was done from the latter. The two in which the growth was not so abundant showed two or three black specks of excrement in each, and the colonies growing at these points were carefully examined, but yielded only sarcines.

The plating media used were the potato-water gelatin, the agar-gelatin of Hiss, and Löwenstein's .05 per cent. carbol-gelatin. Plates were made from the twelve tubes on all these media, and five of them were found so greatly contaminated as to be worthless. Sarcines, yellow, white and orange, liquefying protei, subtilis, megatherium, Zopfii, mesentericus and mycoides were all identified by their colonies, and the attempt to isolate the colon group from among them failed. Better success was obtained from the remaining five bouillon and two agar tubes. Two of them (A 1 and P 1) were so little contaminated with liquefying varieties that the original Holz plates remained usable until they dried. This medium was somewhat disappointing in the other cases, for the liquefying varieties often grew with apparently undiminished vigor, causing complete liquefaction by the time the typhoid colonies should begin to be characteristic. The carbol-gelatin was still less satisfactory, for the addition of .05 per cent. carbolic acid did not prevent the growth of the liquefying forms, but only retarded them; but it also retarded the growth of the typhoid colonies which often at the end of three days showed merely as clear droplets, not distinguishable from colon, or from a slowly liquefying motile bacillus. Moreover, the typhoid surface colonies never took the characteristic grape-leaf appearance, but were always round with smooth edges. On the whole, the agar-gelatin of Hiss gave the most satisfactory results. The tiny, greenish, bristly or thready colonies which developed in eighteen hours were easily distinguishable from the larger, brown, smooth-bordered colon colonies. The differences became more marked in forty-eight hours, for the typhoid colonies increased hardly at all in size, while the colon colonies grew much larger. The acidity of the medium retarded the growth of other forms.

Hiss⁶ has recommended the transplantation of typhoid-like colonies to a semi-solid "tube medium" (containing glucose), in which, he claims, the typhoid bacillus is the only one causing a diffuse cloudiness without forma-

tion of gas, and in this way a rapid positive diagnosis can be made. In my experience this served very well to distinguish the typhoid bacillus from the colon and from various glucose-fermenting intermediate forms, but not from a non-sugar-fermenting bacillus which produces an early and persistent alkalinity in milk and which I have supposed to be identical with the *Bacillus fecalis alkaligenes* of Petruschky. This bacillus caused a diffuse cloudiness in the tube medium, and inasmuch as it resembled the typhoid bacillus morphologically and in its growth on gelatin, agar and bouillon, it could be distinguished from the latter only by its rather more abundant brownish growth on potato and its failure to ferment any of the sugars or to cause even a transient acidity in milk or litmus whey.

Colonies of the colon bacillus were found on every plate, as was to be expected, and two organisms were isolated simply for purposes of comparison. Several intermediate forms were also isolated from the Hiss plates under the supposition that they were typical typhoid colonies, until the formation of gas in glucose media proved their nearer relation to the colon group.

The following is a diagram of the twelve organisms which were isolated from the fly tubes, and of a typhoid bacillus obtained in the laboratory of Rush Medical College from the blood of a typhoid patient and used as control. Those marked A were from the tubes inoculated at Aberdeen street; those marked P, from Polk street. The organisms A 3-c and P 3-b seem to belong to the group intermediate between the typhoid and colon bacilli, resembling the typhoid morphologically, in their growth on potato, gelatin and agar, and in their failure to ferment lactose or coagulate milk; resembling colon in the fermentation of glucose with formation of gas as well as acid, although not as abundantly as does colon. To this group belong the *B. enteritidis* of Gärtner, and the organisms isolated by Widal, Gwyn, Cushing, Schottmueller, Longcope, Hewlett, Johnston, Kurth, and recently Sion and Negel. There are minor variations in the growth of these organisms on culture media as they approach now the typhoid type, now the colon, but in the main the class is well defined.

P 3-b, with its dry, granular, brown growth on potato, early and persistent alkalinity in milk and litmus whey and its slight fermentation of glucose, corresponds closely with the bacillus isolated by Sion and Negel from the blood of five patients and from the well-water in a house epidemic of typhoid-like character. A 3-b and P 3-a ferment no sugars and produce an early and increasing alkalinity in milk and Petruschky's whey, and a relatively abundant growth on potato, characteristics which correspond to those of the *B. fecalis alkaligenes* of Petruschky.

P 3-c and P 1-b are typical members of the colon group, except that P 3-c does not grow characteristically on potato. A 4 should, perhaps, be relegated to the intermediate group, but differs from all those reported, except one of Schottmueller's, in coagulating milk and forming a small amount of indol. It might be regarded as a feebly-growing organism of the colon group were it not for the failure to ferment lactose—the transient faint acidity is soon followed by a persistent alkalinity—and the production of a persistent alkalinity in milk and litmus whey.

The serum of an immunized rabbit was used to test the agglutinability of these organisms. The control agglutinated in a dilution of 1-200, not in a dilution of 1-400. All of the typhoid bacilli isolated from the fly tubes failed to agglutinate in dilutions of 1-400, two

6. Jour. Med. Research, 1902, vol. viii, No. 1, p. 148.

would agglutinate only in 1-100 dilution, and one in 1-150. P 3-b agglutinated in a dilution of 1-70, which is really greater than the dilutions used in the ordinary clinical application of the agglutination test, yet this organism proved by its growth in sugar media and milk to belong to the intermediate group. This agglutinability with typhoid immune serum on the part of an organism of the intermediate group has been observed several times (Durham, Klinger). Durham for this reason questions the value of the agglutination test as compared with cultural tests, for he has even found colon bacilli which agglutinated with this serum, as has also Klinger.

As all parts of the fly came in contact with the culture medium it is impossible to say that any one part was especially responsible for the conveyance of the bacilli. Firth and Horrocks were able to recover the bacillus from the head, wings, legs and body of the flies which had crawled over the typhoid cultures. The failure to isolate it from the excrement of the flies also agrees with the findings of Firth and Horrocks.

CONCLUSIONS.

1. The epidemic of typhoid fever in Chicago during July, August, September and October of 1902 was most severe in the Nineteenth Ward, which, with one-thirty-sixth of the city's population, had over one-seventh of all the deaths from this disease.

2. The concentration of the epidemic in this locality can not be explained by contamination of the drinking water, or of food, or on the ground of ignorance and poverty of the inhabitants, for the Nineteenth Ward does not differ in these respects from several other parts of the city.

3. An investigation of the sanitary conditions of this region shows that many of the street sewers are too small, and that only 48 per cent. of the houses have sanitary plumbing. Of the remaining 52 per cent. 7 per cent. have defective plumbing, 22 per cent. water closets with intermittent water supply, 11 per cent. have privies connected with the sewer but without water supply, and 12 per cent. have privies with no sewer connection.

4. The streets in which the sanitary arrangements are worst had the largest number of cases of typhoid fever during this epidemic, irrespective of the poverty of the inhabitants.

5. Flies caught in two undrained privies, on the fences of two yards, on the walls of two houses, and in the room of a typhoid patient were used to inoculate 18 tubes, and from 5 of these tubes the typhoid bacillus was isolated.

6. When the discharges from typhoid patients are left exposed in privies or yards, flies may be an important agent in the dissemination of the typhoid infection.

Results of Nephrectomy.—V. Carlier describes in *Echo Med. du Nord*, vi., 49, the ultimate results in twelve cases treated by nephrectomy. The reflex irritation from the diseased kidney ceases with the extirpation of the organ, and it frequently happens that a patient afterward will tolerate drugs, mercury for instance, which caused serious disturbances before. The best results from the nephrectomy were attained in cases of renal tuberculosis derived from the blood. One of the patients contracted scarlet fever during convalescence, but the course of the intercurrent disease was normal, with no trace of albuminuria until the third week. In debating the question of intervention he is inclined to distrust patients whose urine contains an abnormal proportion of chlorids, either too little or an excess. A patient in the latter category died three months after a successful nephrectomy, from generalization of the tuberculous process.

PNEUMONIA: THE NEW "CAPTAIN OF THE MEN OF DEATH."

ITS INCREASING PREVALENCE AND THE NECESSITY OF METHODS FOR ITS RESTRICTION.*

ARTHUR R. REYNOLDS, M.D.

Commissioner of Health.

CHICAGO.

Figures are hard reading and harder hearing; but I must ask you to listen to a few in order that you may comprehend the magnitude of the pneumonia problem as I view it.

In the latest (1901) edition of his scholarly and most instructive "Principles and Practice of Medicine" Prof. William Osler, treating of pneumonia in the section of specific infectious diseases, says:

The most widespread and fatal of all acute infectious diseases, pneumonia, is now the "Captain of the Men of Death," to use the phrase applied by John Bunyan to consumption.

He then adds:

In the United States during the census year 1890 there died of it 76,496, a death rate per 100,000 of population of 186.94 [?]. In Chicago during the past ten years it has gradually replaced consumption as the principal cause of death, which A. R. Reynolds attributes to the *predisposing influence of Influenza*. In the last decade the death rate from pneumonia in Chicago was 18.03 per 10,000 of population, against 12.36 per 10,000 in the previous decade. There has been a marked increase in the disease in Baltimore, and Folsom has brought forward evidence to show that there has been a progressive increase in the death rate from pneumonia in the state of Massachusetts. The admission of pneumonia cases to hospitals during the past few years has, in some places, almost doubled.

These passages are cited for two purposes: First, that I may put their reader on his guard against the assumption that the great increase in pneumonia during the last ten or a dozen years is attributed by me solely "to the *predisposing influence of influenza*." Second, that I may emphasize more fully than Professor Osler has done the extent of this increase, both in point of time and of proportion.

As to the first, it will probably be sufficient to call attention to the title of the paper on which the assumption was based. This paper appeared in the March, 1901, Bulletin of the Chicago Health Department, and was entitled "Influenza as a Factor of Recent Mortality"—not the only, but one factor. That influenza has been such a factor and a very important one since December, 1889, is now admitted by etiologists generally. That the increase of pneumonia mortality is due solely to its predisposing influence is disproved by the following figures of deaths from all causes, those from consumption and those from pneumonia during the last thirty years:

In 1880 the death rate from all causes was 14.59 per thousand of the total population of the country; in 1900 it was 13.67—a decrease of 6.3 per cent. in the general mortality rate during the twenty years.

The consumption death rate in 1880 was 18.21 per 10,000 of population and 12.46 per cent. of the total mortality; in 1900 these figures were 14.44 and 10.56 respectively—decreases of 20.7 per cent. in proportion to population and of 15.2 per cent. in proportion to total mortality.

* Read by invitation at the Sixth General Conference of Health Officials in Michigan, under the Auspices of the Michigan State Board of Health, Jan. 15, 1903, at Ann Arbor, Mich.

EDITORIAL NOTE.—See Editorial, "The Deadly Pneumococcus," in THE JOURNAL A. M. A., April 14, 1900, probably the first printed notice of the increasing importance of pneumonia as a factor in the mortality of recent years.