Cowgill⁴ utilised these results as a test for the presence of vitamin B in natural food-stuffs. He found that the potency in promoting appetite ran parallel to the potency in relieving symptoms in polyneuritic animals.

Lumière^{5,6} fed pigeons with decorticated rice and noticed a marked diminution of intake. There was evidence of stasis in the upper part of the digestive tract. The rice was seen to lie as a compact mass in the stomach. Trituration was defective, and the onward passage of the bolus through the pylorus was greatly delayed. There appeared to be a condition of deficient glandular secretion and hypotonus of the intestinal wall. A small dose of extract of rice polishings provoked abundant glandular secretion and stronger intestinal movements which resulted in rapid evacuation of the masses of polished rice lying in the upper intestine. He found that these animals die sooner than starving animals, and quotes Forster, who obtained similar results in dogs.

There appears, therefore, to be a considerable accumulation of evidence that the appetite is adversely affected by the absence of vitamin B from the diet of animals. It is insufficient, however, to state that the vitamin acts by diminishing appetite. Appetite is a general conception, the physical expression of

Chart showing weight and intake curves of two rats. R2: both vitamins administered throughout. R1: both vitamins administered at first. At A both vitamins were removed; at B a trace of marmite was added to diet. Note sudden falling off of food intake at A and rapid recovery at B.

which is the food intake. The problem still remains to explain why the intake is lowered when the vitamin is withheld. It is a general experience that on post-mortem examination of rats fed on diets deficient in vitamins A and B to find the stomach full of food and active fermentation in progress. Sometimes the stomach is definitely dilated. This finding suggests that there is some defect in the power of the stomach to deal with its contents and propel them farther down the intestinal canal. Lumière, too, found the food lying as an inert mass in the stomach of the pigeons experimented on. From a study of the work detailed, it is suggested that the diminished intake may be accounted for quite simply in the following way. The contractions of the stomach wall are weak, the secretion of the gastric glands is defective, and as a result the stomach is never empty. It is always partly distended with fermenting material. Hunger contractions do not take place, the craving for food becomes low, and the intake falls off considerably. This theory, that the defect lies in the stomach, receives valuable support from the fact that the animal responds so very rapidly to the re-addition of the vitamin to the diet. In rats the rise in intake is marked at the end of 24 hours. In pigeons, rapid evacuation of the contents of the intestinal canal occurred immediately after administration of the

vitamin. Admitting the fact of lowered food intake, it must be borne in mind, however, that the loss of weight shown by the animals, and their ultimate death, can still only be partially accounted for. In Drummond's experiments,¹ improved intake was obtained by adding flavouring agents to the diet; growth, however, only occurred when the vitamin was added. In one of Karr's dogs, though the intake remained high in the absence of the vitamin, general symptoms of the vitamin deficiency developed. Forster's experience, that dogs succumbed more rapidly on a vitamin-free diet than when completely deprived of all food, is also very significant.

It may be concluded with some degree of assurance that vitamin B produces its effects by acting primarily, if not solely, on the intestinal canal. In its absence muscle-tone is diminished, peristaltic movements are weak, and there is a reduction in the amount of digestive juice secreted. As a result, stasis occurs, the appetite flags, and the food intake falls off considerably. It is almost certain, however, that the animal is still consuming sufficient to enable it to survive for considerably longer periods than it actually does. The explanation for the early decline may perhaps be found in the changes which occur to the food which is retained for an abnormally long time in the intestine. Fermentation soon sets in and toxic bodies may be produced which adversely affect the general health of the animal, death resulting from the combined effects of all the factors mentioned.

Conclusion.

Vitamin B acts by facilitating the efficient carrying out of the functions of the intestinal canal. The main effects produced by the absence of the vitamin—i.e., diminished food consumption, loss of weight, and ultimate death—are due to intestinal stasis and the absorption of toxic bodies which results therefrom.

The expenses of this research have been partly defrayed by a Government grant made through the Royal Society. The prepared palm-kernel oil was kindly presented to me by Messrs. Lever Bros., to whom I tender my thanks.

References.—1. Drummond, J. C.: *Bioch. J.*, 1918, xii., 25. 2. Drummond, J. C.: *Ibid.*, 1919, xiii., 94. 3. Karr, W. G.: *J. Biol. C.*, 1920, xlv., 255. 4. Cowgill, G. R.: *Proc. Soc. Exp. Biol. and Med.*, xviii., No. 8, 290. 5. Lumière, A.: *Presse Médicale*, 1920, No. 20, May 8th. 6. Lumière, A.: *Bull. de l'Acad. Méd.*, 1920, March.

AORTIC INCOMPETENCE:

A CLINICAL STUDY.

By GEORGE J. LANGLEY, M.D., M.R.C.P. LOND.,

PHYSICIAN, ANCOATS HOSPITAL; ASSISTANT PHYSICIAN, SALFORD ROYAL HOSPITAL, MANCHESTER.

THE following analysis of cases of aortic valvular disease was undertaken largely with a view to discovering, if possible, those untoward signs in routine clinical examination which might possibly be utilised to give some warning of the sudden terminal event which so frequently disconcerts the patient's relatives and the physician in charge of the case.

Material and Incidence.

Fifty cases have been taken in the unselected order of their coming under hospital care, and form one section of a series of about 400 heart cases which have been examined. Thus aortic disease would constitute 12½ per cent. of the cardiac cases, a much lower proportion than that found at the Edinburgh Infirmary (Gillespie, *Edin. Hosp. Rep.*, 1898). The lower percentage is no doubt accounted for by the large number of cases sent to a heart clinic nowadays and classified under the heading "effort syndrome." In no instance in this series was a diagnosis of aortic stenosis made, the only cases showing a thrill at the base of the heart being such as were shown by X ray to have an aneurysm. The average age of the series was 38 years, the youngest patient being 21 and the oldest 67. Only five cases occurred in women and 40 of the cases were married.

Pathogeny.

The pathogeny may be considered under five headings, as follows:—

(1) *Rupture of a valve segment.*—One case may possibly have originated by rupture of a valve segment, but there is no conclusive evidence. The patient fell down a cliff side with a machine-gun on Gallipoli in 1915, was picked up unconscious, found to have a cardiac lesion, invalided home and out of the service, and died in hospital from cardiac failure with general anasarca. The diastolic murmur was always high-pitched and musical in quality. No post-mortem was permitted. No history or evidence of past rheumatism or syphilis was obtainable. The total duration of the disease was five years.

(2) *Acute rheumatism.*—There is considerable difficulty in separating the cases due to rheumatic endocarditis from those not so caused. Two methods were attempted, firstly by careful collection of childhood and adult histories with regard to rheumatic affections, including chorea, tonsillitis, scarlet fever, growing pains, &c. Thus 21 cases gave evidence of rheumatic fever; two other cases gave a history of severe sort-throat or scarlet fever which appeared to stand in direct causal relation to the disease. Two cases had suffered two attacks, and one case three attacks, of rheumatic fever. In only one case of the series was a history of rheumatic affections in the family obtained. Of the female cases all were definitely rheumatic in origin, except one case which at the post-mortem was found not to have aortic incompetence. In only 11 of the 21 cases was the cardiac lesion discovered at the time of the rheumatic attack. The second method attempted for separating the cases of rheumatic origin was from the character of the mitral lesion found. Rheumatic endocarditis is more likely to affect the mitral valve, giving rise to stenosis, than any other ætiological factor at present known. Sixteen cases were found to present a presystolic crescendo murmur at the apex. No attempt was made to differentiate the murmur of true stenosis from the Austin Flint murmur. In my opinion it is not usually possible. Of the 16 cases presenting two murmurs, aortic diastolic and mitral presystolic, there was a definite history of rheumatism in 11 cases, a positive Wassermann reaction (W.R.) in three cases, and evidence of arterio-sclerosis in two cases. It is probable that the presystolic murmur heard in the syphilitic and arterio-sclerotic cases was produced by the method to which Flint drew attention. No post-mortem evidence is yet available.

(3) *Syphilis.*—W.R.'s were tested in 45 cases; 16 of them proved to be positive, 11 being positive 1 in 40 dilution of serum, and 5 positive 1 in 5 dilution. In four cases there was a history of rheumatic fever as well as a positive W.R.; in two of these the cardiac lesion appeared to be the direct and immediate outcome of the rheumatism, but in the other two the heart lesion was found 18 and 26 years after the rheumatic attack, and was probably syphilitic.

(4) *Arterio-sclerosis.*—Twelve cases were found in which neither of the preceding ætiological factors were active. In seven of these the occupation was known to entail unusually heavy work, but in the other five cases the work was light. Two of these cases were found to have deficient renal function.

(5) *Infective endocarditis.*—With regard to an infective active endocarditis without associated rheumatic phenomena, one case had typical acute fulminating ulcerative endocarditis, and four other cases were regarded as subacute bacterial endocarditis.

Duration.

Of the fatal cases the time during which a cardiac lesion was known to have existed averaged only two and two-thirds years. Taking the whole series together the number of years the lesion was known to have existed at the time of examination averaged three and a half years, the period of collection being eight months. Of the 50 cases six were discovered incidentally in the course of general routine medical examination, and were entirely latent, giving no symptoms recognised by the patient. To this number must be added two other cases similarly found, which could not, on account of their nature, be added to this series.

Symptoms.

Pain.—Not severe in character, but recurring at fairly frequent intervals; pain over the præcordium occurred in 26 cases; anginal attacks in two cases; three cases complained of epigastric pain.

Dyspnoea.—Occurred in all cases to some degree, reaching a serious point in six cases; asthmatic attacks repeatedly occurred in two cases, both of

which also had anginal attacks. Twenty-one of these patients had some degree of orthopnoea, but in only five was it distressing.

Cough.—Troublesome from time to time in almost all the cases, but forming only in very few the outstanding feature of the disease, even for a short time.

Vertigo.—An almost constant symptom; in several cases the reason for seeking medical advice.

Predisposing Causes.

Various causes have been regarded as contributing either to the actual production of disease or to the final breakdown.

Tobacco.—So far as this series is concerned, tobacco did not appear to affect the issue. Patients who had been heavy smokers, using more than ten cigarettes per day, or three ounces of tobacco per week, were not found; but in all cases where the cardiac condition was giving rise to symptoms the patients volunteered the information that they had given up smoking as they found that it made them worse.

Alcohol.—There was good reason to believe that alcoholic excess had occurred in five cases. On the other hand, 17 cases stated that they were lifelong abstainers.

Games.—In no case was there reason to suppose that playing games stood in any causal relationship to the disease. Neither could it be used to determine the onset of the lesion; several patients who had suffered an attack of rheumatic fever in early life played games successfully and without any difficulty in adolescence.

Work.—Known to have been unusually heavy in 27 cases—not an unusually high proportion in cases limited to the hospital class of patient. Five cases found in which the work was very light, and neither rheumatism nor syphilis had occurred.

Physical Signs.

The physical signs are as follows:—

Tremor.—Striking rarity of those fine tremors of hands and tongue so constantly seen in the effort syndrome cases. In seven cases tremor of hands was well marked, but all seven patients were obviously and seriously ill. In only one case was tremor of the neurasthenic type seen.

Pallor.—Noticed in 30 cases, slight in eight, very striking in six.

Clubbing.—Definite clubbing of fingers in ten cases, in none of which was any sign of a congenital lesion, but in four cases there were other signs of an active infection.

Petechiæ.—Much difficulty was encountered in determining the occurrence of petechiæ. In many cases, originally seen in the out-patient department, and later taken into the wards for observation, such petechial spots as were found around the neck, lateral aspects of thorax and abdomen, outer side of thighs, &c., entirely disappeared after a week in hospital. In only four cases were petechiæ satisfactorily demonstrated; these all had an infective endocarditis and are already dead.

Vessels.—The apparently enlarged, tortuous, pulsating temporal artery was observed in 23 cases; the violently pulsating neck vessels absent in only five cases. Capillary pulsation found 36 times.

Area of cardiac dullness.—This area, as determined by light percussion, was found materially enlarged in all but four cases, in which four there was other evidence that the aortic defect was only slight.

Position of heart apex beat.—Found to be as follows:—Aortic incompetence alone: fifth space, 11 cases; sixth space, 22 cases. Aortic incompetence with presystolic murmur: fifth space, 6 cases; sixth space, 11 cases.

The degree to which the apex beat extended beyond the left nipple line varied considerably, in four cases reaching as far out as the anterior axillary line. The character of the apex beat was normal in nine cases, in seven of which the aortic second sound was audible in the neck. The beat was powerful and heaving in 38 cases, but wavy and extending over several intercostal spaces in three cases—one of which had ulcerative endocarditis, one is thought to have subacute endocarditis, and the third has active syphilis, very thick vessels, and a small heart.

Murmurs.—Only three cases were found in which no murmur was audible at the apex. In 16 cases a presystolic crescendo murmur was audible. The diastolic murmur was itself heard at the apex in 34 cases, and in 10 cases only a systolic murmur was heard there. At the aortic cartilage a to-and-fro murmur was heard in 33 cases; in the remainder of the cases a diastolic murmur only was heard, among the latter the murmur difficult to hear was encountered. In 21 cases the second sound was audible at the root of the neck; in the remainder it was inaudible. As above stated, a presystolic murmur was found 16 times, and of these 16 cases, 11 gave a clinical history of rheumatic affections.

Pulse.—Varied in frequency at rest from 60 to 120 beats per minute; definitely collapsing in character in 35 cases.

Irregularity occasionally noted in four cases; pulse delay in the radial only noticed twice, but was not particularly looked for. Of the cases not recognised as having a collapsing pulse, all but two had also a presystolic murmur; of these two exceptions, one was found post mortem not to have aortic incompetence, and the other had only a small heart giving rise to no symptoms.

Blood pressure.—Brachial blood-pressure readings (with a Tyco instrument and the auscultatory method) were taken in nearly every case. In eight cases the systolic pressure was 120 mm. Hg or less. In all the rest it was above this normal. The difference between systolic and diastolic pressure was usually extremely well marked, in some cases no diastolic reading being possible. The average difference between systolic and diastolic readings was 85 mm. Hg. No observations were made with regard to arm-leg differences.

Aneurysm.—A definite aortic aneurysm, proven by X ray examination, was found to be present in nine cases, in all of which the W.R. was positive, and the electro-cardiographic left-sided preponderance was very marked indeed.

Lungs.—No lung changes were found in these cases, except the basal crepitations indicating oedema from cardiac failure, and these only in such cases as died in hospital.

Oedema.—The same may be said of oedema generally, and no history of recurrent oedema of ankles was obtained from these patients except in two cases complicated by lesions of other valves and known to be suffering from auricular fibrillation.

Of the three cases which died of slow progressive general anasarca no history of previous cardiac failure was obtainable. One patient, who had advanced renal disease, was said to have occasional variable oedema of the ankles.

Vomiting.—Vomiting was only complained of by patients when there was evidence of grave cardiac disability, and its appearance heralded the end. It occurred—but only as a very late event—in three out of the four patients who died suddenly. One patient ate his breakfast prior to coming to hospital, on his weekly visit, said he felt sick, immediately vomited, and fell dead.

Liver.—Hepatic enlargement was only found when there was other evidence of cardiac failure, except in one case where an alcoholic history, the consistence of the organ and its freedom from tenderness, seemed to justify the additional diagnosis of hepatic cirrhosis.

Spleen.—The spleen was found enlarged and palpable in nine cases. In five of these malaria was known to have occurred in the past, one patient was known to have hepatic cirrhosis, which condition was thought to account for the enlargement. Four cases were diagnosed as subacute bacterial endocarditis, and one as ulcerative endocarditis. In this connexion the frequency with which a past malarial history is obtained in these cases of subacute endocarditis appears to be greater than the fortuitous coexistence of the two diseases would warrant. There seems evidence for the belief that malaria in aortic valve lesions of old standing is liable to determine the onset of this complication.

Eyes.—Routine examination of the fundus oculi was made and pulsating retinal vessels were occasionally observed but in six cases definite retinal changes were found—1 bilateral syphilitic choroiditis, 1 bilateral optic atrophy (tabetic), 2 retinal hæmorrhages (renal; both cases dead), 2 retinal hæmorrhages (in subacute endocarditis; both cases dead).

Kidney.—The renal function was investigated by the urea-concentration test in 39 cases. Response was found to be below normal in eight cases—3 of renal disease, 1 with auricular fibrillation, and 4 with red blood cells also present, and thought to be cases of subacute endocarditis.

Tenderness.—Hyperalgesia was investigated in all cases and in only seven was any tenderness elicited, although so many of the patients complained of præcordial pain or discomfort. In two cases which gave a history resembling, if not identical with, angina, both showed marked tenderness. Two other cases, both with large hearts, showed a tender spot at the left seventh costochondral junction. The rest had only slight præcordial tenderness.

Exertion.—In nine cases the exercise tolerance was quite good, and these patients possessed little or nothing in the way of symptoms except one patient, a tabetic. The rest of the patients had a very poor response to exertion, becoming distressed and breathless on very slight provocation.

X ray.—The routine radiographic examination which was made gave most valuable information, both as finally deciding the presence or absence of an aneurysm and also in corroborating the clinical findings with regard to the cardiac size and shape. The typical aortic form of heart was reported in almost every case.

Other Diseases and Mental Change.

Other diseases were found to coexist in these patients, but always in some way related to the cardiac condition. Thus, to recapitulate: 1 ulcerative endocarditis, 4 subacute endocarditis, 1 active

syphilis, 1 tabes, 1 hepatic cirrhosis, 2 nephritis. In no case was any mental change noticed in the patients except in the tabetic, who constantly professed himself very fit indeed in spite of his manifestly parlous state. He showed no other indications of G.P.I. beyond this and his optic atrophy.

Diagnosis.

Very little evidence is available to show how frequently errors of diagnosis are made in aortic disease. Several cases had murmurs difficult to hear either by direct auscultation or with any form of stethoscope. Much help was obtained by bending the patient well forward and listening along the aortic course, but it is essential to keep a firm grip on these patients, as their vertigo is always increased by this manœuvre, and they are very liable to fall. In one case at least a diagnostic error was probably committed, and this in spite of the fact that the possible fallacy was thought of and discussed. The patient had advanced renal disease with retinal hæmorrhages, urgent dyspnoea, and signs of a very large pericardial effusion. There was a systolic murmur and an impure first sound at the apex. Diastolic murmur was not audible at the base, but was heard at the fifth left costal cartilage, soft and blowing in character. The second sound was easily heard. The possibility of pericardial friction was discussed, but ruled out on account of (a) the character of the sounds, and (b) the size of the effusion. Operation was undertaken later to remove the pericardial fluid in the hope of relieving the respiratory distress. This was to some extent successful, but the patient died four days later. At the post-mortem there was some sclerosis of the aortic valve segments, but no regurgitation to the water test.

Circulatory Conditions.

Much help was obtained from electrocardiographic tracings, conditions being revealed which could not otherwise have been made out.

P.R. interval.—No case of heart-block was found in this series, but the P.R. interval exceeded the normal in four cases—one of rheumatic disease with left-side preponderance who remains well, two of arterio-sclerosis (one died from heart failure, the other shows retinal hæmorrhages), and one case of subacute endocarditis who died suddenly.

Inversion of T.—The inversion of T in lead 2 in the absence of digitalis administration was found in four cases—1 syphilitic aortic disease with right-sided dominance, 1 rheumatic disease with left-sided preponderance, cardiac asthma and angina, 1 arterio-sclerosis, and the case of pericardial effusion which showed no aortic incompetence at post-mortem. All four of these cases were clinically very ill.

Q.R.S. interval.—No case in this series was found showing a Q.R.S. complex exceeding the normal period.

Left-side preponderance.—It was thought that if a left-sided hypertrophy were the response to mechanical aortic damage, its absence should indicate a want of recuperation on the part of the patient; except where a coexisting mitral stenosis might occasion a right-sided preponderance (R.S.P.). Investigation along these lines gave the following results: General condition remarkably good in 8 cases. R.S.P., 3; L.S.P., 3; no change, 2. General condition remarkably bad in 11 cases. R.S.P., 2; L.S.P., 8; no change, 1. Obviously the degree of regurgitation would form a large factor in the resulting hypertrophy. Estimation of the grade of regurgitation was attempted with the following results: Slight, 15 cases: L.S.P., 6; R.S.P., 5; no change, 4. Grave, 25 cases: L.S.P., 17; R.S.P., 4; no change, 4. Of those cases showing a grave degree of regurgitation and no change in the electrocardiogram two are already dead. Among the tracings a left-sided preponderance was found 21 times. In 15 of these the cardiac enlargement was marked and aortic in type, in 2 cases the clinical enlargement was slight, while in 4 cases no enlargement was made out. Of these 21 cases 5 had a presystolic murmur; 8 were thought to be rheumatic, 9 syphilitic, and 4 arterio-sclerotic.

Right-side preponderance.—Nine such cases were found; 4 had a presystolic murmur, 5 were considered rheumatic in origin. There was no evidence for regarding the R.S.P. cases as more serious than those showing L.S.P.

P-wave or auricular wave.—An examination of the P wave as regards magnitude and duration gave no help as regards the presence or absence of true mitral stenosis. Seventeen cases gave a P. deflection exceeding 3 mm. in lead 2, and of these only 6 had a presystolic murmur. In 5 the P. wave was split, but only one had a presystolic murmur, and

only one a history of rheumatism. Of the 17 cases, 6 had a rheumatic history, 5 a positive W.R., and 5 arterio-sclerosis.

Conclusions.

The prognosis in aortic disease remains almost as difficult a problem to-day as it was years ago. Nevertheless modern methods do give invaluable help in certain cases. Pathological increase in the P.R. interval or the duration of the Q.R.S. complex yields information as to myocardial damage which could not be arrived at by other means or could only be guessed at without any definite proof—e.g., P.R. interval increase, four cases, two already dead. Inversion of T. in lead 2, four cases, one already dead, and two others seriously ill. But this matter of “looking for a sign” remains as unprofitable in this century as in any other; it is only by utilising every method of investigation, old as well as new, that a correct estimate is likely to be arrived at in these difficult cases.

Active infections.—The careful and repeated examination of the urine for red blood cells, the systematic search for petechiæ, enlargement of the spleen, pallor and clubbed fingers—as indicative of the subacute bacterial endocarditis described by Libman and by Horder—is perhaps the most important procedure in the whole clinical examination of the patients, and only to a less extent, because less frequent, in all cardiac cases.

Ætiology.—Arteriosclerosis and renal disease, long regarded as ætiological factors, also demand special investigation, more particularly as regards renal function and retinal hæmorrhages. The W.R. and X ray examination both give invaluable help, especially as regards the occurrence of aneurysm, which, although present, may give no signs or symptoms at all.

Correlation of findings.—The correlation of all the findings will probably point the way, and each factor follow the expected line; yet in a certain number of cases this will not be so, and explanation of the anomaly may be extremely difficult. Where the size of the heart does not appear to correspond to the degree of regurgitation, as judged by the collapsing character of the pulse, difference in systolic and diastolic pressures, presence or absence of aortic second sound at the root of the neck, and the presence of L.S.P. in the electrocardiogram, the possibility of an adherent mediastino-pericarditis must be borne in mind; it was thought to be a possibility in only one case of this series. Of the 40 cases fully investigated, all the facts arrived at followed the expected line in 31, but in 9 cases the findings did not correspond in some particular. Of these nine cases 3 are already dead, 1 has an active endocarditis, 1 probably an adherent pericardium, and the other 4 cannot be at present explained. Those factors which herald a lethal result have already been sufficiently insisted upon, but it would seem that cases which show a want of correspondence in all the findings of a routine examination should also be regarded as of serious outlook.

I wish to thank Dr. Arnold Renshaw, pathologist, and Dr. J. M. W. Morison, radiologist, for much help in routine work, and the Pathological Department of the University of Manchester, whose generous grant made this work possible.

ROYAL SUSSEX COUNTY HOSPITAL, BRIGHTON.—

In the last two decades great extensions have been made to this hospital, but circumstances have till now prevented a corresponding extension in the accommodation for nurses. The original nurses' home could provide lodging for 43 of the nursing staff; now the staff numbers 78. As far back as 1907 the matter became pressing, the provision of accommodation in rented houses involving discomfort to the staff in addition to being a costly procedure. In 1912 the governors had plans prepared for an extension of the original nurses' home; a special appeal for funds was made early in 1914, but the project had then to be abandoned, and it remained in abeyance until 1919, when a legacy of over £20,000 was left by Miss Walker. This legacy was treated as capital, and at a cost of £23,000 accommodation has now been found for 88 nurses. The extension to the nurses' home was opened by Princess Mary on Nov. 22nd, on the occasion of her first visit to Brighton in an official capacity. She also presented badges to nurses.

ON THE NON-LACTOSE-FERMENTING BACILLI

IN THE STOOLS OF HEALTHY AND UNHEALTHY
UGANDA NATIVES.

(Preliminary Observations.)

BY H. LYNDBURST DUKE, O.B.E., M.D.,
D.T.M. & H. CAMB.,

BACTERIOLOGIST, UGANDA PROTECTORATE.

THE establishment of a laboratory in a country where the bacteriological aspects of disease have hitherto received practically no attention entails a great deal of preliminary work in order to arrive at a proper appreciation of the limits of normal health for the community. In Uganda we have to deal with a mixed population of Europeans, Asiatics, and natives, many of whom are strangers in the land and only partially adapted to their environment. The native community includes members of many different tribes, accustomed to different dietetic and climatic conditions, and about whose nosology relatively little is known. In the Uganda Laboratory Report for 1920 a number of Arneeth counts were published, which were made with the object of revealing the limits within which variation is compatible with normal health in the various communities. From the little that has been done in this direction it is already clear that considerable care must be used in basing conclusions on this test. The investigations recorded below deal with the non-lactose-fermenting Gram-negative bacilli present in the fæces of healthy and abnormal inhabitants of Uganda—coloured and white. Attention has been confined to non-lactose fermenters, because most of the recognised pathogenic intestinal organisms belong to this category. The work of Douglas and Colebrook and others for the Medical Research Committee has afforded a model upon which the inquiry was roughly moulded, and the results obtained are presented in the briefest manner possible in view of the paucity of the material examined and the relative incompleteness of the examination.

The observations here recorded represent the work of one session terminated by my departure on leave. During my absence the majority of the strains described below will be maintained on agar, to be re-examined when occasion offers. The small number of serological tests applied up to date is explained by the press of routine and other work on the limited laboratory staff. No absorption tests were possible owing to the absence of a suitable centrifuge, but it is hoped to overcome this difficulty next session. It was necessary also to dispense with pathogenicity tests, as the supply of animals was inadequate. The organisms isolated have been grouped provisionally according to the principal sugar reactions. It is realised that the sugar fermentation reactions of an organism do not constitute an infallible means of differentiating species, but they supply a convenient expedient in inquiry of this nature. At present Uganda is fortunate in being relatively free from enteric disease, but the knowledge that must accrue from an investigation of this nature will prove of value should a demand for a prophylactic vaccine arise in the future. It will be interesting to see whether the serological races indicated in some of the groups below will survive the test of absorption. The recognition of serological affinity among strains giving the reactions of Morgan's bacillus No. 1, and isolated in Uganda and Kenya Colony respectively, may prove of some significance in view of Pirie's observations on this organism in Nairobi.¹

Technique.

A small portion of the stool was thoroughly emulsified in sterile distilled water in a test-tube, and after the emulsion had stood for some 15 minutes a loopful was spread on a MacConkey plate. Only one plate was used for each direct plating, and at times only half a plate. At the same time two or three loopfuls of the emulsion were inoculated into peptone-water tubes containing 0.4 and 0.6 c.cm. of a 1/10,000 solution of brilliant green per 10 c.cm. of medium.