to refrain from refilling, without the permission of the prescriber, any prescription which hitherto has been, or in the future may be, given by any of us.

(3.) That we withdraw our patronage from any and all apothecaries who shall wantonly violate our expressed wishes in this matter.

THE DISAPPEARANCE OF TUMORS.¹

BY THOMAS DWIGHT, M. D.

THE fact that tumors occasionally disappear, either spontaneously or in consequence of treatment, rests upon comparatively few authentic observations. The claims of quacks, however powerful with the popular mind, tend to make the profession, if anything, more skeptical than they would otherwise be of phenomena which, though rare, are sufficiently well attested. I regret that I am unable to report the following case in as much detail as could be desired. The notes that I took at the time are by no means as full as they would have been if I could have foreseen what unexpected developments were to occur.

The patient, in June, 1877, was a gentleman of about forty-eight, of a weak constitution, without definite occupation, who led a most unhealthy life, going to bed in the small hours, rising towards noon, and taking little exercise. I think it was nearly a year and a half before this, in the winter of 1875-76, though possibly in that of 1876-77, that he consulted me about a trouble in the rectum. This consisted of a sense of fullness and soreness. The patient stated that some years before (seven or eight, I believe) he had had hæmorrhoids, for which he had been treated by a well-known irregular practitioner of the day, who had applied caustics and effected a cure. It should be stated that my patient was of the kind who are easily deceived by quacks, and I am far from convinced that his trouble, of which he thought he was cured in this manner, ever really existed. On examination I found a somewhat hypertrophied fold of the mucous membrane projecting into the rectum. On withdrawing the finger the end of the fold could be brought into sight. It was evidently congested, but did not deserve to be called a hæmorrhoid. There was no occasion for any operation, and in view of the poor condition of the patient I should have been very unwilling to have undertaken one without imperative indications. I recommended an astringent wash and some simple treatment, under which the discomfort subsided.

In the spring and early summer of 1877 the patient was looking very badly, and complaining a great deal of pains and cramps in his legs. His walk was peculiar; descending stairs seemed especially difficult, and there appeared to be a want of coördination of movements.

Though constantly complaining, he was disinclined to submit to a thorough examination. Towards the end of this period pains returned in the rectum with great severity. He described it as feeling as if full of broken glass. There was a sympathetic difficulty in the bladder, which I thought did not empty itself perfectly.

On June 30th he was eating little, very much run down, weak and emaciated, and suffering from such

profuse night-sweats that his night-clothes were completely saturated two or three times a night. A somewhat superficial examination of the rectum revealed a hard swelling in the right wall. I have rarely seen a patient with cancer who presented the characteristic cachexia more markedly than did this patient at about this time.

July 2d. The rectum was thoroughly examined. The mucous membrane appeared perfectly healthy, but a hard tumor was found occupying the right side of the pelvis, extending from near the concavity of the sacrum to the arch of the pubes. It pressed against the rectum so as nearly to close it when empty, but the finger could be passed round the convexity of the tumor. Morphia suppositories of one fifth of a grain were given when required to relieve the severe pain. The patient for some time had been taking six grains of quinine daily.

On July 3d and 4th the patient seemed to be worse, and on the evening of the 4th Dr. R. M. Hodges saw him with me in consultation.

The tumor was as has been described; if anything, larger than when I examined it, two days before. It was decided that an operation was out of the question, and that the case was inevitably fatal.

It should be stated that there had been no chill nor febrile disturbance. The pulse was of fair strength and abnormally slow. The patient was very weak, however, and the night-sweats were very profuse. Oxide-of-zinc pills were given.

July 5th. I decided to try the effect of iodide of potash, beginning with ten-grain doses three times a day. The bowels moved two or three times during the day.

July 6th. Last night two suppositories were required. Sweats rather less, but pulse and strength weaker. The patient vomited after taking the zinc pill.

July 7th. Again two suppositories during the night. The zinc pill again rejected, and it was consequently stopped. The tumor can now be felt through the abdominal walls. Yesterday and to-day attempts at defecation. Yesterday something was passed which patient describes as a little whitish mucus; to-day nothing. Three suppositories in twenty-four hours is now the rule.

July 8th. Tumor more easily perceptible through abdomen. An enema of water brought away a rather loose discharge.

July 9th. Weaker. Unsuccessful attempt at a movement of the bowels.

July 10th. Had profuse night-sweats, but is pretty comfortable in the morning. Enema. Two or three fair discharges, one with a stain of blood. Morning pulse 48, evening, 75. Apparently losing ground.

July 11th. Looks worse than ever. Three operations during day.

On July 12th a change was noticed. There was less pain. Only two suppositories were used in twenty-four hours, and the patient seemed better.

July 13th. He was evidently better, and had a fair appetite.

July 14th. He was still improving, and the iodide was increased to fifteen grains thrice a day. He still took about three suppositories daily.

July 17th. Still improving. No morphia or bromide since 15th.

He continued improving slowly till July 25th, which was an intensely hot day. He suffered very much from the heat, and had two fits of vomiting.

¹ Read before the Boston Society for Medical Observation, November 1, 1880.

There are no more entries in my notes for a considerable time. It was evident that the patient was improving, but I could not believe that there was any real change of condition. I made no examination of the rectum, not wishing to torment him unnecessarily. I thought it right to let him enjoy his respite, feeling that it must be short, and that an examination could only bring distressing knowledge. I advised him to increase the iodide to twenty grains, and saw him but rarely. To my surprise, no relapse occurred. The patient continued to gain, and though feeble and much of an invalid looked, at the end of September, as well or better than he did before the acute attack. He discontinued the iodide, I believe, early in August.

On September 21st I requested permission to make an examination. The rectum seemed perfectly healthy. It was capacious. There was no tumor and no tenderness.

The patient during the next winter seemed better than usual, but during the summer of 1878 his health declined, and in the autumn it was evident that he was suffering from some grave abdominal trouble. The liver was greatly enlarged; he suffered much from nausea and vomiting, became deeply jaundiced, and died November 29th. The symptoms were quite obscure, and I was inclined to believe that there was malignant disease. A few weeks before his death he again complained of a fullness in the rectum, and on examination I thought I found some undue hardness on the right side. It was nothing, however, of much importance, and my attention was not again called to the rectum in the course of the disease.

The autopsy, made by Dr. E. G. Cutler, revealed that the cause of death was inflammation of the liver. No signs of malignant or venereal disease were discovered. The rectum was carefully examined. The mucous membrane was healthy, and no cicatrix or hint of any opening in it could be found. At each side of the rectum below the lateral reflection of the peritonæum there was a mass of what appeared to be dense hypertrophied connective tissue. This view was confirmed by Dr. Cutler's microscopic examination. The mass on the right side was the larger, and perhaps two thirds of the size of a hen's egg. There were no other signs whatever of an old abscess or of a tumor.

As I have already stated, the record is wanting in many points of detail, but the central fact is beyond question, supported as it is by Dr. Hodges' observation as well as my own, that there was a large hard tumor in the pelvis; and it is also certain that it disappeared. As to its position, though I could not assert that it did not involve the walls of the gut, I believe it to have been external to it. As to its nature, without going into details of the histology of tumors, it may have been one of three things: an abscess, a new growth, or an engorgement of the areolar tissue with blood and cellular elements either from the blood or the connective tissue. The peculiar pain, as of broken glass, is, I think, very characteristic of suppuration, but I am inclined to exclude abscess on account of the absence of fever, which would have been expected when the patient was at his worst; on the contrary, he had a low pulse, was clammy, and almost collapsed. If the abscess had burst, it is hardly credible that its contents could have been completely evacuated at once, so as to have been mistaken for a fecal evacuation. There would have been at least a little discharge for some time that could hardly have escaped notice;

probably, also, the bowel after death would have presented the lesion. It is true that even a large abscess may be absorbed. The New York *Medical Record* of March 13, 1880, contains an account of a curious case mentioned by Dr. Sands while presiding at a meeting of the New York Surgical Society. The case was that of Dr. Hubbel, who left directions that his body should be dissected. "Several years before his death he came to Dr. Sands complaining of a swelling in the left loin, and there was no difficulty in detecting a large fluctuating tumor over the region of the kidney, and extending down to the crest of the ilium. Dr. Sands was uncertain whether the fluctuating swelling had any connection with the urinary passages. Tapping was mentioned, but Dr. Hubbel being somewhat advanced in years, and knowing the possible mischief that might accrue, avoided the operation, and passed from under observation for nearly a year, and at the end of that time he reported himself, saying that he was well. While under observation the abscess extended, and evidence was obtained of matter burrowing down upon the buttock, three or four inches below the crest of the ilium; and it also extended upward to the free border of the ribs, but at the time he reported himself it had entirely disappeared. He subsequently died of Bright's disease. Dr. Sands examined the body very carefully, dissected the left lumbar region, but discovered no abscess. A great deal of connective tissue was found, in which it was difficult to recognize the remains of the kidney. The ureter was imbedded in this mass, which infiltrated the loin, and was impervious down to its junction with the bladder."

Turning from the theory of abscess to that of tumors, the question of syphilis naturally presents itself. I excluded it, because I never recognized any symptom of it during life or after death.

There seems to be little doubt that tumors of various kinds do, though very rarely, disappear without surgical interference. I was induced in this case to give the iodide of potash by seeing a passage from Holmes's Surgery, quoted in Bryant's, in which it is stated that Dr. Esmarch, of Kiel, told Paget that he had seen cases of recurrent fibroid tumor cured, and not again returning, in patients who took large doses of this drug for several weeks. Many of my hearers very probably remember a remarkable case of so-called Inflammatory Fungoid Neoplasm, reported by Dr. Duhring, of Philadelphia. This case appeared in the *Archives of Dermatology* for January, 1879, and a supplement to the case a year later in the same journal. In brief, a woman, otherwise healthy, for some two and a half years before her death had a great number of tumors, chiefly in the chorion or subcutaneous tissue, which presented the most surprising variations of size, sometimes entirely disappearing. The microscope showed them to consist essentially of a hypertrophy of the fibrous elements of the chorion and a varying amount of granular and other cells.

Dr. Coats exhibited for Dr. Gairdner specimens from a similar case before the London Pathological Society in April, 1879. The tumors were found also in the connective tissue at various places inside the abdomen. The growths were held to be lymphadenomatous. Sir James Paget made some remarks in the discussion which deserve to be reproduced. He said "the report of such a case was useful, as likely to help in the explanation of those rare instances in which tumors diagnosed to be cancerous had disappeared after a

time. He suspected that there was a greater number of such cases on record than might be imagined, and the collection of them would be an interesting and important undertaking. Three cases of the disappearance of tumors in this way were known to himself. One was in the person of a young man who had suffered for two or three years from what appeared to be ordinary lymphadenomatous growths, there being clusters of enlarged glands in the neck, axilla, and groins. The patient had also paraplegia, a symptom he had found in another case of lymphadenoma. Within a week these tumors all suddenly disappeared, but the patient then began to suffer from dyspnea, and soon afterwards died, no autopsy being allowed. Another case, mentioned in his lectures at the College of Surgeons, was regarded as one of multiple medullary cancer (what would now be called small-celled sarcoma), and the microscope corroborated this diagnosis. The growths occurred on the neck and axilla. There was a very large mass over one deltoid, which suppurated and sloughed, during which process nearly all the other growths disappeared. The man recovered, and enjoyed good health for some months, but the growth afterwards recurred and caused death. The third case was one which he had diagnosed as medullary cancer of an undescended testis. There was a tumor as large as two fists, and he had prescribed liquor potassæ and iodide of potassium, under which treatment the mass soon entirely disappeared. In eight or ten weeks, however, it recurred, but disappeared again under the same treatment. This also happened a third time, but having recurred a fourth time it was no longer amenable to treatment, and the patient died. The microscope confirmed his original diagnosis as to the nature of the growth." Other gentlemen mentioned somewhat similar cases.

An interesting paper on the sudden disappearance of tumors, by Dr. Fischer, of Breslau, is to be found in the *Deutsche Zeitschrift für Chirurgie*, vol. xii., 1879. Dr. Fischer calls attention to the fact that in certain very prostrating diseases tumors of some kinds suffer a great reduction in size, and among these are sarcoma, adenoma, and swellings of lymphatic glands. On the other hand, carcinoma, lipoma, fibroma, and some other tumors are not affected under like circumstances. The cases he reports are chiefly ones of enlarged lymphatic glands and of enlarged thyroids. Some of the glandular tumors were greatly affected by the removal of other tumors. One of the enlarged thyroids returned to its proper size during a light attack of scarlatina.

To return to the case I have reported, if I am justified in excluding abscess and syphilis, what remains? Many, I suppose, would object to the tumor being classed among the sarcomata on the ground that if it had belonged to that group it would not have disappeared. This, however, is begging the question. The engorgement of the liver had no doubt already begun before the time of the tumor. The direct venous connection of the liver with the rectum favors the production of hæmorrhoids when there is an obstacle to the circulation, but it is perhaps not impossible that through some peculiarity of the vascular distribution this effect took place in this instance around the rectum instead of in its walls. This explanation, I admit, is barely satisfactory to myself, but the masses of hypertrophied connective tissue beside the rectum suggest the sponge-like framework of an inflammatory growth, into the meshes of

which cellular elements might make their way, causing a great expansion, which would disappear on their withdrawal. Is this process essentially different from the development of a sarcoma? Billroth remarks that if any one wishes to define the inflammatory new formations in their different stages as types of sarcoma he is ready to agree to it. Whether the disappearance of the tumor was due to the use of the iodide of potassium is of course a matter of speculation; the tumors in Dr. Duhring's case disappeared without it. Still, under similar circumstances, I should certainly try it again.

MASSACHUSETTS LAWS REGARDING ANATOMICAL SCIENCE.

BY E. M. HARTWELL, M. A.,

Fellow of Johns Hopkins University.

It is not yet one hundred years since Dr. John Warren delivered the first course of public anatomical lectures ever given in Massachusetts, in compliance with a vote of invitation passed by the Boston Medical Society, November 3, 1781. It is scarcely fifty years since the Massachusetts Medical Society began to agitate the question of legalizing the study of anatomy. The Harvard Medical School, in the ninety-eight years of its history, has had but three professors of anatomy, namely, Dr. John Warren, professor of anatomy and surgery from 1782 till 1815, when he died; Dr. John C. Warren, professor of anatomy and surgery from 1815 to 1847, when he resigned; and Dr. Oliver Wendell Holmes, professor of anatomy, who, like the elder Warren, has held his chair thirty-three years.

Dr. John Warren's son and successor, Dr. John C. Warren, was three years old in 1781, the year the Massachusetts Medical Society was incorporated. Fifty years later, as one of the most prominent members of that society, February 2, 1831, he lectured before the members of the Massachusetts legislature, in the representatives' chamber, on the Study of Anatomy, in accordance with a vote of the house of representatives, passed January 29, 1831. At the time of this lecture the anatomy bill, which became a law on the 28th of that month, was still pending.

No better testimony concerning the obstacles which beset the pursuit of anatomical science during those fifty years can be given than is found in the Biographical Notes of Dr. John C. Warren, from which we quote: "No occurrences in the course of my life have given me more trouble and anxiety than the procuring of subjects for dissection. My father began to dissect early in the Revolutionary War. He obtained the office of army surgeon when the Revolution broke out, and was able to procure a multitude of subjects from having access to the bodies of soldiers who had died without relations. In consequence of these opportunities he began to lecture on anatomy in 1781. After the peace there was great difficulty in getting subjects. Bodies of executed criminals were occasionally procured, and sometimes a pauper subject was obtained, averaging not more than two a year. While in college I began the business of getting subjects in 1796. Having understood that a man without relations was to be buried in the North Burying-Ground, I formed a party. . . . When my father came up in the morning to lecture, and found that I had been engaged in this scrape, he was very much alarmed, but