treatment, when the clinical features of the case warrant us in declaring that such a chest is prepared for the bacillus, through imperfect function, or through exudations resulting therefrom. In this respect we have been leaning too much upon the evidence of the microscope, and too little upon equally as positive clinical evidence, of far more valuable diagnostic importance, viz.: imperfect function, as shown by the absence or muffling of the vesicular murmur. If we accept the clinical evidence, the fight with the bacillus can be easily carried to a successful issue outside the walls. But if we find that the bacillus has already taken possession, the fight will be longer and harder, and accompanied with many dangers, although it may yet be carried to a successful result if we follow the teachings of Nature, so far as we know.

It is impossible to describe in words the sound of the vesicular murmur. Every individual physician must acquire a knowledge of it through his own independent discovery, by carefully listening to many healthy chests, until his own senses are forcefully impressed with its character and language. It is a gentle breezy sound, exactly like nothing else in the world, but when thoroughly appreciated one of the most delightful sounds the ear can listen to, for it means absolute health; its absence, absolute disease sooner or later, unless restored by appropriate means intelligently directed.

The methods which I have used for many years were described in an article entitled "Personal Observations in Pulmonary Phthisis," published in the *Medical Record*, March 14, 1898, and in *THE JOURNAL A. M. A.*, Feb. 5, 1898. In the absence of more effective means, I still hold to the same methods, for the reasons there given.

The principle of the action of the pneumatic cabinet is correct, and indispensable in many cases, and should be applied where it is possible to obtain it, for it is a very difficult thing to wake up the almost paralytic chest muscles to perform their natural function, and is only obtained by persistent and continuous effort.

I have observed a very great lack of appreciation among medical men of the importance of this fundamental symptom of the vesicular murmur, and many lives are irretrievably lost by neglect of the methods required for its restoration, which otherwise might have been saved by prompt and early mechanical expansion.

ON THE RELATION BETWEEN THE VARIETY OF MICRO-ORGANISMS AND THE COM- POSITION OF STONE IN CALCULOUS PYELONEPHRITIS.

THOMAS R. BROWN, M.D.

BALTIMORE, MD.

No subject in renal diseases has been more discussed than calculus, especially as regards the etiologic factors involved in the production of the stone. Many theories have been held regarding the origin of stone in nephrolithiasis, but of these most have been finally abandoned from insufficient evidence; thus the idea that telluric conditions play an important rôle in this connection has been practically abandoned by all, as also the belief that gout was the most important etiologic factor in the development of such a condition, while Senator, Von Noorden and numerous other observers have definitely shown that the method of living and the variety of diet play, speaking generally, but a small part.

Our ideas regarding the formation of uric acid in

the body have undergone a marked change, through the work of Horbaczewski and Kossel, who showed that in a large number of cases, at least, the uric acid output was directly dependent on the amount of nucleins destroyed in the various vital and metabolic processes of the body.

That, however, simply an increase in the uric acid output is in itself not sufficient to bring about calculus formation is shown in the first place by the fact that in childhood, when the uric acid output is extremely high, renal calculi are extremely uncommon, while in various conditions, pathologic and otherwise, which are associated with an extremely high output of uric acid, such as splenomyelogenous leukemia, calculus formation is rarely met with. Also in cases of experimental phosphaturia, we practically never meet with phosphatic stones.

It will thus be seen that some other factor besides the formation of excessively large amounts of various of the chemical constituents of the urine is necessary for the formation of the different varieties of stone, at least in the great majority of cases. This conclusion and also the discovery of an albuminoid center or nucleus in the case of a number of stones, has led some observers to believe that the stone is formed by deposition of some of the urinary constituents about a proteid mass made up of epithelial cells, pus cells, red blood cells, etc. Comparatively recently, however, bacteria have been assigned an important rôle in this connection, due in the first place to the frequent association of calculus with certain forms of renal infection, and in the second place to the demonstration of bacteria in the center of a number of renal calculi. Thus, according to Harris, who has recently reviewed the subject, various micro-organisms have been found in the center of several renal calculi, by Galippe, in one case by Begoyne, and in an especially interesting case by Lenander, who demonstrated zoögleal masses of the colon bacillus in the center of a calculus composed of calcium oxalate and urates. Harris himself also reports three cases in which cocci or bacilli or both were found in the center of uric acid calculi; in these cases, however, cultures of the bacteria were not made, but their presence was demonstrated only by microscopic examination.

In all the cases which I wish to report, careful bacteriologic examinations were made of the urine, while the chemical composition of the stone was definitely determined in every case by the qualitative methods of chemical analysis usually in vogue. These cases, seven in number, were all those in which infection of the renal pelvis or the substance of the kidney was associated with the presence of stone, and in all cases except one the symptoms definitely pointed to the condition being unilateral, and this was subsequently verified by the bacteriologic, chemic and microscopic examination of the urine on either side. The means of making this examination was as follows: Through the cystoscope a sterile ureteral catheter with a sterile rubber cuff on its distal end was introduced into the ureter and, after the urine had been allowed to flow for a short space of time, the rubber cuff was withdrawn and a certain amount of the urine collected in a sterile test-tube from which, subsequently, the bacteriologic, chemic and microscopic studies were made. This method absolutely eliminates any chance of contamination from the bladder or from external sources if carried out with rigorous care. From the urine thus obtained, cultures were made upon agar plates, and from the colonies that grew thereon the species of micro-organism was determined.

by the usual methods. As stated before, ureteral catheterization was practiced in all cases on the side which gave no symptom, and in all cases except one the urine from that side was absolutely normal, showing that the calculous pyelonephritis was unilateral while in the case of the infected side the urine always contained large numbers of pus cells and a smaller number of red blood and epithelial cells and a considerable amount of albumin, while the reaction of the urine from that side depended entirely upon the variety of micro-organism met with, being acid in the one case of calculous pyelonephritis due to the colon bacillus met with, and alkaline in five cases due to various micro-organisms which possess in a marked degree the ability to decompose urea. These seven cases are therefore subdivided into: 1, those where the urine from the affected kidney is alkaline and 2, where the urine is acid.

WHERE THE URINE IS ALKALINE.

Five cases have been met with and studied bacteriologically and chemically, in which the urine was alkaline, while in one other case, although there had been marked urinary alkalinity at the beginning of the trouble the infection had subsequently died out spontaneously and the urine, when examined by us, was acid and sterile. The history of the development of the infection in these cases is of extreme importance; in Case 1 the condition came on insidiously, without any apparent cause, and was subsequently followed by several typical attacks of renal colic. In Case 2 the onset had also been insidious, while the only symptoms present were dull pain in the back and right side for several years, and loss of strength and weight. In Case 3, where the condition had lasted for over fifteen years, the etiologic factors had been quite obscure, the condition developing after a protracted anemia, the exact cause of which could not be definitely determined although it was probably chlorotic in nature. The symptoms in this case had been pyuria and pain, first on the left side, subsequently on the right side also, and in the former case they became so great as to necessitate the removal of the kidney which was converted into a large pyelonephrotic sac in whose center a calculus was found. In Case 4, five months after a difficult instrumental labor, the patient complained of severe pain in the right renal region, which had been constantly present ever since (3½ years) associated with occasional attacks of sharp pain. In Case 5 the pyelitis was secondary to a cystitis occurring 2½ years after the development of the latter condition. The symptoms were almost constant severe pain in the right renal region, associated with a high grade of pyuria; while in the sixth case, where the infection had spontaneously died out, the patient had developed an acute ammoniacal cystitis after severe instrumental labor thirty years ago, followed shortly by symptoms of an infection of the left kidney; the urine remained alkaline or ammoniacal for many years, but for the past twenty years has been almost constantly acid, although it still at times contains enormous numbers of pus cells as if from some large sac which occasionally empties itself. The bacteria found in these cases were respectively as follows: Case 1, *B. proteus vulgaris*; Case 2, *B. proteus vulgaris*; Case 3, a white staphylococcus which rapidly decomposed urea and liquefied gelatin but slowly; in Case 4, the same white staphylococcus just mentioned; in Case 5, the *B. proteus vulgaris*; while in Case 6, as stated before, the urine was sterile.

In all cases except one, nephrectomy was performed,

the stone removed and the renal pelvis irrigated and drained; in the one exception mentioned several calculi were voided with the urine, so that it was possible in all cases to make a careful chemical examination of the stone. The chemical composition of the stone in all these cases was the same, the calculus being made up of the phosphates and carbonates of calcium and magnesium, although the proportion of the various salts differed in the different cases. In one case, that of the calculous pyelonephritis due to the infection of the *B. proteus vulgaris*, a culture was made under the most careful aseptic precautions from the center of the stone, and the *B. proteus vulgaris* was obtained therefrom in pure culture; in another case microscopic examination of the center demonstrated staphylococci—in this case the renal infection was due to the white staphylococcus previously mentioned—while in the third case, that in which the infection began thirty years ago and in which the urine had apparently been sterile for upwards of twenty years, no bacterial center was able to be demonstrated. In the other three cases the stones were not examined to see what was the exact condition present in their respective centers.

In one case the diagnosis was made in a rather unique fashion. Through the ureteral catheter there was injected into the renal pelvis of the affected side 15 c.c. of a .2 per cent. solution of hydrochloric acid, which was allowed to remain in the pelvis between ten and fifteen minutes; it was then analyzed chemically, when the marked amount of calcium and magnesium phosphate and carbonate present showed that the pelvis contained a calculus of that composition.

The condition in all these cases was, as far as could be judged from the careful study of the symptoms and the urine, one of infection of the kidney subsequently followed, sooner or later, by stone formation. A consideration of the findings in certain of the cases warrants us in the belief that the nucleus of the stone was a zoögleal mass made up mostly of the micro-organisms causing the infection, while the constant alkalinity of the urine caused by the urea-decomposing power of the micro-organism furnish the salts necessary for formation of the stone.

The extreme frequency of calculus in infections of the kidney brought about by such urea-decomposing bacteria is shown by the fact that, of all the cases of renal infection due to this class of bacteria, stone was found in all the cases but one which was an acute pyelonephritis of but two weeks' duration where, obviously, sufficient time had not elapsed for the deposition of the salts necessary to form the stone.

WHERE THE URINE IS ACID.

Only one case was met with in which stone was associated with a pyelonephritis due to a micro-organism which retained the normal acidity of the urine. This case was one of infection arising immediately after the formation of a uretero-vaginal fistula, which in turn had been produced by the trauma of a very severe, protracted instrumental labor.

The infection was due to the *B. coli communis*, and had been present for one year; the urine from this kidney was discharged entirely through the fistula, contained pus, blood and epithelial cells and considerable albumin and was always very markedly acid; from this urine a pure culture of the colon bacillus in considerable number was obtained. There had been absolutely no symptoms of stone, but, following upon a careful vaginal and rectal examination, a small calculus was passed

through the fistula, the chemical analysis of which was a mixture of uric acid and urates, the former largely in excess. The consideration of this case makes it highly probable that the stone was formed definitely because of the urinary infection.

Albarran has divided the cases of renal calculi into primary stones where the kidney is not the seat of infection, and secondary stones where bacteria are found in the renal pelvis; according to this classification our cases fall into the second group, although recently many have ascribed to bacteria the formation of stones of the first group, believing that a renal bacteriuria furnishes the focus about which the stones are built.

My series of cases, therefore, besides showing the definite relationship existing between the variety of micro-organism causing the infection and the chemical composition of the calculus, also adds more cases to the ever-increasing list in which to the presence of bacteria is to be ascribed the formation of stone; this is not only so in the case of renal calculi, but the causal relationship between bacteria and stone has been shown also in cholelithiasis, salivary, lacrimal and pancreatic stones, in rhinoliths and otoliths and various stones found in pathologic conditions of the respiratory tracts.

1033 Cathedral Street.

COMPLETE INGUINAL EXTRAPERITONEAL HERNIA OF THE BLADDER; RECOVERY.

J. F. BALDWIN, A.M., M.D.

Surgeon to Grant Hospital; Fellow of the American Association of
Obstetricians and Gynecologists, Etc.
COLUMBUS, OHIO.

Mr. D. H., West Mansfield, Ohio, aged 51, 5 ft. 6 inches in height and weighing 245 lbs., consulted me May 26, 1900. He stated that he had had an inguinal rupture for eight years, which he had been unable to retain with any form of truss. Ordinarily he had been able to reduce his hernia, but it had now been irreducible for forty-eight hours. The day before he consulted his local physician, Dr. Wanser, who administered an anesthetic but was unable to affect a reduction. He stated that he had noticed many times that he was unable to empty his bladder completely without lifting up the tumor.

The tumor was about the size of two fists and extended to the bottom of the scrotum. It was somewhat oval in outline and conveyed a sensation of fluctuation. It was tender and entirely irreducible. Although he was experiencing some pain, there was no evidence of an intestinal disturbance. I therefore explained to him that I thought the main mass of the tumor was omental in character, but that I should not be surprised to find his bladder involved. I based this statement of the involvement of his bladder on the fact that he had to lift up the tumor in order to completely empty that viscus. I advised immediate operation and assent was given. He was admitted to the hospital and the operation made an hour or two later.

I was assisted in it by my regular assistants, Drs. Shepard and Chapman, Dr. Willey, of Baltimore, Md., being present. The usual incision was made over the tumor, but this had to be somewhat longer than usual, owing to the great amount of fat which was present, an incision about 2 inches in depth being required to reach the external fascia. The tissues were carefully divided until what corresponded to a hernial sac was opened. This disclosed a fleshy mass adherent throughout. The adhesions, however, were easily separated and the mass brought up into view. The fluctuation which I had noticed and which I supposed was due to free serum in the hernial sac, such as is usually met with in cases of strangulation, was now found to be inside the extruded mass. Drawing the tumor down the finger was passed up to the neck when it was found that the extrusion had taken place apparently as in a direct inguinal hernia. The ring being enlarged somewhat, and the finger passed through, the diagnosis was arrived at very

promptly that the projecting mass was practically the entire bladder, since all that was left inside the pelvis was merely the neck. The bladder walls contained a thick deposit of fat so that the entire thickness seemed to be about one inch.

With a good deal of difficulty a pelvic receptacle was made by separating the peritoneum, and the bladder replaced. The replacement was only accomplished with the utmost difficulty, owing to the large amount of adipose tissue present. The patient was placed in the Trendelenburg position and so maintained for a considerable length of time before complete reduction was accomplished. The opening was then carefully closed in layers with kangaroo tendon. The hernial sac, which consisted merely of condensed connective tissue instead of peritoneum, was trimmed up and the fat brought together by plain catgut in two layers, with a subcuticular stitch for the skin. The wound was then sealed with collodion and a spika bandage applied.

Recovery was immediate and in every respect satisfactory. The patient reported at my office four months later, when everything was found in normal condition, the tissues being apparently perfectly strong and firm.

I have been able to find in the literature at my disposal but very little bearing on this most unusual form of hernia. In the *Philadelphia Medical Journal*, of June 30, 1900, there is a synopsis of an article by E. Martin, in the *Deutsche Zeitschrift für Chirurgie*, for February, 1900. In this Martin reports a case briefly as follows: A man, 51 years of age, had had a hernia the size of a small walnut in the right groin for some eight years past, present only at intervals and always easily reducible. Twenty-four hours previously the hernia had reappeared, this time larger and irreducible: the pain and acute symptoms led to the diagnosis of incarcerated omental or intestinal hernia. At the operation it was found that the hernia consisted of a portion of the bladder which had been protruded extraperitoneally, and which was easily replaced after the constriction had been relieved. The patient recovered. Martin, in this article, has collected from the pages of literature nine cases of strangulated hernia in which, as in his own case, the bladder alone was involved. In only one of these cases was a hernia of the bladder suspected before the operation was started, and it is plainly to be inferred from this report that in most of these cases, as well as in his own, the bladder was wounded before the mistake in diagnosis was discovered.

According to Martin's article, therefore, my own case makes the eleventh to be recorded. In my case, fortunately, the involvement of the bladder was suspected before beginning the operation, and hence that organ was looked after.

Cases in which the bladder has been involved in a hernia together with omentum and intestine are not so uncommon, but that the bladder alone should become extruded is certainly quite surprising, and for that reason such cases are well worth recording. In the case reported the thick deposit of fat in the bladder wall caused it to so closely resemble omentum that the surgeon, unless on his guard, would more than likely have incised it.

Taxing Poisons.—According to the *New York Times*, in an editorial with this caption. Sir Michael Hicks-Beach, Chancellor of the Exchequer, England, has received a suggestion from the food reformers that a tax be laid on poisons found in food and drinks. Arsenic in beer appears to be the main inspiration of the proposition. Parliament has certainly deemed it of enough importance to create a government commission, much to the consternation of the brewers. The value of the public laboratories is pointed out from the fact that the presence of arsenic in beer is a very recent discovery. The combination of a tax and a benefit is a rare one, as maintained by the editorial in question, but the argument is advanced that not much aid would be given to the replenishment of the imperial coffers since what it was sought to tax would disappear. "To smile but put the question by," virtually concludes the article, is the immemorial habit of a chancellor. Reforms, however, it may be added, are seldom advanced when criticisms regarding methods become too rampant.