tery; but coming on in the advanced stage of convalescence from disease, and yet depending on the morbid condition alone, is what during my professional life, which is now advanced many degrees beyond its meridian, I have *never* seen.

Dec. 20th, 1855.

CASE OF SCROFULOUS DISEASE.

[Communicated for the Boston Medical and Surgical Journal.]

MESSRS. EDITORS,—I transmit the following statement of a case which came under my charge a few weeks since (it being quite novel to me, and may be so to some of the readers of the Journal), for the reason that it showed a remarkable extent and variety of development of this formidable disease, in so young a child, and also because life was so prolonged while inhabiting such a diseased and corrupted “tenement of clay.”

The history of the child as given me by the parents, is as follows: Some weeks after birth it was attacked by a cough, but, as no other urgent symptom was attending to demand *rational* treatment to preserve life, he was treated by their family physician for “a cold”—a malady which has a great account to render at the last summing up, if guilty of half that is ascribed to it in these parts. In a few weeks the cough subsided, and the child was apparently well, though not strong. The parents remarked to me that it was “a very bright, sweet child,” which was symptomatic of that precociousness attending this fatal disease.

The physical development was pretty good up to the age of two years, though not keeping pace with the mental. It continued tolerably healthy, nothing appearing to indicate disease excepting that once or twice there were swellings in the neck, but these disappeared in a few days.

I should have mentioned that the boy was 2 years and 9 months old when I first saw him, and he was under my care three months before he died.

When two years of age, the right side of the scrotum was observed to be swollen, and the “family physician” was again consulted. He diagnosed a hydrocele, and ordered a wash of *acetat, plumbi*, although the swelling was, in the commencement, uniform, hard, and attended with some pain. It was treated on the “lead principle” for nearly six months, when, as the physician thought, it had attained sufficient size to warrant evacuation; so he proceeded to perform the operation of “*dry tapping*.” After this, the child had no treatment for three months, the parents assigning as a reason that “the family physician had not been there since.”

When called to see the child, I found it presenting the general scrofulous diathesis, having a slight cough, capricious appetite, torpid bowels, enlarged abdomen. It was considerably emaciated and exceedingly peevish. The tongue and pulse were pretty good, and

there was no pain except during the erect position. The scrotal enlargement was now seven inches in length and ten and three quarters in circumference, nearly uniform, though smallest at its inferior portion. The enlargement was so great, that at first sight there appeared to be no penis. At the anterior superior portion there was something very much resembling the navel, though less in size, which was found to be the prepuce, with no apparent passage through which the urine was voided. Upon examination of the tumor, I found the left testicle of its normal size, at the superior part, but the right one was not to be found; or, in other words, the tumor was *all* testicle. There was a glandular enlargement in the right groin, of the size of a hen's egg, bluish in color, and a small one in the left.

Diagnosis—Scrofula. *Prognosis*—very doubtful.

Treatment,—first, Lugol's solution, which was tolerated but for a few days. After this, I gave iod. pot., syr. sarsap. and iod. pot., and syr. iod. ferri, each in turn, but all were rejected after a short trial. Some local applications of iodine were used, but all to no avail.

The testicle remained about the same size, changing in appearance only at its inferior portion, where it became purple, and ulcerated slightly. The tumors in the groin enlarged rapidly—the right to the size of a pint cup, and the left to two thirds the size of the right. The skin presented that peculiar purplish or livid hue, showing the languor of its capillary circulation, and its proneness to be destroyed by gangrenous erosion. Four weeks before death, the skin over the right tumor ulcerated and considerable hemorrhage ensued. The left soon followed the same course. The right opened in two places, forming ulcers of one and a half inch in diameter, from which there appeared a growth much resembling fungus hæmatodes or encephaloid cancer. They discharged a bloody ichorous fluid. Before death, their surface seemed to have passed on to mortification. The abdomen increased in size rapidly, and the superficial veins were blue and large, like varicose vessels. Swellings appeared in the neck. The general symptoms remained about as they were when I first saw him. There was gradual emaciation, with some more pain.

Autopsy, twelve hours after death. Head not examined. Louis states that he never found but one case of scrofulous disease without finding tubercles in the lungs.* I could find none in the lungs, though there were extensive adhesions of the right lung; therefore I think this is another. Some effusion in the pericardium; tubercles in the liver, from the size of a nutmeg to that of a pea, somewhat softened and of curdy consistence. The mesentery contained a tumor of five or six pounds in weight, firmly adherent on its posterior surface, and to which the intestines were so firmly attached as

* This statement is not correct. Out of 358 subjects *above the age of fifteen*, in whom tubercles existed in various organs of the body, they were found in the lungs in every instance but one.—*Recherches sur la Phtisie*, 2nd edition, page 182.—EDITORS.

to prevent all motion. The mesenteric covering of the tumor was quite thick, and nearly natural in color. Upon opening it there was no organized structure found. It was composed of a semi-transparent substance, of about the consistence of jelly, with some traces of pus. The lacteals and absorbents were generally diseased. The disease in the testicle had so far advanced that there was scarcely organized structure enough to be recognized. It was light in color, soft, and composed partly of thin, light-colored pus, and partly of slightly softened tuberculous matter. There was in the inferior part, and also in a portion of the mesenteric tumor, a quantity of fluid resembling thin molasses. The tumor in the neck was of the ordinary kind of hard, scrofulous enlargement. Those parts of the glandular system not already passed, seemed to be rapidly passing on to malignant degeneration.

By this time perhaps you will be disposed to say I am troubled with *cacoethes scribendi*; but I wish to accomplish two things—to give a statement of the case, and to show that the time is very often suffered to pass by when suitable remedies *might* have arrested the disease.

H. P. STRONG, M.D.

Beloit, Wis., Nov., 1855.

CASE OF A FOREIGN BODY IN THE AIR-PASSAGES.

[Communicated for the Boston Medical and Surgical Journal.]

MESSRS. EDITORS,—The following case occurred about three weeks ago in my ride.

The patient, a girl about seven years old, inhaled a piece of hickory nut-shell, of very irregular and angular form. At the time, the accident occasioned but moderate annoyance. After a pretty severe paroxysm of coughing, she resumed her play, and remained well for a period of three days. On the fourth day, in the morning, she had a very severe paroxysm of coughing, during which suffocation was pending. At this the parents became alarmed, and I was sent for. I did not arrive until after this violent coughing had ceased, though the breathing was still quite embarrassed. These violent paroxysms did not return again until the final one, when the foreign body was expelled. From the time of my first visit, the whole case appeared very much like a sudden and severe development of bronchial inflammation. It was treated according to the usual mode in such cases (I should say vigorously), and there was a favorable abatement of all the symptoms; the patient, after the first week, would have been regarded as convalescent, were it not for the fact that there was a foreign body in the lungs.

After the decline of the inflammatory condition, the case seemed to be progressing so favorably that the idea of a foreign body yet existing in the lungs, was about discarded. But the reality of its presence was soon revealed by the appearance of swelling, with